Representability of pairing relation algebras depends on your ontology

Ágnes KURUCZ
Department of Computing
Imperial College, London
e-mail: kuag@doc.ic.ac.uk
István NÉMETI
Rényi Institute Budapest
e-mail: nemeti@renyi.hu *

June 26, 2000

Abstract

We consider classes of relation algebras expanded with new operations based on the formation of ordered pairs. Examples for such algebras are pairing (or projection) algebras of algebraic logic and fork algebras of computer science. It is proved in Sain–Németi [38] that there is no 'strong' representation theorem for all abstract pairing algebras in most set theories including ZFC as well as most non-well-founded set theories. Such a 'strong' representation theorem would state that every abstract pairing algebra is isomorphic to a set relation algebra having projection elements which are defined with the help of the real (set theoretic) pairing function. Here we show that, by choosing an appropriate (non-well-founded) set theory as our metatheory, pairing algebras and fork algebras admit such 'strong' representation theorems.

Keywords: relation algebra, projection elements, non-well-founded set theories

*Research supported by Hungarian National Foundation for Scientific Research grants T30314, F17452 and T23234. Á.Kurucz is also supported by UK EPSRC Grant GR/L85978.

1 Introduction

This paper is about representation (or axiomatization) problems of algebraic logic. In particular, we discuss the representation problem of classes of relation algebras expanded with extra operations based on the formation of ordered pairs. Our results are also related to completeness problems of finite variable fragments of first order logic. In the present paper we prove that in a particular non-well-founded set theory, a natural variant of these expanded relation algebras admits a so called 'strong representation theorem'. We will also see that for such a theorem we need this particular variant of our algebras as well as this particular choice of set theory. Namely: (i) the other variants which came up in the literature do not admit a strong representation theorem even in this set theory, and (ii) all known variants (including ours) of these algebras behave badly in many other set theories (non-well-founded as well as well-founded ones). This way this paper can also be considered as a contribution to the field of so called 'reverse mathematics' or 'inverse set theory', cf. e.g. Simpson [41], Andréka et al [4]. In this field one tries to find out what is the exact fragment of background set theory really needed to prove some theorems of certain mathematical disciplines, in our case theorems of algebraic logic.

The motivation for studying relation algebras expanded with extra operations based on forming ordered pairs comes from the following two areas (or 'traditions').

(1) In Tarskian algebraic logic, these kinds of algebras are called 'relation algebras with projection elements' or 'pairing relation algebras'. They have been studied beginning with the early 1940's, when Tarski observed that certain theories of first-order logic with countably many variables are 'algebraizable' with the help of relation algebras having 'projection elements' (Tarski [42], Tarski–Givant [43], Maddux [24]). The 'projection elements' were taken as distinguished constants of the similarity type in Németi [30], Andréka–Jónsson–Németi [3], Maddux [22], [23], Mikulás–Sain–Simon [28], Sain–Simon [39].

(2) Recently in computer science, these kinds of expanded relation algebras were found useful in various ways (e.g. in the theory of programming). Therefore they have been studied in the computer science literature extensively, cf. Veloso–Haeberer [44], [45], [17], Baum–Haeberer–Veloso [7], Berghammer et al [8], Frias et al [10], Baum et al [6], Frias–Gordillo [11]. The name 'fork algebra' was suggested in computer science for

a relatively comprehensive class, where relation algebras are expanded with a binary operation (based on pairing) called 'fork'.

Analogously to the situation with Boolean algebras, groups, semigroups, polyadic algebras etc., one can contemplate the idea of developing a representation theory for the above kinds of expanded relation algebras, too. Desirability of a representation theory also comes from logic (cf. section 5 for a discussion). By the above analogy, such a representation theory would consist of

- defining a *concrete* class of algebras, i.e. where the universe of an algebra completely determines the operations of the algebra (analogously to Boolean set algebras, transformation semigroups, polyadic set algebras, etc.); and
- an axiomatic (*abstract*) class of algebras, defined by finitely many first-order formulas (preferably equations) valid in the concrete class; and then
- a *strong representation theorem*[1] stating that every member of the abstract class is isomorphic to a member of the concrete class.

Both traditions (1) and (2) above mention as *standard examples* those expanded set relation algebras in which the extra operations (based on pairing) are based on the standard set theoretic definition of ordered pairs $\langle a, b\rangle \stackrel{\text{def}}{=} \{\{a\}, \{a, b\}\}$. These algebras are concrete in the above sense: given such an algebra $\mathfrak{A}$, to each operation g of $\mathfrak{A}$ (in particular, to the extra operation [fork or the projections] based on pairing) there is an explicit set theoretic definition of g using no other parameter than A (see sections 2 and 3 for definitions). Therefore it would be nice to find an axiomatic (abstract) class of, say, fork algebras such that their representation theory could be based

[1]There are cases when strong representation theorems are not available, and therefore so called *weak representation theorems* are used as a substitute. These representation theorems are called weak because the class K of algebras they use to represent an abstract class is not concrete in the sense defined above. In the case of fork algebraic weak representation theorems, all the relation algebraic operations are represented in a concrete way, but the extra operation 'fork' based on pairing remains abstract in the sense that it is represented only via some abstract class of groupoids, which cannot be recovered from the universes of the algebras in question, cf. Gyuris [15], [14], Sain–Németi [38], Baum et al [6]. Weak and strong representation theorems were discussed and carefully compared in detail (with examples from the literature) in e.g. Sain–Németi [38], Henkin–Monk–Tarski [18, pp.459–461, item 2.7.46].

on these standard algebras. This was indeed experimented with in the earlier papers of the computer science tradition, cf. e.g. Haeberer et al [7], [17], [44], [16]. Several variants of 'standard' algebras were introduced and studied in the literature, e.g. in Németi [30], Andréka–Jónsson–Németi [3], Veloso–Haeberer [45], Baum–Haeberer–Veloso [7], Maddux [22], [23], Sain–Németi [38]. For all of these variants, it was proved in Sain et al [38], [36], [35], [39], [28] that if one uses the usual Zermelo-Fraenkel set theory ZFC as his metatheory then this task cannot be solved for the following reason. The equational theories of these concrete classes are not finitely axiomatizable; further, these concrete classes are not axiomatizable by any set of first-order formulas. (There are even further negative results in the quoted papers, see section 4 for a summary.) These results hold if one works in our usual set theory ZFC as metatheory.

This might look discouraging, but if we are motivated by computer science applications of fork algebras, then the idea comes up that there might be alternative set theories to ZFC which are just as suitable for computer science as ZFC is. Indeed, Aczel [1] and Barwise–Etchemendy [5] discuss many so called *non-well-founded* set theories which are suitable for computer science, for theories of natural language as well as for parts of Philosophical Logic. These set theories are obtained from ZFC by dropping the Axiom of Foundation and adding some new axioms to replace it. It was also proved in Sain–Németi [38] that for many of these non-well-founded set theories the above quoted negative results carry over (for all variants of concrete classes of expanded relation algebras in the literature). In the present paper we discuss a particular non-well-founded set theory, denoted here as $ZFC^- + BAFA_0$, and we look at some particular concrete classes of expanded relation algebras (see Defs. 4 and 10 below), such that in our set theory our concrete classes are axiomatizable by finitely many equations. Thus we can provide an abstract class of, say, fork algebras, defined by finitely many equations such that every member of the class can be strongly represented. On the other hand, by [38], even our classes behave 'badly' in ZFC as well as in many other non-well-founded set theories. (Our concrete classes are exceptional in the sense that the *other* variants in the literature remain non-finitely axiomatizable even in $ZFC^- + BAFA_0$.) The results of this paper were announced in [32].

For completeness, we note that the word *ontology* in the title refers to the fact that the anti-foundation axiom used in this paper states the existence of certain kinds of sets. Therefore, at least in a sense, accepting or rejecting this axiom might depend on one's ontological commitment. (The basic question seems to be whether we want to have sets which are different from one another

but undistinguishable in the language of set theory.) In this connection, we note that we are under the impression that one can do 'ordinary mathematics' in $ZFC^- + BAFA_0$ reasonably comfortably[2].

The paper is structured as follows. In section 2 we introduce the particular non-well-founded set theory $ZFC^- + BAFA_0$ we use as our metatheory. We state, in $ZFC^- + BAFA_0$, our main technical lemma about the characterization of set theoretical projection functions. We introduce the various classes of expanded relation algebras and discuss their strong representation theory in $ZFC + BAFA_0$. We also prove that the approaches (1) and (2) above can be considered as equivalent ones in a very strong sense. Section 3 is devoted to some properties of our metatheory $ZFC + BAFA_0$, and to the proof of our main lemma. Section 4 gives a detailed summary of the negative results achieved in the above quoted papers of Sain et al. Section 5 discusses the connections with the so called 'finitization problem' of algebraic logic. We also discuss briefly some motivations coming from logic.

2 Representing relation algebras expanded with operations based on forming ordered pairs

In this section we develop a representation theory for some classes of expanded relation algebras. Since the validity of the statements may depend on the metatheory used, we carefully indicate, after the number of each theorem and claim, the meta (set)theory in which the theorem in question is proved. In particular, ZFC denotes Zermelo-Fraenkel Set Theory with the Axiom of Choice (which includes the Axiom of Foundation) and ZFC^- is ZFC without the Axiom of Foundation. We do not even assume the Collection Principle as an axiom of ZFC^- (which is used as an axiom of non-well-founded set theories in Aczel [1]). We use ZF and ZF^- similarly for set theories without the Axiom of Choice.

[2]As it is proved in section 3, if ZFC is consistent then the set theory ${BAFA_0}^+ \stackrel{\text{def}}{=} (ZFC^- + BAFA_0 + \text{collection principle})$ is consistent, too. Since our theorems are proved in $ZFC^- + BAFA_0$ as metatheory, they also remain true in ${BAFA_0}^+$. We mention this because, on the basis of [4], we conjecture that practically all theorems in e.g. the textbook Grätzer [13] are true in ${BAFA_0}^+$ (in place of ZFC). We believe that we have a proof that all the "isomorphism invariant" theorems of universal algebra and abstract algebra are equi-provable in ZFC and ${BAFA_0}^+$. We think that these equi-provability results can be pushed far beyond "isomorphism invariance", e.g. we checked that the known non-finitizability results of algebraic logic (cf. [29], [2]) also remain true in ${BAFA_0}^+$. E.g. the class of representable relation algebras remains non-finitely axiomatizable in ${BAFA_0}^+$.

Notation 1. $\langle x, y\rangle$ denotes the usual set theoretic *ordered pair of x and y*, i.e.

$$\langle x, y\rangle \stackrel{\text{def}}{=} \{\{x\}, \{x, y\}\}.$$

For any set U, $U \times U \stackrel{\text{def}}{=} \{\langle x, y\rangle : x, y \in U\}$. A *binary relation on U* is some subset of $U \times U$. For any binary relation E, dom E denotes the *domain of E* that is, dom $E \stackrel{\text{def}}{=} \{a : \langle a, b\rangle \in E \text{ for some } b\}$. We consider functions as special binary relations. A *partial function* on some set U is a function f with dom $f \subseteq U$. In case of some partial function f we will use notation $f(a) = b$ instead of writing $\langle a, b\rangle \in f$.

For any class K of algebras, $\mathbb{I}\,\mathsf{K}$, $\mathbb{S}\mathsf{K}$ and $\mathbb{P}\mathsf{K}$ denote, respectively, the classes of all isomorphic copies, of subalgebras, and of isomorphic copies of direct products of members of K. □

Definition 2. An algebraic structure $\mathfrak{A} = \langle A, +, \cdot, 1, 0, -, ;, ^\smile, 1'\rangle$ is a *relation algebra* if equations (R0)–(R7) below are valid in $\mathfrak{A}$ (cf. e.g. Henkin–Monk–Tarski [19], or Tarski–Givant [43]).

(R0) the Boolean algebraic equations for $+, \cdot, 1, 0$, and $-$
(R1) $(x\,;\,y)\,;\,z = x\,;\,(y\,;\,z)$
(R2) $(x + y)\,;\,z = (x\,;\,z) + (y\,;\,z)$
(R3) $x\,;\,1' = x$
(R4) $x^{\smile\smile} = x$
(R5) $(x + y)^\smile = x^\smile + y^\smile$
(R6) $(x\,;\,y)^\smile = y^\smile\,;\,x^\smile$
(R7) $x^\smile\,;\,[-(x\,;\,y)] \leq -y$

A relation algebra $\mathfrak{A}$ is called a *set* (or *proper*) *relation algebra* if A is a set of binary relations on some set U and the operations $+, \cdot, 1, 0, -, ;, ^\smile, 1'$ coincide with the usual set theoretic operations of union, intersection, $\bigcup A$, empty set, complementation relative to $\bigcup A$, relation composition, converse, and the identity relation on U, respectively. The set U is called the *base* of $\mathfrak{A}$. $\bigcup A$ (which is, being the biggest element of $\mathfrak{A}$, always an equivalence relation with domain U) is called the *unit*[3] of $\mathfrak{A}$. If A consists of all the subsets of $U \times U$ then $\mathfrak{A}$ is called a *full set relation algebra*. □

Recall that a representation theory usually consists of

- defining a concrete class of algebras, i.e. where the universe of an algebra determines every operation of the algebra in question;

[3]If $\mathfrak{A}$ is simple then the unit of $\mathfrak{A}$ can be taken to be $U \times U$, cf. [18], [19].

- an axiomatic (abstract) class of algebras, defined by finitely many first-order formulas (preferably equations) valid in the concrete class; and then
- a strong representation theorem stating that every member of the abstract class is isomorphic to a member of the concrete class.

In this sense, the class of relation algebras is an abstract class, while the class of set relation algebras is a concrete one. Thus we call a relation algebra *representable* if it is isomorphic to a set relation algebra.

An abstract subclass of relation algebras can be given as follows. A relation algebra $\mathfrak{A}$ is called a QRA (cf. Tarski–Givant [43])[4] iff the following (existential) formula is valid in $\mathfrak{A}$.

$$\exists p \exists q \ (p^{\smile} ; p \leq 1' \quad \wedge \quad q^{\smile} ; q \leq 1' \quad \wedge \quad p^{\smile} ; q = 1)$$

Such elements p, q are called *quasi-projections of* $\mathfrak{A}$.

Theorem 3. [Tarski][5] (in ZFC^-) *Every* QRA *is representable.*

Proof. The proof can be found e.g. in Tarski–Givant [43] or in Maddux [24], see also Simon [40]. Those proofs are done in ZFC, but it can be checked that they do not use the Axiom of Foundation. We note that for *countable* QRA's, the Axiom of Choice is not needed in the proof either. □

This theorem is a strong representation theorem in the above sense, since all the operations are represented in a concrete way. However, one can expand the similarity type with the constant symbols p and q, as it is done in Németi [30], Andréka–Jónsson–Németi [3], Maddux [24], [22], [23], Mikulás–Sain–Simon [28], Sain–Simon [39]. That is, let

$$\mathsf{K} \stackrel{\text{def}}{=} \{\langle \mathfrak{A}, p, q \rangle \ : \ \mathfrak{A} \text{ is a } \mathsf{QRA}, \text{ and } p, q \text{ are quasi-projections of } \mathfrak{A}\}.$$

Then Tarski's theorem does not imply that every member of K is strongly representable. Though, in the representing set relation algebras, quasi-projections p and q are partial functions with $p^{\smile} ; q = 1$, their behaviour cannot be 'found out of' the universe of the algebra, they can be any pair of partial functions on the base of the algebra in question which satisfy

[4]Throughout, we use notation like QRA in two different sense. First, a QRA is an algebra defined above. Second, QRA also denotes the class of all QRA's.

[5]In 1943–44, a theory was built around this result and was presented at Tarski's Relation Algebra Seminar in 1945, Univ. of California at Berkeley, cf. [43, pp.xvi line 6 bottom up–xvii line 12, and footnote 3 on p.168].

$p^\smile ; q = 1$. However, if such an algebra belonged to some concrete class then its universe would have to determine p and q in a unique way.

We want to find classes of algebras which will be candidates for being concrete classes in a representation theory of relation algebras expanded with quasi-projections. To this end, for any set U one can define the binary relations

$$P_U \stackrel{\text{def}}{=} \{\langle\langle u, x\rangle, u\rangle \ : \ u \in U, \ \langle u, x\rangle \in U\}, \text{ and}$$

$$Q_U \stackrel{\text{def}}{=} \{\langle\langle x, u\rangle, u\rangle \ : \ u \in U, \ \langle x, u\rangle \in U\}.$$

P_U and Q_U are called the *left* and *right projections on* U, respectively. The structure $\langle U, P_U, Q_U\rangle$ is called the *projection structure of* U.

We say that an equivalence relation E on some set U is *closed* if whenever $\langle u, v\rangle \in E$ then $\langle\langle u, v\rangle, u\rangle \in E$ also holds. (This implies that also $\langle\langle u, v\rangle, v\rangle \in E$, since E is an equivalence relation.) We note that if there is some closed E on U then U must be closed under forming ordered pairs 'relative to' E that is, for any $u, v \in U$, if $\langle u, v\rangle \in E$ then $\langle u, v\rangle \in U$ must hold. On the other hand, if $E \subseteq U \times U$ is closed then U is not necessarily closed under the projections P_U and Q_U that is, there can be some $\langle u, v\rangle \in U$ such that $u \notin U$ and/or $v \notin U$. But if $\langle u, v\rangle \in E$ holds then closedness requires not only $u, v \in U$ but the stronger $\langle\langle u, v\rangle, u\rangle, \langle\langle u, v\rangle, v\rangle \in E$.

The following classes of algebras were suggested first in [30].

Definition 4. An algebra $\mathfrak{A} = \langle A, +, \cdot, 1, 0, -, \ ; \ , ^\smile, 1', p, q\rangle$ is called a *partial projection set algebra* (a SetPPA) if conditions (i)–(iii) below hold.

(i) $\langle A, +, \cdot, 1, 0, -, \ ; \ , ^\smile, 1'\rangle$ is a set relation algebra, say with base U and unit E;

(ii) E is closed;

(iii) $p = P_U \cap E = \{\langle\langle u, x\rangle, u\rangle \in E \ : \ u \in U, \ \langle u, x\rangle \in U\}$
$q = Q_U \cap E = \{\langle\langle x, u\rangle, u\rangle \in E \ : \ u \in U, \ \langle x, u\rangle \in U\}$.

That is, p and q are the projections on U, restricted (as relations) to the binary relation E.

A partial projection set algebra is called a *projection set algebra* (a SetPA) if dom p = dom q also holds. Thus SetPA $\subseteq$ SetPPA clearly holds. □

Claim 5. (in ZF^-) *If $\langle\mathfrak{A}, p, q\rangle$ is a projection set algebra (with base U and unit E) then*

$p = \{\langle\langle u, v\rangle, u\rangle \ : \ \langle u, v\rangle \in E\}$, *and*
$q = \{\langle\langle u, v\rangle, v\rangle \ : \ \langle u, v\rangle \in E\}$.

Proof. $p = P_U \cap E \supseteq \{\langle\langle u,v\rangle,u\rangle \ : \ \langle u,v\rangle \in E\}$, because E is closed. For the other inclusion, assume that $\langle\langle u,v\rangle,u\rangle \in E$ for some $\langle u,v\rangle, u \in U$. Then $\langle u,v\rangle \in \mathrm{dom}\ (P_U \cap E) = \mathrm{dom}\ (Q_U \cap E)$, thus $\langle\langle u,v\rangle,v\rangle \in (Q_U \cap E) \subseteq E$. Therefore, since E is an equivalence relation, $\langle u,v\rangle \in E$ also holds. Similar argument works for q. □

The classes SetPA and SetPPA are certainly concrete classes. However, as it is proved in Sain–Németi [38], in ZFC these classes cannot support a representation theory. It is shown in [38] that the equational theories of the classes SetPPA and SetPA are not finitely axiomatizable (cf. section 4 below). Roughly speaking, these negative results are 'due to' the fact that the Axiom of Foundation does not allow the existence of 'not nice' projection structures. E.g. an infinite p-chain $u_0 \stackrel{p}{\to} u_1 \stackrel{p}{\to} u_2 \stackrel{p}{\to} \dots$ would result in an infinite descending $\in$-chain $u_0 \ni \{u_1\} \ni u_1 \ni \{u_2\} \ni \dots$, thus it cannot exist in ZFC. Therefore the idea comes up to experiment with non-well-founded set theories where such objects do exist. The anti-foundation axiom we introduce below postulates the existence of as many non-well-founded sets as possible without harming the Axiom of Extensionality.

Let U be a set and let E be a binary relation on U, i.e. E is a set with $E \subseteq U \times U$. The structure $\langle U,E\rangle$ is called *extensional* if it satisfies the Axiom of Extensionality of ZFC:

$$(\forall x,y \in U)[(\forall z \in U)(\langle z,x\rangle \in E \ \Leftrightarrow \ \langle z,y\rangle \in E) \ \Rightarrow \ x = y].$$

Boffa's weak anti-foundation axiom ($BAFA_0$) is the following:

($BAFA_0$) Every extensional structure is isomorphic to the 'set theoretical $\in$-structure' of some transitive[6] set. That is,

$$\begin{aligned}(\forall \text{ set } U) \ \ & (\forall \text{ set } E \subseteq U \times U) \ (\langle U,E\rangle \text{ is extensional } \Rightarrow \\ & (\exists \text{ transitive } W)\ \langle U,E\rangle \text{ is isomorphic to } \langle W, \in_W\rangle)\,,\end{aligned}$$

where $\in_W \stackrel{\text{def}}{=} \{\langle x,y\rangle \in W \times W : x \in y\}$.

Now we show that, in $ZFC^- + BAFA_0$, both $\mathbb{I}$ SetPPA and $\mathbb{I}$ SetPA are axiomatizable by finitely many equations. To this end, we define a *partial projection algebra* (a PPA) to be an algebra $\mathfrak{A} = \langle A, +, \cdot, 1, 0, -, \ ;\ , ^\smile, 1', p, q\rangle$ satisfying equations (P0)–(P4) below.

(P0) (R0)–(R7)
(P1) $p^\smile ; p \leq 1'$

[6] x is *transitive* iff $(\forall z \in x)\ z \subseteq x$.

(P2) $q^\smile ; q \leq 1'$
(P3) $(p ; p^\smile) \cdot (q ; q^\smile) \leq 1'$
(P4) $p^\smile ; q = 1$

Theorem 6. (in $ZFC^- + BAFA_0$) *Every partial projection algebra is representable that is, it is isomorphic to some partial projection set algebra. Consequently,*

$$\mathsf{PPA} = \mathbb{I}\,\mathsf{SetPPA}$$

that is, the class of representable partial projection algebras is axiomatizable by finitely many equations, namely by (P0)–(P4).

Before proving the theorem we discuss some standard properties of projection structures.

(p1) P_U is a partial function on U.

(p2) Q_U is a partial function on U.

(p3) For any $a, b \in (\mathrm{dom}\ P_U \cap \mathrm{dom}\ Q_U)$, if $P_U(a) = P_U(b)$ and $Q_U(a) = Q_U(b)$ hold then $a = b$.

Claim 7. (in ZF^-) *Let $\mathfrak{A}$ be a set relation algebra with base U and unit E, and let $p, q \in A$. Then*

(i) $p^\smile ; p \leq 1'$ *holds in* $\mathfrak{A}$ *iff* *p satisfies* (p1) *i.e., p is a partial function on U;*

(ii) $q^\smile ; q \leq 1'$ *holds in* $\mathfrak{A}$ *iff* *q satisfies* (p2) *i.e., q is a partial function on U;*

(iii) $(p ; p^\smile) \cdot (q ; q^\smile) \leq 1'$ *holds in* $\mathfrak{A}$ *iff* *p and q satisfy* (p3) *i.e., for any $u, v \in (\mathrm{dom}\ p \cap \mathrm{dom}\ q)$, if there are some $w, z \in U$ such that $\langle u, w\rangle, \langle v, w\rangle \in p$ and $\langle u, z\rangle, \langle v, z\rangle \in q$ then $u = v$;*

(iv) $p^\smile ; q = 1$ *holds in* $\mathfrak{A}$ *iff* $(\forall \langle u, v\rangle \in E)\ (\exists w \in U)\ \langle w, u\rangle \in p$ *and* $\langle w, v\rangle \in q$;

(v) $p ; 1 = q ; 1$ *holds in* $\mathfrak{A}$ *iff* $(\forall u \in U)\ [(\exists v \in U)\langle u, v\rangle \in p \ \Leftrightarrow\ (\exists v \in U)\langle u, v\rangle \in q]$ *(that is,* $\mathrm{dom}\ p = \mathrm{dom}\ q$*).*

Proof of Claim 7. The implications from right to left are obvious by the definitions. The other directions are also easy, but they need some extra consideration. E.g. $p^\smile ; p \leq 1'$ clearly implies, that for any $\langle u, v\rangle \in E$, if $\langle w, u\rangle \in p$ and $\langle w, v\rangle \in p$ for some $w \in U$ then $u = v$ must hold. However, for any $u, v \in U$, if $\langle w, u\rangle \in p$ and $\langle w, v\rangle \in p$ for some $w \in U$ then, since E is an equivalence relation, $\langle u, v\rangle \in E$ must hold, proving that p is a partial function on U. Similar argument works for cases (ii) and (iii). In case (v), $p^\smile ; q = 1$ clearly implies that dom $p \cap$ dom $E =$ dom $q \cap$ dom E. But dom $E = U$, since E is reflexive. ∎

Lemma 8 below is our basic technical tool in proving Theorem 6 and the other representation theorems below. It states that, in $ZF^- + BAFA_0$, properties (p1)–(p3) do characterize projection structures up to isomorphism.

Lemma 8. (in $ZF^- + BAFA_0$) *Let p and q be binary relations on some set U. Then the structure $\langle U, p, q\rangle$ is isomorphic to some projection structure $\langle W, P_W, Q_W\rangle$ iff p and q have properties* (p1)–(p3) *above.*

The proof of Lemma 8 is in section 3. We note that its non-trivial direction obviously fails in ZFC. E.g. consider some one-element set $U = \{u\}$ with dom $p =$ dom $q = \{u\}$ and with $p(u) = q(u) = u$. Then $\langle U, p, q\rangle$ clearly have properties (p1)–(p3). On the other hand, every set which is in the domain of both projections is the ordered pair of its left and right projections. Thus, if $\langle U, p, q\rangle$ were isomorphic to the projection structure of some set $W = \{w\}$ then $w = \langle P_W(w), Q_W(w)\rangle = \langle w, w\rangle = \{\{w\}\}$ would hold. However, such a set w cannot be well-founded. See section 4 for some non-well-founded set theories where this lemma also fails.

Proof of Theorem 6. Let $\mathfrak{A} = \langle A, +, \cdot, 1, 0, -, ;, ^\smile, 1', p, q\rangle$ be a partial projection algebra. Then by (P0)–(P2) and (P4), the relation algebra type reduct $\mathfrak{A}^-$ of $\mathfrak{A}$ is a QRA. Therefore, by Theorem 3, $\mathfrak{A}^-$ is representable, i.e. there is some isomorphism g to a set relation algebra $\mathfrak{B}$, say with base U and unit E. By Claim 7(i)–(iii), $g(p)$ and $g(q)$ satisfy properties (p1)–(p3). Therefore, by Lemma 8, there is some isomorphism F from the structure $\langle U, g(p), g(q)\rangle$ to the projection structure $\langle W, P_W, Q_W\rangle$ of some set W. Let $\mathfrak{C}$ be the set relation algebra on base W that is isomorphic to $\mathfrak{B}$ via the mapping f from binary relations on U to binary relations on W, induced by the bijection $F : U \to W$. (E.g. the unit of $\mathfrak{C}$ is $f(E) = \{\langle f(u), f(v)\rangle : \langle u, v\rangle \in E\}$ and $f(g(p)) = P_W$, $f(g(q)) = Q_W$.) Then the algebra $\langle \mathfrak{B}, g(p), g(q)\rangle$ is certainly isomorphic to $\langle \mathfrak{C}, P_W, Q_W\rangle$. It remains to show that $\langle \mathfrak{C}, P_W, Q_W\rangle \in \mathsf{SetPPA}$. First, since $g(p) \subseteq E$ and $P_W = f(g(p))$, $P_W \subseteq f(E)$ (and similarly, $Q_W \subseteq f(E)$), thus $\langle \mathfrak{C}, P_W, Q_W\rangle = \langle \mathfrak{C}, P_W \cap f(E), Q_W \cap f(E)\rangle$. Finally, we

show that $f(E)$ is closed. Indeed, take some $\langle a, b\rangle \in f(E)$. Since equation (P4) is valid in $\langle \mathfrak{C}, P_W, Q_W\rangle$, Claim 7(iv) implies that there is some $c \in W$ with $P_W(c) = a$ and $Q_W(c) = b$. Since P_W and Q_W are the projections on W, this c must be equal to $\langle a, b\rangle$ that is, $\langle\langle a, b\rangle, a\rangle \in P_W = P_W \cap f(E) \subseteq f(E)$ and similarly, $\langle\langle a, b\rangle, b\rangle \in f(E)$ hold. We proved that the partial projection algebra $\mathfrak{A}$ is isomorphic to the partial projection set algebra $\langle \mathfrak{C}, P_W, Q_W\rangle$.

$$\begin{array}{lll} \langle \mathfrak{A}^-, p, q\rangle \xrightarrow{\ g\ } \langle \mathfrak{B}, g(p), g(q)\rangle & \Big| & \langle U, g(p), g(q)\rangle \\ \qquad\qquad\quad \downarrow f & & \quad \downarrow F \\ \qquad\qquad\quad \langle \mathfrak{C}, P_W, Q_W\rangle \in \mathsf{SetPPA} & & \quad \langle W, P_W, Q_W\rangle \\ \qquad\qquad\quad = \\ \qquad\qquad\quad \langle \mathfrak{C}, P_W \cap f(E), Q_W \cap f(E)\rangle & & \end{array}$$

It is routine to check that every partial projection set algebra satisfies equations (P0)–(P4). We note that the unit of a SetPPA is taken to be closed in order to satisfy (P4). ∎

A partial projection algebra is called a *projection algebra* (a PA) if it also satisfies equation (P5) below.

(P5) $\quad p\,;1 = q\,;1$

Obviously, $\mathsf{PA} \subseteq \mathsf{PPA}$ holds.

Theorem 9. (in $ZFC^- + BAFA_0$) *Every projection algebra is representable that is, it is isomorphic to some projection set algebra. Consequently,*

$$\mathsf{PA} = \mathbb{I}\,\mathsf{SetPA}$$

that is, the class of representable projection algebras is axiomatizable with finitely many equations, namely with (P0)–(P5).

Proof. Using the notation of the previous proof, if (P5) holds in $\mathfrak{A}$ then (P5) also holds in $\langle \mathfrak{C}, P_W, Q_W\rangle$ (since it is isomorphic to $\mathfrak{A}$ via mapping f). Thus, by Claim 7(v), dom P_W = dom Q_W holds. On the other hand, projection set algebras clearly satisfy (P5), also by Claim 7(v). □

Instead of expanding the similarity type of relation algebras with the constant symbols p and q, one can introduce a binary function symbol $\triangledown$, called *fork*, as it is done in e.g. Veloso–Haeberer [44], [45], [17], Baum–Haeberer–Veloso [7], Berghammer et al [8], Frias et al [10], Baum et al [6], Frias–Gordillo [11]. Let R, S be binary relations on some set U. Then $R \triangledown_U S$ is the following binary relation.

$$R\bigtriangledown_U S \stackrel{\text{def}}{=} \{\langle u, \langle v, w\rangle\rangle \ : \ \langle u, v\rangle \in R,\ \langle u, w\rangle \in S\}$$



The picture above motivates the name 'fork'. We note that $R\bigtriangledown_U S$ is not necessarily a binary relation on U, since $\langle v, w\rangle \notin U$ can happen for some $v, w \in U$.

Definition 10. A *fork algebra* (an FA) is an algebra

$$\langle A, +, \cdot, 1, 0, -, \ ;\ , \breve{}, 1', \bigtriangledown\rangle$$

satisfying equations (F0)–(F3) below.

(F0) (R0)–(R7)
(F1) $x\bigtriangledown y = (x\ ;\ (1'\bigtriangledown 1)) \cdot (y\ ;\ (1\bigtriangledown 1'))$
(F2) $(x\bigtriangledown y)\ ;\ (z\bigtriangledown w)\breve{} = (x\ ;\ z\breve{}) \cdot (y\ ;\ w\breve{})$
(F3) $(1'\bigtriangledown 1)\breve{}\bigtriangledown(1\bigtriangledown 1')\breve{} \leq 1'$

A *fork set algebra* (a SetFA) is an algebra $\langle \mathfrak{A}, \bigtriangledown\rangle$ such that

(i) $\mathfrak{A}$ is a set relation algebra, say with base U and unit E;

(ii) $\bigtriangledown$ coincides with $\bigtriangledown_U$;

(iii) E is *closed under* $\bigtriangledown_U$ i.e., for any $R, S \subseteq E$, $R\bigtriangledown_U S \subseteq E$ also holds. □

Claim 11. (in ZF^-) $\quad \mathbb{I}\,\mathsf{SetFA} \subseteq \mathsf{FA}$. ∎

We note that algebras satisfying (F0)–(F3) are called *pre-fork algebras* in Gyuris [15], [14]. Recently, the fork algebra literature started to call the models of (F0)–(F3) *abstract fork algebras*, cf. Frias et al [11], [12].

Next we discuss the connection between the two ways of expanding relation algebras. Namely, we show that fork algebras and projection algebras are equivalent in the strongest possible sense. To this end, consider the terms

$p^{\bigtriangledown} \stackrel{\text{def}}{=} (1'\bigtriangledown 1)\breve{}$ and
$q^{\bigtriangledown} \stackrel{\text{def}}{=} (1\bigtriangledown 1')\breve{}$

of the similarity type of fork algebras; and the term

$x\bigtriangledown^{pq} y := (x\ ;\ p\breve{}) \cdot (y\ ;\ q\breve{})$

of the similarity type of projection algebras.

Claim 12. (in ZF^-) *For any set relation algebra $\mathfrak{A}$, say with base U and unit E, the following hold.*

(i) *E is closed iff E is closed under $\bigtriangledown_U$.*

(ii) *Let $p = P_U \cap E$, $q = Q_U \cap E$. Then for any $R, S \in A$, $R\bigtriangledown^{pq}S = R\bigtriangledown_U S$ holds. Consequently, if $\langle \mathfrak{A}, p, q\rangle \in \mathsf{SetPPA}$ then $\langle \mathfrak{A}, \bigtriangledown^{pq}\rangle \in \mathsf{SetFA}$.*

Proof. For (i): Assume E is closed and let $R, S \subseteq E$. For any $\langle u, \langle v, w\rangle\rangle \in R\bigtriangledown_U S$, $\langle u, v\rangle \in R \subseteq E$ and $\langle u, w\rangle \in S \subseteq E$, thus $\langle v, w\rangle \in E$ also holds. This implies, by E being closed, that $\langle\langle v, w\rangle, v\rangle \in E$ and therefore, by $\langle u, v\rangle \in E$, $\langle u, \langle v, w\rangle\rangle \in E$.

In the other direction, for any $\langle u, v\rangle \in E$, $\langle\langle u, v\rangle, u\rangle \in (1'\bigtriangledown_U 1)^\smile$ thus, by E being closed under $\bigtriangledown_U$, $\langle\langle u, v\rangle, u\rangle \in E$ holds.

Item (ii) can be proved by a routine computation. □

Theorem 13. (in ZF^-) *The classes* FA *and* PA *are equivalent in the following sense (cf.* McKenzie–McNulty–Taylor [27, Def. 4.138]*).*

(i) [cf. Gyuris [15], [14]] *For any $\langle \mathfrak{A}, \bigtriangledown\rangle \in \mathsf{FA}$, $\langle \mathfrak{A}, p^\triangledown, q^\triangledown\rangle \in \mathsf{PA}$.*

(ii) *For any $\langle \mathfrak{A}, p, q\rangle \in \mathsf{PPA}$, $\langle \mathfrak{A}, \bigtriangledown^{pq}\rangle \in \mathsf{FA}$.*

(iii) *The equation $x\bigtriangledown y = x\bigtriangledown^{p^\triangledown q^\triangledown} y$ is valid in* FA.

(iv) *The equations $p = p^{\triangledown^{pq}}$ and $q = q^{\triangledown^{pq}}$ are valid in* PA.

Proof. For (i): We quote from [15], [14] the derivations of (P1), (P2) and (P4):

$$p^{\triangledown^\smile} ; p^\triangledown = (1'\bigtriangledown 1)^{\smile\smile} ; (1'\bigtriangledown 1)^\smile \overset{\text{(F0),(F2)}}{=} (1' ; 1'^\smile) \cdot (1 ; 1^\smile) \overset{\text{(F0)}}{\leq} 1'$$

$$q^{\triangledown^\smile} ; q^\triangledown = (1\bigtriangledown 1')^{\smile\smile} ; (1\bigtriangledown 1')^\smile \overset{\text{(F0),(F2)}}{=} (1 ; 1^\smile) \cdot (1' ; 1'^\smile) \overset{\text{(F0)}}{\leq} 1'$$

$$p^{\triangledown^\smile} ; q^\triangledown = (1'\bigtriangledown 1)^{\smile\smile} ; (1\bigtriangledown 1')^\smile \overset{\text{(F0),(F2)}}{=} (1' ; 1^\smile) \cdot (1 ; 1'^\smile) \overset{\text{(F0)}}{=} 1.$$

(P3) can be derived as follows.

$$(p^\triangledown ; p^{\triangledown^\smile}) \cdot (q^\triangledown ; q^{\triangledown^\smile}) \overset{\text{(F2)}}{=} (p^\triangledown \bigtriangledown q^\triangledown) ; (p^\triangledown \bigtriangledown q^\triangledown)^\smile =$$
$$= [(1'\bigtriangledown 1)^\smile \bigtriangledown (1\bigtriangledown 1')^\smile] ; [(1'\bigtriangledown 1)^\smile \bigtriangledown (1\bigtriangledown 1')^\smile]^\smile \overset{\text{(F0),(F3)}}{\leq} 1' ; 1'^\smile \overset{\text{(F0)}}{=} 1'.$$

Finally, for (P5):

$$1 ; (1'\bigtriangledown 1) \overset{\text{(F0),(F1)}}{\leq} 1 ; (1\bigtriangledown 1) \overset{\text{(F1)}}{=} 1 ; [(1 ; (1'\bigtriangledown 1)) \cdot (1 ; (1\bigtriangledown 1'))]$$
$$\overset{\text{(F0)}}{\leq} 1 ; (1\bigtriangledown 1').$$

Therefore $p^{\triangledown} ; 1 = (1' \triangledown 1)^{\smile} ; 1 \stackrel{(F0)}{=} (1 ; (1' \triangledown 1))^{\smile} \stackrel{(F0)}{\leq} (1 ; (1 \triangledown 1'))^{\smile} \stackrel{(F0)}{=} q^{\triangledown} ; 1$.

For (ii): Assume that some of the equations (F0)–(F3) (with $\triangledown^{pq}$ in place of $\triangledown$) fails in some PPA. Then it also fails in some countable subalgebra $\mathfrak{B}$ of the PPA in question. The relation algebraic reduct of $\mathfrak{B}$ is a QRA thus, by Theorem 3, it is representable that is, isomorphic to an algebra whose elements are binary relations. (For countable algebras, Theorem 3 can be proved without the Axiom of Choice.) Now it is routine to check that equations (F0)–(F3) (with $\triangledown^{pq}$) hold in those PPA's whose elements are binary relations.

For (iii): $x \triangledown^{p^{\triangledown} q^{\triangledown}} y = (x ; p^{\triangledown \smile}) \cdot (y ; q^{\triangledown \smile}) = [(x ; (1' \triangledown 1)^{\smile \smile}] \cdot [y ; (1 \triangledown 1')^{\smile \smile}] \stackrel{(F0),(F1)}{=} x \triangledown y$.

For (iv): $p^{\triangledown^{pq}} = (1' \triangledown^{pq} 1)^{\smile} = [(1' ; p^{\smile}) \cdot (1 ; q^{\smile})]^{\smile} \stackrel{(P0)}{=} (1' ; p^{\smile})^{\smile} \cdot (1 ; q^{\smile})^{\smile} \stackrel{(P0)}{=} = p \cdot (q ; 1) \stackrel{(P5)}{=} p \cdot (p ; 1) \stackrel{(P0)}{=} p$. One can prove $q^{\triangledown^{pq}} = q$ in a similar way. □

It is proved in Sain–Németi [38] that, in ZFC, the class SetFA cannot support a representation theory (cf. section 4 below). However, in $ZFC^- + BAFA_0$, we have a representation theorem for partial projection algebras (Theorem 6 above) which, by Claim 12 and Theorem 13, can be 'converted to' a representation theorem of fork algebras.

Theorem 14. (in $ZFC^- + BAFA_0$) *Every fork algebra is representable that is, it is isomorphic to some set fork algebra. Consequently,*

$$\mathsf{FA} = \mathbb{I}\,\mathsf{SetFA}$$

that is, the class of representable fork algebras is axiomatizable with finitely many equations, namely with (F0)–(F3).

Proof. Let $\langle \mathfrak{A}, \triangledown \rangle \in \mathsf{FA}$. Then, by Theorem 13(i), $\langle \mathfrak{A}, p^{\triangledown}, q^{\triangledown} \rangle \in \mathsf{PPA}$. Thus, by Theorem 6, $\langle \mathfrak{A}, p^{\triangledown}, q^{\triangledown} \rangle$ is isomorphic to some $\langle \mathfrak{B}, p, q \rangle \in \mathsf{SetPPA}$. Therefore, by Claim 12, $\langle \mathfrak{B}, \triangledown^{pq} \rangle \in \mathsf{SetFA}$ which is certainly isomorphic to $\langle \mathfrak{A}, \triangledown^{p^{\triangledown} q^{\triangledown}} \rangle$. But, by Theorem 13(iii), the latter is just $\langle \mathfrak{A}, \triangledown \rangle$. □

Remark 15. Besides their finite axiomatizability, the classes $\mathbb{I}\,\mathsf{SetPPA}$, $\mathbb{I}\,\mathsf{SetPA}$ and $\mathbb{I}\,\mathsf{SetFA}$ have other 'nice' properties in the set theory $ZFC^- + BAFA_0$. Namely, in $ZFC^- + BAFA_0$, these classes enjoy the so called *super amalgamation property* (and hence the so called *strong amalgamation property*), in the sense of Madarász [21], Marx [25]. The proofs of these theorems are analogous with the proof (in ZFC) of the strong amalgamation property of QRA, given in Sain [34].

3 Projection functions in non-well-founded set theories

Theorem 16. [cf. Aczel [1]] *If ZFC^- is consistent then 'ZFC^-+$BAFA_0$+ collection principle' is also consistent.*

Proof. Since in [1] this theorem is not explicitly stated, we list the main 'landmarks' of its proof in [1]. Our formulation of $BAFA_0$ is denoted by BA_1 in [1]. *Boffa's Anti-foundation Axiom* $BAFA$ is formulated on p.59, and Exercise 5.14 on p.65 states that $BAFA$ implies BA_1. On p.61 it is stated that '$BAFA$ holds in the universe iff the universe is *superuniversal*' (see p.13 for the definition of a *system*, and p.59 for superuniversality). Theorem 5.9 on p.61 states that every superuniversal system is *full* (see p.34 for definition). Rieger's theorem on p.119 states that any full system is a class model of 'ZFC^-+collection principle'. Next, Theorem 5.18 on p.66 states that, assuming that there is a definable well-ordering of the universe, there exists a superuniversal system. Finally, the assumption that 'there is a definable well-ordering of the universe' holds in the constructible universe L, and $V = L$ is consistent with ZFC^-. □

Proof of Lemma 8 of section 2. We start with introducing some notation. Let E be a binary relation on some set V. For $u, v_1, \ldots, v_n \in V$,

$$u = \{v_1, \ldots, v_n\}_E \stackrel{\text{def}}{\iff} (\forall z \in V)\ (\langle z, u\rangle \in E \ \Leftrightarrow\ (z = v_1 \text{ or} \ldots \text{or } z = v_n)).$$

In case of $u = \{v_1, \ldots, v_n\}_E$ we say that $v_1, \ldots, v_n$ are the *E-elements of u*. Similarly, for any $u \in V$ and $v \neq w \in V$,

$$\begin{array}{lll} u = \langle v, v\rangle_E & \stackrel{\text{def}}{\iff} & (\exists x \in V)\ x = \{v\}_E \text{ and } u = \{x\}_E \\ u = \langle v, w\rangle_E & \stackrel{\text{def}}{\iff} & (\exists x, y \in V)\ x = \{v\}_E,\ y = \{v, w\}_E \text{ and } u = \{x, y\}_E. \end{array}$$

If $u = \langle v, w\rangle_E$ we say that u is the *E-pair of v and w*.

Now assume that some structure $\langle U, p, q\rangle$ satisfying (p1)–(p3) is given. We will construct a structure $\langle V, E\rangle$ for some set $V \supseteq U$ such that the following hold.

$$(\sharp) \quad (\forall u, v \in U)\ (p(u) = v \iff u = \langle v, x\rangle_E \text{ for some } x \in V)$$

$$(\sharp\sharp) \quad (\forall u, v \in U)\ (q(u) = v \iff u = \langle x, v\rangle_E \text{ for some } x \in V).$$

The motivation for the construction is as follows. The structure $\langle V, E\rangle$ is intended to 'mimic' the $\in$-structure of the transitive closure[7] of some set

[7] The *transitive closure* of some set x is the smallest transitive set containing x as a subset.

whose projection structure is isomorphic to $\langle U, p, q\rangle$. If we want to find some transitive set such that its $\in$-structure is isomorphic to $\langle V, E\rangle$ then we also have to ensure the extensionality of $\langle V, E\rangle$. In order to achieve extensionality, in the construction of V we always take new sets unless it is 'explicitly required' that some set coincides with one of the sets already taken.

What kind of new sets do we need in V to have properties ($\sharp$) and ($\sharp\sharp$)? Certainly, for each $v \neq w \in U$ such that $p(u) = v$ and $q(u) = w$ for some $u \in \operatorname{dom} p \cap \operatorname{dom} q$, we need a new[8] set, call it pair_{vw}, which 'mimics' the set $\{v, w\}$. We also take the following new sets:

- singl_v, for each $v \in U$ such that $v = p(u)$ for some $u \in \operatorname{dom} p$;
- $\mathsf{pair}_{?u}$ and $?_u$, for each $u \in \operatorname{dom} p - \operatorname{dom} q$;
- pair_{*u}, singl_{*u} and $*_u$, for each $u \in \operatorname{dom} q - \operatorname{dom} p$;
- $\clubsuit_u$ and $\spadesuit_u$, for each $u \in U - (\operatorname{dom} p \cup \operatorname{dom} q)$.

Now let V be the expansion of U with these new sets.

Next we are defining a binary relation E on V. First, for each $u \in \operatorname{dom} p \cap \operatorname{dom} q$, add the following pairs to E (in the figures below $\langle x, y\rangle \in E$ is illustrated by $x \to y$):





[8]Such new sets certainly exist, even in non-well-founded set theories. However, to be explicit, e.g. one can choose pair_{vw} to be the set $\{\alpha, v, w\}$, where α is the smallest ordinal which is not in the transitive closure of U.

For each $u \in \text{dom } p - \text{dom } q$, add the following pairs to E:



For each $u \in \text{dom } q - \text{dom } p$, add the following pairs to E:



For each $u \in U - (\text{dom } p \cup \text{dom } q)$, add the following pairs to E:



Claim 16.1. *$\langle V, E\rangle$ satisfies properties (♯) and (♯♯) above.*

Proof of Claim 16.1. To prove (♯), take some $u, v \in U$ and assume $p(u) = v$. If $u \in \text{dom } p \cap \text{dom } q$ and also $q(u) = v$ then $u = \{\mathsf{singl}_v\}_E$ and $\mathsf{singl}_v = \{v\}_E$, thus $u = \langle v, v\rangle_E$. If $q(u) \neq v$ then $u = \{\mathsf{singl}_v, \mathsf{pair}_{vq(u)}\}_E$ and $\mathsf{singl}_v = \{v\}_E$, $\mathsf{pair}_{vq(u)} = \{v, q(u)\}_E$, thus $u = \langle v, q(u)\rangle_E$. If $u \in \text{dom } p - \text{dom } q$ then $u = \{\mathsf{singl}_v, \mathsf{pair}_{?u}\}_E$ and $\mathsf{singl}_v = \{v\}_E$, $\mathsf{pair}_{?u} = \{v, ?_u\}_E$, thus $u = \langle v, ?_u\rangle_E$.

For the other direction of (♯), assume that $u = \langle v, x\rangle_E$, for some $u, v \in U$ and $x \in V$. Then u must belong to dom $p \cup$ dom q, since inspection shows that elements of $U - (\text{dom } p \cup \text{dom } q)$ are not E-pairs. We show that $u \in \text{dom } p$ must hold. Indeed, assume $u \in \text{dom } q - \text{dom } p$. Then $u = \{\mathsf{singl}_{*u}, \mathsf{pair}_{*u}\}_E$. Thus either $\mathsf{singl}_{*u} = \{v\}_E$ or $\mathsf{pair}_{*u} = \{v\}_E$. In the first case $v = *_u$, contradicting $v \in U$. The second case cannot happen, since

pair_{*u} has two E-elements. So we have $u \in \mathrm{dom}\ p$. Then $\langle \mathsf{singl}_{p(u)}, u\rangle \in E$ and $\mathsf{singl}_{p(u)}$ has exactly one E-element. Therefore it follows from $u = \langle v, x\rangle_E$ that $\mathsf{singl}_{p(u)} = \{v\}_E$, implying $p(u) = v$.

Property ($\sharp\sharp$) can be proved similarly. ∎

Claim 16.2. *$\langle V, E\rangle$ is extensional.*

Proof of Claim 16.2. We will say that $z \in V$ *E-distinguishes* $x \in V$ from $y \in V$ if $\langle z, x\rangle \in E$ and $\langle z, y\rangle \notin E$ or the other way round. We say that $x, y \in V$ are *E-distinguishable* if there is some $z \in V$ that E-distinguishes between them. We have to prove that any $x \neq y \in V$ are E-distinguishable. To this end, let the subset $V_0 \subseteq V$ be defined as follows.

$$V_0 \stackrel{\text{def}}{=} U \cup \{\mathsf{singl}_v \ : \ v \in U \text{ and } \exists u \in U \text{ with } p(u) = v\} \cup \\ \cup \{\mathsf{pair}_{vw} \ : \ v \neq w \in U \text{ and } \exists u \in U \text{ with } p(u) = v,\ q(u) = w\}.$$

First, we show that any $x \in V_0$ and $y \in V - V_0$ are E-distinguishable. Indeed, $?_u$ E-distinguishes members of V_0 from $?_u$ and also from $\mathsf{pair}_{?u}$; $*_u$ E-distinguishes members of V_0 from $*_u$, from pair_{*u} and also from singl_{*u}; $\spadesuit_u$ E-distinguishes members of V_0 from $\spadesuit_u$ and also from $\clubsuit_u$.

Second, we show that any $x \neq y \in V - V_0$ are E-distinguishable. Indeed, for any $u \in \mathrm{dom}\ p - \mathrm{dom}\ q$, $\mathsf{pair}_{?u}$ has two E-elements, while $?_u$ has only one, thus they must be E-distinguishable. For any $u \neq v \in \mathrm{dom}\ p - \mathrm{dom}\ q$, $?_u$ E-distinguishes $?_u$ from $?_v$, and also $\mathsf{pair}_{?u}$ from $\mathsf{pair}_{?v}$. For any $u \in \mathrm{dom}\ q - \mathrm{dom}\ p$, both pair_{*u} and $*_u$ have two E-elements, while singl_{*u} has only one, thus singl_{*u} is E-distinguishable from pair_{*u} and also from $*_u$; and e.g. $q(u)$ E-distinguishes pair_{*u} from $*_u$. For any $u \neq v \in \mathrm{dom}\ q - \mathrm{dom}\ p$, $*_u$ E-distinguishes $*_u$ from $*_v$, singl_{*u} from singl_{*v}, and also pair_{*u} from pair_{*v}. $?_u$ E-distinguishes $?_u$ from $*_v$, from singl_{*v}, and from pair_{*v}; and also $\mathsf{pair}_{?u}$ from $*_v$, from singl_{*v}, and from pair_{*v}. $\clubsuit_u$ E-distinguishes $\clubsuit_u$ from $\spadesuit_u$, and $\spadesuit_u$ E-distinguishes other members of $V - V_0$ from $\spadesuit_u$ and from $\clubsuit_u$.

Third, we show that any $x \neq y \in V_0$ are E-distinguishable. Indeed, for any $u, v, w \in U$, u and singl_v, and also u and pair_{vw} are E-distinguishable, since every E-element of singl_v and pair_{vw} belongs to U, while u always has some E-element which does not belong to U. pair_{vw} has two E-elements, while singl_v has only one, thus they must be E-distinguishable. For any $v_1 \neq v_2$ such that $v_1 = p(u_1)$, $v_2 = p(u_2)$ for some $u_1, u_2 \in U$, v_1 E-distinguishes singl_{v_1} from singl_{v_2}. Take any $v_1 \neq w_1$ and $v_2 \neq w_2$ such that $\{v_1, w_1\} \neq \{v_2, w_2\}$ and $v_1 = p(u_1)$, $w_1 = q(u_1)$, $v_2 = p(u_2)$, $w_2 = q(u_2)$ for some $u_1, u_2 \in U$. If $v_1 \neq v_2$ then v_1 E-distinguishes $\mathsf{pair}_{v_1 w_1}$ from $\mathsf{pair}_{v_2 w_2}$. If $v_1 = v_2$ then $w_1 \neq w_2$ must hold, thus w_1 E-distinguishes $\mathsf{pair}_{v_1 w_1}$ from

$\mathsf{pair}_{v_2w_2}$. It remains to E-distinguish the elements of U from each other. U can be partitioned to the following four parts:

$U = (\mathrm{dom}\ p \cap \mathrm{dom}\ q) \ \cup\ (\mathrm{dom}\ p - \mathrm{dom}\ q) \ \cup\ (\mathrm{dom}\ q - \mathrm{dom}\ p) \ \cup\ (U - (\mathrm{dom}\ p \cup \mathrm{dom}\ q))$.

Then for any $u \in U$,

$$u \in \mathrm{dom}\ p - \mathrm{dom}\ q \iff \langle \mathsf{pair}_{?u}, u\rangle \in E,$$

$$u \in \mathrm{dom}\ q - \mathrm{dom}\ p \iff \langle \mathsf{singl}_{*u}, u\rangle \in E$$

$$u \in U - (\mathrm{dom}\ p \cup \mathrm{dom}\ q) \iff \langle \clubsuit_u, u\rangle \in E.$$

Thus, if u and v are in different blocks of the partition above then they must be E-distinguishable.

If $u \neq v \in \mathrm{dom}\ p \cap \mathrm{dom}\ q$ then, by property $(p3)$, either $p(u) \neq p(v)$ or $q(u) \neq q(v)$. If $p(u) \neq p(v)$ then $\mathsf{singl}_{p(u)}$ E-distinguishes u from v. If $p(u) = p(v)$ and $q(u) \neq q(v)$ then either $p(u) = q(u)$ or not. If $p(u) = q(u)$ then $p(v) \neq q(v)$, thus v has two E-elements and u has only one, thus they must be E-distinguishable. If $p(u) \neq q(u)$ then $\mathsf{pair}_{p(u)q(u)}$ E-distinguishes u from v.

If $u \neq v \in \mathrm{dom}\ p - \mathrm{dom}\ q$ then $\mathsf{pair}_{?u}$ E-distinguishes between them. If $u \neq v \in \mathrm{dom}\ q - \mathrm{dom}\ p$ then singl_{*u} E-distinguishes between them. If $u \neq v \in U - (\mathrm{dom}\ p \cup \mathrm{dom}\ q)$ then u E-distinguishes between them, completing the proof of Claim 16.2. ∎

So we have an extensional structure $\langle V, E\rangle$. Therefore, by $BAFA_0$, $\langle V, E\rangle$ is isomorphic to $\langle Z, \in_Z\rangle$ for some transitive set Z. Call this isomorhpism f. Then it is easy to check that

$(\sharp\sharp\sharp)$ $\quad (\forall x, y, z \in V)\ (x = \langle y, z\rangle_E \iff f(x) = \langle f(y), f(z)\rangle)$.

Let $W \stackrel{\mathrm{def}}{=} \{f(u) : u \in U\}$. We claim that the restriction of f to U is an isomorphism between $\langle U, p, q\rangle$ and the projection structure $\langle W, P_W, Q_W\rangle$. Indeed, for any $u, v \in U$, if $\langle u, v\rangle \in p$ then, by $(\sharp)$, $u = \langle v, x\rangle_E$ for some $x \in V$ therefore, by $(\sharp\sharp\sharp)$, $f(u) = \langle f(v), f(x)\rangle$. Since $u, v \in U$ implies $f(u), f(v) \in W$, we obtain $\langle f(u), f(v)\rangle \in P_W$. For the other direction, assume $\langle f(u), f(v)\rangle \in P_W$ for some $u, v \in U$. Then $f(u) = \langle f(v), y\rangle$ for some set y. But $f(u) \in W \subseteq Z$ and Z is transitive, thus there is some $x \in V$ with $f(x) = y$. This implies, by $(\sharp\sharp\sharp)$ and $(\sharp)$, that $\langle u, v\rangle \in p$.

$(\langle u, v\rangle \in q \iff \langle f(u), f(v)\rangle \in Q_W)$ can be proved in a similar way, completing the proof of Lemma 8. ∎

4 A summary of negative results

In this section we summarize some negative results concerning the representation theory of the expanded relation algebras introduced in section 2. We will see that it is not really the Axiom of Foundation that is 'responsible' for the failure of the representation theorems in ZFC, but rather the following version of Extensionality, called the *Axiom of Strong Extensionality* (SE). This axiom 'helps to identify' non-well-founded sets.

(SE) For any transitive sets x, y, if the structures $\langle x, \in\rangle$ and $\langle y, \in\rangle$ are isomorphic then $x = y$.

SE is an axiom of 'forbidding' type. It does not state the existence of anything, but it forbids to exist 'too many' or 'too ugly' sets. E.g. one can prove from SE that there are no infinite descending $\in$-chains of distinct singletons as follows. Assume we have such a chain, say

$$u_0 \ni u_1 \ni u_2 \ni \dots \quad .$$

Then the transitive closure $trc(u_0)$ of u_0 is not equal to $trc(u_1)$, since $u_1 \in trc(u_0) - trc(u_1)$. But $\langle trc(u_0), \in\rangle$ and $\langle trc(u_1), \in\rangle$ are isomorphic structures, contradicting SE.

SE is a theorem both of ZF and of the non-well-founded set theories in Barwise–Etchemendy [5]. To unify the various non-well-founded set theories, Aczel [1, section 4] introduces the flexible axiom $AFA^{\sim}$, where the special equivalence relation $\sim$ is a parameter of the axiom. For various choices of $\sim$, we obtain different anti-foundation axioms, e.g. Scott's anti-foundation axiom, the basic variant AFA of Aczel, Finsler's anti-foundation axiom, etc. [1, Prop.4.26, p.52] shows that SE is a theorem in all of these set theories. (Our axiom SE is a part of Finsler's axiom and is denoted by $FAFA_2$ in [1].)

The following claim states that Lemma 8 of section 3 fails in strongly extensional set theories.

Claim 17. (in $ZF^{-}+SE$) *There is some structure $\langle U, p, q\rangle$ satisfying properties* (p1)–(p3) *which is not isomorphic to any projection system.*

Proof. Consider the set ω of natural numbers and the usual successor function *suc*. Then $\langle \omega, suc, suc\rangle$ certainly satisfies (p1)–(p3). Assume that $\langle \omega, suc, suc\rangle$ is isomorphic to some projection system $\langle U, P_U, Q_U\rangle$. Then U is a countable set, say $U = \{u_n : n \in \omega\}$, and for any $n \in \omega$, $u_n = \langle P_U(u_n), Q_U(u_n)\rangle = \langle u_{n+1}, u_{n+1}\rangle = \{\{u_{n+1}\}\}$. Therefore

$$u_0 \ni \{u_1\} \ni u_1 \ni \{u_2\} \ni u_2 \ni \dots$$

is an infinite descending $\in$-chain of distinct singletons, contradicting SE. □

Our main representation theorems also fail if one uses some strongly extensional metatheory: the classes of representable algebras are not axiomatizable by finitely many equations. Moreover, the following stronger theorem holds.

Theorem 18. [Sain-Németi [38]] (in $ZF^- + SE$) *The classes* $\mathbb{I}\,\mathsf{SetPPA}$, $\mathbb{I}\,\mathsf{SetPA}$ *and* $\mathbb{I}\,\mathsf{SetFA}$ *are not axiomatizable by any set of first-order sentences.*

Idea of proof. It is proved in [38] that one can find a 'bad' ultrapower of some set algebra (belonging to any of the above classes) such that any set algebra which is isomorphic to that 'bad' ultrapower must contain an infinite descending $\in$-chain of distinct singletons. The Axiom of Choice can be eliminated by replacing ultrapowers with using the compactness theorem for countable languages. ∎

The following theorem states that, in strongly extensional set theories, even the equational theories of the above classes are very far from being computable. This theorem was first proved in Sain–Simon [39] in ZF. The proof in $ZF^- + SE$ is published in [38].

Theorem 19. [cf. Sain-Németi [38]] (in $ZF^- + SE$) *The equational theories of the classes* SetPPA, SetPA *and* SetFA *are* Π^1_1*-hard. Consequently, they are not recursively enumerable. In particular, they are not finitely axiomatizable.*

Idea of proof. It is proved in [38] that there is a recursive translation, sending each Π^1_1-formula φ of the language of arithmetic to some equation e_φ of the similarity type of projection algebras (or of fork algebras), such that

$$\varphi \text{ holds in the standard model of arithmetic} \quad \text{iff} \quad (\text{say}) \ \mathbb{I}\,\mathsf{SetPA} \models e_\varphi.$$

The main idea behind the above translation is that in any $\mathfrak{A} \in \mathsf{SetPA}$, one can 'find' a structure isomorphic to some model of arithmetic. In particular, some subset of $1'^{\mathfrak{A}}$ plays the role of ω, and $(p^{\mathfrak{A}} \cap q^{\mathfrak{A}})^\smile$ plays the role of the successor function. SE comes into the picture when one tries to prove that $p^{\mathfrak{A}} \cap q^{\mathfrak{A}}$ is not 'nonstandard' in the sense that there is no infinite 'predecessor-chain' that is, there is no infinite sequence of distinct elements $u_0, u_1, u_2, \cdots \in U$ with

$$(\forall n \in \omega)\ (p^{\mathfrak{A}} \cap q^{\mathfrak{A}})(u_n) = u_{n+1}.$$

Since $\mathfrak{A} \in \mathsf{SetPA}$, $p^{\mathfrak{A}}$ and $q^{\mathfrak{A}}$ are the projections on U, thus such a chain would result in an infinite descending $\in$-chain of distinct singletons

$$u_0 \ni \{u_1\} \ni u_1 \ni \{u_2\} \ni u_2 \ni \dots$$

(cf. the proof of Claim 17 above), contradicting SE. ∎

We note that the non-recursive enumerability of the equational theories in Theorem 19 above remains even if one considers bigger classes where only some weakening of the associativity of relation composition holds, see [20].

Next we would like to mention some negative results of [38] which already hold in ZF^-. These theorems are about certain subclasses of our classes of representables. A set projection algebra $\mathfrak{A}$ with base U is called a *true pairing algebra* (a TPA) (cf. Maddux [23], [22]) if its unit is $U \times U$ and[9] there is some nonempty set U_0 (called the set of *urelements*) with the following properties:

(i) U is the smallest set such that $U_0 \subseteq U$ and $U \times U \subseteq U$;

(ii) $U - U_0 = U \times U$.

We note that for any $\mathfrak{A} \in \mathsf{SetPA}$ with base U, $\mathfrak{A} \in \mathsf{TPA}$ iff there is some set U_0 of urelements which does not contain ordered pairs and $U = \bigcup_{n\in\omega} U_n$, where $U_{n+1} = (\bigcup_{i\leq n} U_i) \times (\bigcup_{i\leq n} U_i)$. Thus, the base of some TPA cannot contain infinite $(p \cup q)$-chains. From this it follows that

(∗) (in ZF^-) the base of some TPA cannot contain infinite descending $\in$-chains of distinct singletons.

Indeed, if $u_0 \ni u_1 \ni u_2 \ni \ldots$ were such a chain then $u_0 = \{u_1\} = \{\{u_2\}\} = \langle u_2, u_2\rangle$, $u_2 = \langle u_4, u_4\rangle$, etc. would hold. Therefore $\langle u_0, u_2, u_4, \ldots\rangle$ would be an infinite $(p \cup q)$-chain.

The following theorem was first proved in Sain–Simon [39], cf. also [28]. The proof is published in [38].

Theorem 20. (in ZF^-)

(i) $\mathbb{I}\,\mathsf{TPA}$ *and* $\mathbb{SP}\mathsf{TPA}$ *are not axiomatizable by any set of first-order sentences.*

(ii) The equational theory of TPA *is* Π^1_1*-hard. Therefore it is not finitely axiomatizable.*

Idea of proof. It is in [38]. In fact, the proofs of Theorem 18 and 19 go through without SE in this case, because of (∗) above. ∎

[9] In [23], notation 'TPA' denoted the class of those TPA's which consist of all the subsets of some set $U \times U$.

Finally, we note that one can define a class, called the class of *true fork algebras*, in a way similar to TPA. Then the equational theory of true fork algebras is provably non-finitely axiomatizable, already in ZF^- (hence the same holds in any extension, like $ZFC^- + BAFA_0$, of ZF^-). The same applies to the so called 'concatenation version' (cf. Haeberer–Veloso [17]), to the 'free groupoid version' (cf. Baum–Haeberer–Veloso [7]) and to the 'finite tree version' (cf. Haeberer–Baum–Schmidt [16]) of fork algebras, because the equational theory of TPA is interpretable in the equational theories of these classes. These results as well as some further negative results in this line are proved in [38].

5 The finitization problem, connections with logic

The *finitization problem* of algebraic logic (cf. e.g. [29]) asks whether the reason for non-finite axiomatizability of the classes of representable relation algebras or finite dimensional representable cylindric algebras is caused only by Tarski's not including enough operations on relations into the definitions of these classes. A possible formulation of this problem is the following.

For any set U, let $\mathcal{R}(U \times U)$ denote the full set relation algebra with base U (cf. Definition 2). Consider the question whether

> one can add to the operations of set relation algebras finitely many *natural* operations $f_1, \ldots, f_n$ such that the class
>
> $$\mathbb{SP}\{\langle \mathcal{R}(U \times U), f_1, \ldots, f_n\rangle \ : \ U \text{ is a set}\}$$
>
> would be a finitely axiomatizable variety (or quasivariety).

A key issue here is what 'naturalness' requirements the extra operations $f_1, \ldots, f_n$ should satisfy. The so called *soft* version of the finitization problem requires $f_1, \ldots, f_n$ to be set theoretically definable (without parameters). This requirement is motivated by the fact that the other operations $\cap$, $;$, $\breve{}$, etc. of set relation algebras are set theoretically definable operations (see below for further motivations coming from logic). Typical examples for set theoretically definable operations are the projections P_U, Q_U and the fork operation $\bigtriangledown_U$. The *hard* version of the finitization problem requires $f_1, \ldots, f_n$ to be invariant under all permutations of U.[10] A typical example for such

[10] A, say, unary operation f is called *permutation invariant* on U if for any permutation $\pi : U \to U$, and $R \subseteq U \times U$, $\pi(f_i(R)) = f_i(\pi(R))$ holds (where $\pi(R) \stackrel{\text{def}}{=} \{\langle \pi(u), \pi(v)\rangle : \langle u, v\rangle \in R\}$).

an f_i is taking the transitive closure R^* of a relation R. In ZFC, no solution is known for either the soft or the hard version of the finitization problem, cf. e.g. [29].[11] As a contrast, by Theorems 6, 9 and 14 way above, in $ZFC^- + BAFA_0$ the soft version of the finitization problem does admit positive solutions.

There are various motivations for the finitization problem coming from 'pure' logic.

(1) First order logic $L_{\omega\omega}$ is such that its valid formula schemas cannot be axiomatized by any finite schema of axioms and rules. The question is whether there is a new logic $L^+_{\omega\omega}$, containing $L_{\omega\omega}$ as a (definable) fragment, such that $L^+_{\omega\omega}$ admits a finite axiomatization of schemas. The new connectives of $L^+_{\omega\omega}$ are subject to some 'naturalness' conditions (this is where the above conditions on f_i come from).

(2) The finite variable fragments L_n of $L_{\omega\omega}$ are strongly incomplete in the sense that they admit no finite schema axiomatization, cf. [29].[12] The task here is to expand L_n with new logical connectives (e.g. with generalized quantifiers) such that the so obtained L^+_n would admit a complete finite axiomatization.

(3) In Marx–Venema [26, pp.115–123], it is described how $L_{\omega\omega}$ and L_n can be translated to equivalent propositional modal logics. It turns out there that these modal counterparts of $L_{\omega\omega}$ and L_n are not finitely axiomatizable. The question is whether these modal logics can be expanded with new (natural) connectives such that they would become finitely axiomatizable.

Using the (algebraic) results in this paper, the soft versions of the above problems (of logic) can receive positive answers, cf. e.g. [31], [32, section 5], [33].

Acknowledgement: A shorter, preliminary version of this paper appeared as [32]. (It was intended to be a pre-publication of this material and therefore was submitted to the Bulletin of the IGPL – which, at that time, was declared to be a pre-publication forum. In its "General Description" it was written: "Publishing in the Bulletin ... will not exclude the author from submitting a better version to a journal".)

[11] This applies only to the so called finite dimensional version of the problem. The infinite dimensional version was solved positively in Sain [35], [36], Sain–Gyuris [37].

[12] The same applies for the versions $L^n_{\infty,\omega}$ and $L^n_{\kappa,\omega}$ (with $\kappa \geq \omega$) of L_n which allow infinite conjunctions built up from L_n formulas.

References

[1] P. Aczel. *Non–well–founded Sets*. Volume 14 of *Lecture Notes*, CSLI Publications, Stanford, 1988.

[2] H. Andréka. Complexity of the Equations Valid in Algebras of Relations. Part I: Strong non-finitizability; Part II: Finite axiomatizations. *Annals of Pure and Applied Logic*, 89:149–209 and 211–229, 1997.

[3] H. Andréka, B. Jónsson, and I. Németi. Free Algebras in Discriminator Varieties. *Algebra Universalis*, 28:401–447, 1991. Abstracted in [9], 1–14.

[4] H. Andréka, Á. Kurucz, and I. Németi. Connections between Axioms of Set Theory and Basic Theorems of Universal Algebra. *Journal of Symbolic Logic*, 59(3):912–922, 1994.

[5] J. Barwise and J. Etchemendy. *The Liar*. Oxford University Press, 1989.

[6] G. A. Baum, M. F. Frías, A. M. Haeberer, and P. E. M. López. From Specifications to Programs: a Fork Algebraic Approach to Bridge the Gap. In *Document 731-HKO-4, 47th Meeting of the IFIP Working Group 2.1 "Algorithmic Languages and Calculi"; Hong Kong*, January 1995.

[7] G. A. Baum, A. M. Haeberer, and P. A. S. Veloso. On the representability of the $\bigtriangledown$–abstract relational algebra. *IGPL Newsletter*, 1(3):3–4, 1992.

[8] R. Berghammer, A. M. Haeberer, G. Schmidt, and P. A. S. Veloso. Comparing Two Different Approaches to Products in Abstract Relation Algebras. In T. Rus M. Nivat, C. Rattray and G. Scollo, editors, *Algebraic Methodology and Software Technology (AMAST'93)* (Proceedings of the Third International Conference on Algebraic Methodology and Software Technology, University of Twente, The Netherlands, 21-25 June 1993), Springer–Verlag, London, 1993.

[9] C. H. Bergman, R. D. Maddux, and D. L. Pigozzi, editors. *Algebraic Logic and Universal Algebra in Computer Science*. Volume 425 of *Lecture Notes in Computer Science*, Springer–Verlag, Berlin, 1990. xi+292 pp.

[10] M. F. Frías, G. A. Baum, A. M. Haeberer, and P. A. S. Veloso. Fork Algebras are Representable. *Bulletin of the Section of Logic*, 24(2):64–75, 1995.

[11] M. F. Frías and S. Gordillo. Semantic Optimization of Queries in Deductive Object-oriented Database. In *Proc. of ADBIS'95*, Springer Verlag, Berlin, 1995. To appear.

[12] M. F. Frías, A. M. Haeberer, and P. A. S. Veloso. A Finite Axiomatization of Fork Algebras. *Bulletin of the Section of Logic*, 24(4):193–200, 1995.

[13] G. Grätzer. *Universal Algebra. Lecture Notes*, Springer Verlag, Berlin, 1979. 2nd edition.

[14] V. Gyuris. Representability of Fork Algebras. *Theoretical Computer Science*, 188(1–2):211–220, 1997.

[15] V. Gyuris. A Short Proof for Representability of Fork Algebras. *Logic Journal of the IGPL*, 3(5):791–796, 1995.

[16] A. M. Haeberer, G. A. Baum, and G. Schmidt. *Dealing with Non-constructive Specifications involving Quantifiers*. Technical Report, Pontifícia Universidade Católico do Rio de Janeiro, Dept. of Inf., 1993.

[17] A. M. Haeberer and P. A. S. Veloso. Partial Relations for Program Derivation: Adequacy, Inevitability and Expressiveness. In B. Möller, editor, *Constructing programs from specifications*, pages 319–371, 1991. Proc. of the IFIP TC2 Working Conference on Constructing Programs from Specifications, N.H., IFIP WG 2.1.

[18] L. Henkin, J. D. Monk, and A. Tarski. *Cylindric Algebras Part I.* North–Holland, Amsterdam, 1971 and 1985.

[19] L. Henkin, J. D. Monk, and A. Tarski. *Cylindric Algebras Part II.* North–Holland, Amsterdam, 1985.

[20] Á Kurucz. Decision Problems in Algebraic Logic. 1997.

[21] J. X. Madarász. Interpolation in Algebraizable Logics; Semantics for Non-normal Multi-modal Logic. *Journal of Applied Non-Classical Logics*, 8(1–2):67–105, 1998.

[22] R. D. Maddux. Finitary Algebraic Logic. *Zeitschrift f. Math. Logic u. Grundl. Math.*, 35:321–332, 1989.

[23] R. D. Maddux. Finitary Algebraic Logic, II. *Mathematical Logic Quarterly* (formerly: *Zeitschr. f. math. Logik und Grundlagen d. Math.*), 39:566–569, 1993.

[24] R. D. Maddux. Some Sufficient Conditions for Representability of Relation Algebras. *Algebra Universalis*, 8:162–172, 1978.

[25] M. Marx. *Algebraic Relativization and Arrow Logic*. PhD thesis, University of Amsterdam, appeared as ILLC Dissertation Series 1995–3. ILLC, Plantage Muidergracht 24, 1018 TV Amsterdam, e-mail: illc@fwi.uva.nl, 1995.

[26] M. Marx and Y. Venema. *Multi-Dimensional Modal Logic*. Volume 4 of *Applied Logic Series*, Kluwer Academic Publishers, 1997.

[27] R. N. McKenzie, G. F. McNulty, and W. F. Taylor. *Algebras, Lattices, Varieties, Vol. I*. The Wadsworth and Brooks/Cole Mathematics Series, Monterey, California, 1987.

[28] Sz. Mikulás, I. Sain, and A. Simon. Complexity of Equational Theory of Relational Algebras with Projection Elements. *Bulletin of the Section of Logic*, 21(3):103–111, 1992. Full paper is [39].

[29] I. Németi. *Algebraization of Quantifier Logics, An Introductory Overview. 12th version*. Technical Report 13/1996, Math. Inst. Hungar. Acad. Sci., Budapest, 1996. electronically available as: http://circle.math-inst.hu/pub/algebraic-logic/survey.dvi, survey.ps. An extended abstract appeared in *Studia Logica*, vol. 50 (3,4) (1991), pp. 485–569.

[30] I. Németi. Free Algebras and Decidability in Algebraic Logic. Dissertation with the Academy of Hungarian Sciences. Budapest, 1986. In Hungarian, xviii+169 pp.

[31] I. Németi. Ontology Can Turn Negative Results to Positive. *Bulletin of the Section of Logic*, 25(1):29–40, 1996.

[32] I. Németi. Strong Representability of Fork Algebras, A Set Theoretic Foundation. *Logic Journal of the IGPL*, 5(1):3–23, 1997.

[33] I. Németi and A. Simon. *Weakly Higher Order Cylindric Algebras and Finite Schema Axiomatization of the Representables*. Technical Report, Math. Inst. Hungar. Acad. Sci., Budapest, 1995.

[34] I. Sain. Beth's and Craig's Properties via Epimorphisms and Amalgamation in Algebraic Logic. In R. D. Maddux C. H. Bergman and D. L. Pigozzi, editors, *Algebraic Logic and Universal Algebra in Computer Science*, pages 209–226, Springer–Verlag, Berlin, 1990.

[35] I. Sain. A Finitizable Algebraization of First Order Logic. *This volume*, 2000. Abstracted in *Bulletin of the Section of Logic*, Univ. of Łódz, vol. 23(2) (1994), pp. 66–79. Extended abstract appeared as [36].

[36] I. Sain. On the Problem of Finitizing First Order Logic and its Algebraic Counterpart (A Survey of Results and Methods). In D. M. Gabbay L. Csirmaz and M. de Rijke, editors, *Logic Colloquium'92* (Proc. Veszprém, Hungary, 1992), pages 243–292, CSLI Publications, Stanford, 1995.

[37] I. Sain and V. Gyuris. Finite Schematizable Algebraic logic. *Logic Journal of the IGPL*, 5(5):699–751, 1997.

[38] I. Sain and I. Németi. Fork Algebras in Usual and in Non-well-founded Set Theories (An Overview). In E. Orlowska, editor, *Logic at Work (Essays Dedicated to the Memory of Helena Rasiowa)*, pages 669–694, Physica-Verlag, a Springer-Verlag Company, Heidelberg New York, 1999.

[39] I. Sain and A. Simon. *The Complexity of Equational Theory of Relational Algebras with Projection Elements*. Technical Report, Math. Inst. Hungar. Acad. Sci., Budapest, 1991.

[40] A. Simon. *A Purely Algebraic Proof of Representability of Quasi-projective Relation Algebras (*QRA*'s)*. Technical Report, Math. Inst. Hungar. Acad. Sci., Budapest, 1995.

[41] S. G. Simpson. Which Set Existence Axioms are Needed to Prove the Cauchy/ Peano Theorem? *Journal of Symbolic Logic*, 49:783–802, 1984.

[42] A. Tarski. On the Calculus of Relations. *Journal of Symbolic Logic*, 6:73–89, 1941.

[43] A. Tarski and S. Givant. *A Formalization of Set Theory Without Variables*. Volume 41 of *AMS Colloquium Publications*, Providence, Rhode Island, 1987.

[44] P. A. S. Veloso and A. M. Haeberer. A Finitary Relational Algebra for Classical First-order Logic. *Bulletin of the Section of Logic*, 20(2):52–62, June 1991.

[45] P. A. S. Veloso and A. M. Haeberer. A New Algebra of First-order Logic. In *Methodology and Philosophy of Science* (Proc. 9th International Congress on Logic, Upsala, Sweden), 1991.

View publication stats