

# MARKOV CHAIN APPROXIMATIONS TO SCALE FUNCTIONS OF LÉVY PROCESSES

ALEKSANDAR MIJATOVIĆ, MATIJA VIDMAR, AND SAUL JACKA

ABSTRACT. We introduce a general algorithm for the computation of the scale functions of a spectrally negative Lévy process  $X$ , based on a natural weak approximation of  $X$  via upwards skip-free continuous-time Markov chains with stationary independent increments. The algorithm consists of evaluating a finite linear recursion with, what are nonnegative, coefficients given explicitly in terms of the Lévy triplet of  $X$ . Thus it is easy to implement and numerically stable. Our main result establishes sharp rates of convergence of this algorithm providing an explicit link between the semimartingale characteristics of  $X$  and its scale functions, not unlike the one-dimensional Itô diffusion setting, where scale functions are expressed in terms of certain integrals of the coefficients of the governing SDE.

## 1. INTRODUCTION

It is well-known that, for a spectrally negative Lévy process  $X$  [5, Chapter VII] [33, Section 9.46], fluctuation theory in terms of the two families of scale functions,  $(W^{(q)})_{q \in [0, \infty)}$  and  $(Z^{(q)})_{q \in [0, \infty)}$ , has been developed [23, Section 8.2]. Of particular importance is the function  $W := W^{(0)}$ , in terms of which the others may be defined, and which features in the solution of many important problems of applied probability [22, Section 1.2]. It is central to these applications to be able to evaluate scale functions for any spectrally negative Lévy process  $X$ .

The goal of the present paper is to define and analyse a very simple novel algorithm for computing  $W$ . Specifically, to compute  $W(x)$  for some  $x > 0$ , choose small  $h > 0$  such that  $x/h$  is an integer. Then the approximation  $W_h(x)$  to  $W(x)$  is given by the recursion:

$$W_h(y+h) = W_h(0) + \sum_{k=1}^{y/h+1} W_h(y+h-kh) \frac{\gamma_{-kh}}{\gamma_h}, \quad W_h(0) = (\gamma_h h)^{-1} \quad (1.1)$$

for  $y = 0, h, 2h, \dots, x-h$ , where the coefficients  $\gamma_h$  and  $(\gamma_{-kh})_{k \geq 1}$  are expressible directly in terms of the Lévy measure  $\lambda$ , (possibly vanishing) Gaussian component  $\sigma^2$  and drift  $\mu$  of the Lévy process  $X$  (see (2.4)–(2.5) in Subsection 2.1 below).

---

2010 *Mathematics Subject Classification.* 60G51.

*Key words and phrases.* Spectrally negative Lévy processes, algorithm for computing scale functions, sharp convergence rates, continuous-time Markov chains.

The support of the Slovene Human Resources Development and Scholarship Fund under contract number 11010-543/2011 is acknowledged.

Indeed, the algorithm just described is based on a purely probabilistic idea of weak approximation: for small positive  $h$ ,  $X$  is approximated by what is a random walk  $X^h$  on a lattice with spacing  $h$ , skip-free to the right, and embedded into continuous time as a compound Poisson process (see Definition 3.1). Then, in recursion (1.1),  $W_h$  is the scale function associated to  $X^h$  — it plays a probabilistically analogous rôle for the process  $X^h$ , as does  $W$  for the process  $X$ . Thus  $W_h$  is computed as an approximation to  $W$  (see Corollary 3.1).

When it comes to existing methods for the evaluation of  $W$ , note that analytically  $W$  is characterized via its Laplace transform  $\widehat{W}$ ,  $\widehat{W}$  in turn being a certain rational function of the Laplace exponent  $\psi$  of  $X$ . However, already  $\psi$  need not be given directly in terms of elementary/special functions, and less often still is it possible to obtain closed-form expressions for  $W$  itself. The user is then faced with a Laplace inversion algorithm [13] [22, Chapter 5], which (i) necessarily involves the evaluation of  $\psi$ , typically at complex values of its argument and requiring high-precision arithmetic due to numerical instabilities; (ii) says little about the dependence of the scale function on the Lévy triplet of  $X$  (recall that  $\psi$  depends on a parametric complex integral of the Lévy measure, making it hard to discern how a perturbation in the Lévy measure influences the values taken by the scale function); and (iii) being a numerical approximation, fails *a priori* to ensure that the computed values of the scale function are probabilistically meaningful (e.g. given an output of a numerical Laplace inversion, it is not necessary that the formulae for, say, exit probabilities, involving  $W$ , should yield values in the interval  $[0, 1]$ ).

By contrast, it follows from (1.1) and the discussion following, that our proposed algorithm (i) requires no evaluations of the Laplace exponent of  $X$  and is numerically stable, as it operates in nonnegative real arithmetic [31, Theorem 7]; (ii) provides an explicit link between the deterministic semimartingale characteristics of  $X$ , in particular its Lévy measure, and the scale function  $W$ ; and (iii) yields probabilistically consistent outputs. Further, the values of  $W_h$  are so computed by a simple finite linear recursion and, as a by-product of the evaluation of  $W_h(x)$ , values  $W_h(y)$  for all the grid-points  $y = 0, h, 2h, \dots, x - h, x$ , are obtained (see Matlab code for the algorithm in [29]), which is useful in applications (see Section 6 below).

Our main results will (I) show that  $W_h$  converges to  $W$  pointwise, and uniformly on the grid with spacing  $h$  (if bounded away from 0 and  $+\infty$ ), for any spectrally negative Lévy process, and (II) establish sharp rates for this convergence under a mild assumption on the Lévy measure.

Due to the explicit connection between the coefficients appearing in (1.1) and the Lévy triplet of  $X$ , (1.1) also has the spirit of its one-dimensional Itô diffusion analogue, wherein the computation of the scale function requires numerical evaluation of certain integrals of the coefficients of the SDE driving said diffusion (for the explicit formulae of the integrals see e.g. [9, Chapters 2 and 3]). Indeed, we express  $W$  as a single limit, as  $h \downarrow 0$ , of nonnegative terms explicitly given in terms of the Lévy triplet. This is more direct than the Laplace inversion of a rational transform of the Laplace exponent, and hence may be of purely theoretical significance (see Remark 3.3 on how

the scale functions are affected by a perturbation of the Lévy measure, following directly from a transformed form of (1.1).

Finally, note that an algorithm, completely analogous to (1.1), for the computation of the scale functions  $W^{(q)}$ , and also  $Z^{(q)}$ ,  $q \geq 0$ , follows from our results (see Corollary 3.1, Eq. (3.3) and (3.4)) and presents no further difficulty for the analysis of convergence (see Theorem 1.1 below). Indeed, our discretization allows naturally to approximate other quantities involving scale functions, which arise in application: the derivatives of  $W^{(q)}$  by difference quotients of  $W_h^{(q)}$ ; the integrals of a continuous (locally bounded) function against  $dW^{(q)}$  by its integrals against  $dW_h^{(q)}$ ; expressions of the form  $\int_0^x F(y, W^{(q)}(y))dy$ , where  $F$  is continuous locally bounded, by the sums  $\sum_{k=0}^{\lfloor x/h \rfloor - 1} F(kh, W_h^{(q)}(kh))h$  etc. (See Section 6 for examples.)

**1.1. Overview of main results.** The key idea leading to the algorithm in (1.1) is best described by the following two steps: (i) approximate the spectrally negative Lévy process  $X$  by a continuous-time Markov chain (CTMC)  $X^h$  with state space  $\mathbb{Z}_h := \{hk: k \in \mathbb{Z}\}$  ( $h \in (0, h_*)$  for some  $h_* > 0$ ), as described in Subsection 2.1; (ii) find an algorithm for computing the scale functions of the chain  $X^h$ . The approximation in Subsection 2.1 implies that  $X^h$  is a compound Poisson (CP) process, which is not spectrally negative. However, since the corresponding jump chain of  $X^h$  is a skip-free to the right  $\mathbb{Z}_h$ -valued random walk, it is possible to introduce (right-continuous, nondecreasing) scale functions  $(W_h^{(q)})_{q \geq 0}$  and  $(Z_h^{(q)})_{q \geq 0}$  (with measures  $dW_h^{(q)}$  and  $dZ_h^{(q)}$  supported in  $\mathbb{Z}_h$ ), in analogy to the spectrally negative case. Moreover, as described in Corollary 3.1, a straightforward recursive algorithm is readily available for evaluating *exactly* any function in the families  $(W_h^{(q)})_{q \geq 0}$  and  $(Z_h^{(q)})_{q \geq 0}$  at any point. More precisely, it emerges, that for each  $x \in \mathbb{Z}_h$ ,  $W_h^{(q)}(x)$  (resp.  $Z_h^{(q)}(x)$ ) obtains as a *finite* linear combination of the preceding values  $W_h^{(q)}(y)$  (resp.  $Z_h^{(q)}(y)$ ) for  $y \in \{0, h, \dots, x - h\}$ ; with the starting value  $W_h^{(q)}(0)$  (resp.  $Z_h^{(q)}(0)$ ) being known explicitly. This is in spite of the fact that the state space of the Lévy process  $X^h$  is in fact the *infinite* lattice  $\mathbb{Z}_h$ .

In order to precisely describe the rates of convergence of the algorithm in (1.1), we introduce some notation. Fix  $q \geq 0$  and define for  $K, G$  bounded subset of  $(0, \infty)$ :

$$\Delta_W^K(h) := \sup_{x \in \mathbb{Z}_h \cap K} \left| W_h^{(q)}(x - \delta^0 h) - W^{(q)}(x) \right| \quad \text{and} \quad \Delta_Z^G(h) := \sup_{x \in \mathbb{Z}_h \cap G} \left| Z_h^{(q)}(x) - Z^{(q)}(x) \right|,$$

where  $\delta^0$  equals 0 if  $X$  has sample paths of finite variation and 1 otherwise. We further introduce:

$$\kappa(\delta) := \int_{[-1, -\delta)} |y| \lambda(dy), \quad \text{for any } \delta \geq 0.$$

If the jump part of  $X$  has paths of infinite variation, i.e. in the case the equality  $\kappa(0) = \infty$  holds, we assume (throughout the paper we shall make it explicit when this assumption is in effect):

**Assumption 1.1.** *There exists  $\epsilon \in (1, 2)$  with:*

- (1)  $\limsup_{\delta \downarrow 0} \delta^\epsilon \lambda(-1, -\delta) < \infty$  and
- (2)  $\liminf_{\delta \downarrow 0} \int_{[-\delta, 0)} x^2 \lambda(dx) / \delta^{2-\epsilon} > 0$ .

Note that this is a fairly mild condition, fulfilled if e.g.  $\lambda(-1, -\delta)$  “behaves as”  $\delta^{-\epsilon}$ , as  $\delta \downarrow 0$ ; for a precise statement see Remark 5.9.

Here is now our main result:

**Theorem 1.1.** *Let  $K$  and  $G$  be bounded subsets of  $(0, \infty)$ ,  $K$  bounded away from zero when  $\sigma^2 = 0$ . If  $\kappa(0) = \infty$ , suppose further that Assumption 1.1 is fulfilled. Then the rates of convergence of the scale functions are summarized by the following table:*

$\lambda(\mathbb{R}) = 0$	$\Delta_W^K(h) = O(h^2)$ and $\Delta_Z^G(h) = O(h)$
$0 < \lambda(\mathbb{R}) \notin \kappa(0) < \infty$	$\Delta_W^K(h) + \Delta_Z^G(h) = O(h)$
$\kappa(0) = \infty$	$\Delta_W^K(h) + \Delta_Z^G(h) = O(h^{2-\epsilon})$

Moreover, the rates so established are sharp in the sense that for each of the three entries in the table above, examples of spectrally negative Lévy processes are constructed for which the rate of convergence is no better than stipulated.

*Remark 1.1.* (1) The rates of convergence depend on the behaviour of the tail of the Lévy measure at the origin; by contrast behaviour of Laplace inversion algorithms tends to be susceptible to the degree of smoothness of the scale function (for which see [12]) itself [1].

(2) More exhaustive and at times general statements are to be found in Propositions 5.5–5.11. In particular, the case  $\sigma^2 > 0$  and  $\kappa(0) = \infty$  does not require Assumption 1.1 to be fulfilled, although the statement of the convergence rate is more succinct under its proviso.

(3) The proof of Theorem 1.1 consists of studying the differences of the integral representations of the scale functions. The integrands decay only according to some power law, making the analysis much more involved than was the case in [30], where the corresponding decay was exponential. In particular, one cannot in the pure-jump case directly apply the integral triangle inequality. The structure of the proof is explained in detail in Subsection 5.1.

(4) Since scale functions often appear in applications (for which see Section 1.2 below) in the form  $W^{(q)}(x)/W^{(q)}(y)$  ( $x, y > 0$ ,  $q \geq 0$ ), we note that the rates from Theorem 1.1 transfer directly to such quotients, essentially because  $W_h^{(q)}(y) \rightarrow W^{(q)}(y) \in (0, \infty)$ , as  $h \downarrow 0$ , and since for all  $h \in (0, h_*)$ ,  $\frac{1}{W^{(q)}(y)} - \frac{1}{W_h^{(q)}(y)} = \frac{W_h^{(q)}(y) - W^{(q)}(y)}{W^{(q)}(y)W_h^{(q)}(y)}$ .

(5) For a result concerning the derivatives of  $W^{(q)}$  see Subsection 5.10.

**1.2. Overview of the literature and of the applications of scale functions.** For the general theory of spectrally negative Lévy processes and their scale functions we refer to [23, Chapter 8] and [5, Chapter VII], while an excellent account of available numerical methods for computing them can be found in [22, Chapter 5]. Examples, few, but important, of processes when the scale functions *can* be given analytically, appear e.g. in [18]; and in certain cases it is possible to construct them *indirectly* [22, Chapter 4] (i.e. not starting from the basic datum, which we consider here to be the characteristic triplet of  $X$ ). Finally, in the special case when  $X$  is a positive drift minus a compound Poisson subordinator, we note that numerical schemes for (finite time) ruin/survival probabilities

(expressible in terms of scale functions), based on discrete-time Markov chain approximations of one sort or another, have been proposed in the literature (see [36, 16, 11, 15] and the references therein).

In terms of applications of scale functions in applied probability, there are numerous identities concerning boundary crossing problems and related path decompositions in which scale functions feature [22, p. 5]. They do so either (a) indirectly (usually as Laplace transforms of quantities which are ultimately of interest), or even (b) directly (then typically, but not always, as probabilities in the form of quotients  $W(x)/W(y)$ ). For examples of the latter see the two-sided exit problem [5, Chapter VII, Theorem 8]; ruin probabilities [23, p. 217, Eq. (8.15)] and the Gerber-Shiu measure [24, Section 5.4] in the insurance/ruin theory context; laws of suprema of continuous-state branching processes [8, Proposition 3.1]; Lévy measures of limits of continuous-state branching processes with immigration (CBI processes) [21, Eq. (3.7)]; laws of branch lengths in population biology [25, Eq. (7)]; the Shepp-Shiryaev optimal stopping problem (solved for the spectrally negative case in [4, Theorem 2, Eq. (30)]); [27, Proposition 1] for an optimal dividend control problem. A further overview of these and other applications of scale functions (together with their derivatives and the integrals  $Z^{(q)}$ ), e.g. in queuing theory and fragmentation processes, may be found in [22, Section 1.2], see also the references therein. A suite of identities involving Laplace transforms of quantities pertaining to the reflected process of  $X$  appears in [28].

**1.3. Organisation of the remainder of the paper.** Section 2 gives the setting and fixes general notation. Section 3 introduces upwards skip-free Lévy chains (they being the continuous-time analogues of random walks, which are skip-free to the right), describes their scale functions and how to compute them. In Section 4 we demonstrate pointwise convergence of the approximating scale functions to those of the spectrally negative Lévy process. Then Section 5 establishes the rate at which this convergence transpires. Finally, Section 6 provides some numerical illustrations and further discusses the computational side of the proposed algorithm. Appendices A and B contain the proofs of technical results from Subsection 5.2, while Appendix C provides some additional numerical examples.

## 2. SETTING AND GENERAL NOTATION

Throughout this paper we let  $X$  be a spectrally negative Lévy process (i. e.  $X$  has stationary independent increments, is càdlàg,  $X_0 = 0$  a.s., the Lévy measure  $\lambda$  of  $X$  is concentrated on  $(-\infty, 0)$  and  $X$  does not have a.s. monotone paths). The Laplace exponent  $\psi$  of  $X$ , defined via  $\psi(\beta) := \log \mathbf{E}[e^{\beta X_1}]$  ( $\beta \in \{\gamma \in \mathbb{C} : \Re \gamma \geq 0\} =: \overline{\mathbb{C}^+}$ ), can be expressed as (see e.g. [5, p. 188]):

$$\psi(\beta) = \frac{1}{2}\sigma^2\beta^2 + \mu\beta + \int_{(-\infty, 0)} \left( e^{\beta y} - \beta\tilde{c}(y) - 1 \right) \lambda(dy), \quad \beta \in \overline{\mathbb{C}^+}. \quad (2.1)$$

The Lévy triplet of  $X$  is thus given by  $(\sigma^2, \lambda, \mu)_{\tilde{c}}$ ,  $\tilde{c} := \text{id}_{\mathbb{R}} \mathbb{1}_{[-V, 0)}$  with  $V$  equal to either 0 or 1, the former only if  $\int_{[-1, 0)} |x|\lambda(dx) < \infty$ . Further, when the Lévy measure satisfies  $\int_{[-1, 0)} |x|\lambda(dx) < \infty$ ,

we may always express  $\psi$  in the form  $\psi(\beta) = \frac{1}{2}\sigma^2\beta^2 + \mu_0\beta + \int_{(-\infty,0)} (e^{\beta y} - 1) \lambda(dy)$  for  $\beta \in \overline{\mathbb{C}^{\rightarrow}}$ . If in addition  $\sigma^2 = 0$ , then necessarily the drift  $\mu_0$  must be strictly positive,  $\mu_0 > 0$  [23, p. 212].

**2.1. The approximation.** We now recall from [30], specializing to the spectrally negative setting, the spatial discretisation of  $X$  by the family of CTMCs  $(X^h)_{h \in (0, h_*)}$  (where  $h_* \in (0, +\infty]$ ). This family weakly approximates  $X$  as  $h \downarrow 0$ . As in [30] we will use two approximating schemes, scheme 1 and 2, according as  $\sigma^2 > 0$  or  $\sigma^2 = 0$ . Recall that two different schemes are introduced since the case  $\sigma^2 > 0$  allows for a better (i.e. a faster converging) discretization of the drift term, but the case  $\sigma^2 = 0$  (in general) does not [30, Paragraph 2.2.1]. Let also  $V = 0$ , if  $\lambda$  is finite and  $V = 1$ , if  $\lambda$  is infinite. Notation-wise, define for  $h > 0$ ,  $c_y^h := \lambda(A_y^h)$  with  $A_y^h := [y - h/2, y + h/2)$  ( $y \in \mathbb{Z}_h^- := \mathbb{Z}_h \cap (-\infty, 0)$ );  $A_0^h := [-h/2, 0)$ ;

$$c_0^h := \int_{A_0^h} y^2 \mathbb{1}_{[-V,0)}(y) \lambda(dy) \quad \text{and} \quad \mu^h := \sum_{y \in \mathbb{Z}_h^-} y \int_{A_y^h} \mathbb{1}_{[-V,0)}(z) \lambda(dz).$$

Now, it is clear from [30] that, for each  $h \in (0, h_*)$ ,  $X^h$  is a CP process, with  $X_0^h = 0$ , a.s., and whose positive jumps do not exceed  $h$ . Thus each  $X^h$  admits its Laplace exponent  $\psi^h(\beta) := \log \mathbb{E}[e^{\beta X_1^h}]$  ( $\beta \in \overline{\mathbb{C}^{\rightarrow}}$ ), which in turn uniquely determines its law. Moreover,  $\psi^h$  may be obtained from the characteristic exponent by analytic continuation, and we have from [30, Eq. (3.1) and (3.2)] (for  $\beta \in \overline{\mathbb{C}^{\rightarrow}}$ ) under scheme 1,

$$\psi^h(\beta) = (\mu - \mu^h) \frac{e^{\beta h} - e^{-\beta h}}{2h} + (\sigma^2 + c_0^h) \frac{e^{\beta h} + e^{-\beta h} - 2}{2h^2} + \sum_{y \in \mathbb{Z}_h^-} c_y^h (e^{\beta y} - 1), \quad (2.2)$$

and under scheme 2,

$$\psi^h(\beta) = (\mu - \mu^h) \frac{e^{\beta h} - 1}{h} + c_0^h \frac{e^{\beta h} + e^{-\beta h} - 2}{2h^2} + \sum_{y \in \mathbb{Z}_h^-} c_y^h (e^{\beta y} - 1). \quad (2.3)$$

To properly appreciate where the different terms appearing in these Laplace exponents come from, we refer the reader to our paper [30], especially Section 2.1 therein.

Lévy measure/diffusion part	$\sigma^2 > 0$	$\sigma^2 = 0$
$\lambda(\mathbb{R}) < \infty$	$V = 0$ , scheme 1	$V = 0$ , scheme 2
$\lambda(\mathbb{R}) = \infty$	$V = 1$ , scheme 1	$V = 1$ , scheme 2

TABLE 1. The usage of schemes 1 and 2 and of  $V$  depends on the nature of  $\sigma^2$  and  $\lambda$ .

Indeed, note that, starting directly from [30, Eq. (3.2)], the term  $(\mu - \mu^h) \frac{e^{\beta h} - 1}{h}$  in (2.3) should actually read as:

$$(\mu - \mu^h) \left( \frac{e^{\beta h} - 1}{h} \mathbb{1}_{[0, \infty)}(\mu - \mu^h) + \frac{1 - e^{-\beta h}}{h} \mathbb{1}_{(-\infty, 0]}(\mu - \mu^h) \right).$$

However, when  $X$  is a spectrally negative Lévy process with  $\sigma^2 = 0$ , we have  $\mu - \mu^h \geq 0$ , at least for all sufficiently small  $h$ . Indeed, if  $\int_{[-1,0)} |y|\lambda(dy) < \infty$ , then  $\mu_0 > 0$  and by dominated convergence  $\mu - \mu^h \rightarrow \mu_0$  as  $h \downarrow 0$ . On the other hand, if  $\int_{[-1,0)} |y|\lambda(dy) = \infty$ , then we deduce by monotone convergence  $-\mu^h \geq \frac{1}{2} \int_{[-1,-h/2)} |y|\lambda(dy) \rightarrow \infty$  as  $h \downarrow 0$ . We shall assume throughout that  $h_\star$  is already chosen small enough, so that  $\mu - \mu^h \geq 0$  holds for all  $h \in (0, h_\star)$ .

In summary, then,  $h_\star$  is chosen so small as to guarantee that, for all  $h \in (0, h_\star)$ : (i)  $\mu - \mu^h \geq 0$  and (ii)  $\psi^h$  is the Laplace exponent of some CP process  $X^h$ , which is also a CTMC with state space  $\mathbb{Z}_h$  (note that in [30, Proposition 3.9] it is shown  $h_\star$  can indeed be so chosen, viz. point (ii)). Eq. (2.2) and (2.3) then determine the *weak* approximation  $(X^h)_{h \in (0, h_\star)}$  precisely. Finally, for  $h \in (0, h_\star)$ , let  $\lambda^h$  denote the Lévy measure of  $X^h$ . In particular,  $\psi^h(\beta) = \int (e^{\beta y} - 1) \lambda^h(dy)$ ,  $\beta \in \overline{\mathbb{C}^-}$ ,  $h \in (0, h_\star)$ , so that the jump intensities, equivalently the Lévy measure, of  $X^h$  can be read off directly from (2.2)-(2.3). It will also be convenient to define  $\psi^0 := \psi$ .

We conclude this subsection by expressing explicitly the coefficients of the linear recursion in (1.1) in terms of the Lévy triplet of  $X$ . Define:

$$\tilde{\sigma}_h^2 := \frac{1}{2h^2} (\sigma^2 + c_0^h), \quad \tilde{\mu}^h := \frac{1}{2h} (\mu - \mu^h),$$

and note that  $\tilde{\mu}^h$  is non-negative for  $h \in (0, h_\star)$ . Recall that  $V$  equals 0 or 1 according as to whether  $\lambda$  is finite or infinite and note that, if  $V = 0$ , we have  $\tilde{\sigma}_h^2 = \sigma^2/2h^2$  and  $\tilde{\mu}^h = \mu/2h$ . We can now define the coefficients in (1.1) by:

$$\gamma_h := \tilde{\sigma}_h^2 + \mathbb{1}_{(0,\infty)}(\sigma^2) \tilde{\mu}^h + \mathbb{1}_{\{0\}}(\sigma^2) 2\tilde{\mu}^h, \quad \gamma_{-h} := \tilde{\sigma}_h^2 - \mathbb{1}_{(0,\infty)}(\sigma^2) \tilde{\mu}^h + \lambda(-\infty, -h/2] \quad (2.4)$$

$$\gamma_{-kh} := \lambda(-\infty, -kh + h/2], \quad \text{where } k \geq 2. \quad (2.5)$$

**2.2. Connection with integro-differential equations.** Note that an alternative representation of (1.1) (as generalized to the case of arbitrary  $q \geq 0$ ; see (3.1) of Proposition 3.2) is the analogue of the relation  $(L - q)W^{(q)} = 0$  in the spectrally negative case, the latter holding true under sufficient regularity conditions on  $W^{(q)}$  (see e.g. [7, Eq. (12)]). This suggests there might be a link between our probabilistic approximation and solutions to integro-differential equations.

Indeed, one can check, for each  $q \geq 0$ , by taking Laplace transforms (using (2.1), the expression for the Laplace transform of  $W^{(q)}$  [22, p. 100, Eq. (4)], nondecreasingness of  $W^{(q)}$ , and the theorem of Fubini), that the function  $W^{(q)}$  satisfies the following integro-differential equation (cf. [3, Corollary IV.3.3] for the case  $\sigma^2 = 0$ ,  $\lambda(\mathbb{R}) < +\infty$ , and survival probabilities):

$$\frac{1}{2\sigma^2} \frac{dW^{(q)}}{dx} = 1 - \mu W^{(q)}(x) + \int_0^\infty \left( W^{(q)}(x-y) (\lambda(-\infty, -y) + q) - W^{(q)}(x) \lambda[-V, -y] \mathbb{1}_{(0,V]}(y) \right) dy \quad (2.6)$$

(for the value of  $W^{(q)}(0)$  see [22, p. 127, Lemma 3.1]). Note that in the last integral of (2.6), the two terms appearing in its integrand cannot be separated when  $\kappa(0) = +\infty$ .

Furthermore, (1.1) as generalized to arbitrary  $q \geq 0$  (see Corollary 3.1, Eq. 3.3) can be rewritten, when  $\sigma^2 = 0$ , as (for  $x \in \mathbb{Z}_h^{++}$ ):

$$\frac{1}{2c_0^h} \frac{W_h^{(q)}(x) - W_h^{(q)}(x-h)}{h} = 1 + (\mu^h - \mu) W_h^{(q)}(x) + h \sum_{k=1}^{x/h} W_h^{(q)}(x-kh) (\lambda(-\infty, -(k-1/2)h) + q), \quad (2.7)$$

with  $W_h^{(q)}(0) = \frac{1}{\frac{1}{2}c_0^h/h + \mu - \mu^h}$ , and when  $\sigma^2 > 0$ , as (again for  $x \in \mathbb{Z}_h^{++}$ ):

$$\begin{aligned} \frac{1}{2}(\sigma^2 + c_0^h) \frac{W_h^{(q)}(x) - W_h^{(q)}(x-h)}{h} &= 1 + (\mu^h - \mu) \frac{W_h^{(q)}(x) + W_h^{(q)}(x-h)}{2} \\ &\quad + h \sum_{k=1}^{x/h} W_h^{(q)}(x-kh) (\lambda(-\infty, -(k-1/2)h) + q), \end{aligned} \quad (2.8)$$

with  $W_h^{(q)}(0) = \frac{2h}{\sigma^2 + c_0^h + (\mu - \mu^h)h}$ .

Thus (2.7) and (2.8) can be seen as (simple) approximation schemes for the integro-differential equation (2.6).<sup>1</sup> However, from this viewpoint alone, it would be very difficult indeed to “guess at” the correct discretization, which would also yield meaningful generalized scale functions of approximating chains — the latter being our starting point and precisely the aspect of our schemes, which we wish to emphasize. Indeed, higher-order schemes for (2.6), if and should they exist, would (likely) no longer be connected with Lévy chains.

**2.3. General notation.** With regard to miscellaneous notation, we let  $\mathbb{R}_+$  (respectively  $\mathbb{R}^+$ ) be the nonnegative (respectively strictly positive) real numbers;  $\mathbb{Z}_+ = \mathbb{N}_0$  (respectively  $\mathbb{Z}^+$ ,  $\mathbb{Z}_-$ ,  $\mathbb{Z}^-$ ) the nonnegative, (respectively strictly positive, nonpositive, strictly negative) integers;  $\mathbb{Z}_h^+$  (respectively  $\mathbb{Z}_h^{++}$ ,  $\mathbb{Z}_h^-$ ,  $\mathbb{Z}_h^{--}$ ) the nonnegative (respectively strictly positive, nonpositive, strictly negative) elements of  $\mathbb{Z}_h := \{hk : k \in \mathbb{Z}\}$ ;  $\mathbb{C}^{\leftarrow} := \{z \in \mathbb{C} : \Re z < 0\}$  (respectively  $\mathbb{C}^{\rightarrow} := \{z \in \mathbb{C} : \Re z > 0\}$ ) with  $\overline{\mathbb{C}^{\leftarrow}}$  (respectively  $\overline{\mathbb{C}^{\rightarrow}}$ ) denoting the closure of this set (note that the arrow notation is suggestive of which halfplane is being considered). To a nondecreasing right-continuous function  $F : \mathbb{R} \rightarrow \mathbb{R}$  a measure  $dF$  may be associated in the Lebesgue-Stieltjes sense; stating that  $F$  is of  $M$ -exponential order means  $|F|(x) \leq Ce^{Mx}$  for all  $x \geq 0$ , for some  $C < \infty$ . On the other hand, a real-valued function  $f$ , defined on some  $(a, \infty)$ ,  $a \in \mathbb{R}$  is said to grow asymptotically  $M$ -exponentially,  $M \in (0, \infty)$ , if  $\{\liminf_{x \rightarrow \infty} f(x)e^{-Mx}, \limsup_{x \rightarrow \infty} f(x)e^{-Mx}\} \subset (0, \infty)$ . Further, for functions  $g \geq 0$  and  $h > 0$  defined on some right neighborhood of 0,  $g \sim h$  (resp.  $g = O(h)$ ,  $g = o(h)$ ) means  $\lim_{0+} g/h \in (0, \infty)$  (resp.  $\limsup_{0+} g/h < \infty$ ,  $\lim_{0+} g/h = 0$ ). Next, the Laplace transform of a measurable function  $f : \mathbb{R} \rightarrow \mathbb{R}$  of  $M$ -exponential order, with  $f|_{(-\infty, 0)}$  constant (respectively measure  $\mu$  on  $\mathbb{R}$ , concentrated on  $[0, \infty)$ ) is denoted  $\hat{f}$  (respectively  $\hat{\mu}$ ):  $\hat{f}(\beta) = \int_0^\infty e^{-\beta x} f(x) dx$  for  $\Re \beta > M$  (respectively  $\hat{\mu}(\beta) = \int_{[0, \infty)} e^{-\beta x} \mu(dx)$  for all  $\beta \geq 0$  such that this integral is finite). For  $\{x, y\} \subset \mathbb{R}$ ,  $[x] := \max\{k \in \mathbb{Z} : k \leq x\}$ ,  $x \vee y = \max\{x, y\}$  and  $x \wedge y = \min\{x, y\}$ . A sequence  $(h_n)_{n \in \mathbb{N}}$  of non-zero real numbers is said to be nested, if  $h_n/h_{n+1} \in \mathbb{N}$  for all  $n \in \mathbb{N}$ .  $\delta_x$  denotes the Dirac measure at  $x \in \mathbb{R}$ . Finally, increasing will mean strictly increasing; DCT stands for the Dominated Convergence Theorem; and we interpret  $\pm a/0 = \pm \infty$  for  $a > 0$ .

<sup>1</sup>We are grateful to an anonymous referee for pointing out this connection, on an earlier draft of this paper.



## 3. UPWARDS SKIP-FREE LÉVY CHAINS AND THEIR SCALE FUNCTIONS

In the sequel, we will require a fluctuation theory (and, in particular, a theory of scale functions) for random walks, which are skip-free to the right, once these have been embedded into continuous-time as CP processes (see next definition and remark). Indeed, this theory has been developed in full detail in [35] and we recall here for the readers convenience the pertinent results.

**Definition 3.1.** A Lévy process  $Y$  with Lévy measure  $\nu$  is said to be an *upwards skip-free Lévy chain*, if it is compound Poisson and the Lévy measure  $\nu$  satisfies  $\text{supp}(\nu) \subset \mathbb{Z}_h$  and  $\text{supp}(\nu|_{\mathcal{B}((0,\infty))}) = \{h\}$  for some  $h > 0$ .

*Remark 3.1.* For all  $h \in (0, h_*)$ ,  $X^h$  is an upwards skip-free Lévy chain.

For the remainder of this subsection we let  $Y$  be an upwards skip-free Lévy chain with Lévy measure  $\nu$ , such that  $\nu(\{h\}) > 0$  ( $h > 0$ ).

The following is either clear or else can be found in [35, Subsection 3.1]:

- (i) One can introduce the Laplace exponent  $\varphi : \overline{\mathbb{C}^\rightarrow} \rightarrow \mathbb{C}$ , given by  $\varphi(\beta) := \int_{\mathbb{R}} (e^{\beta x} - 1)\nu(dx)$  ( $\beta \in \overline{\mathbb{C}^\rightarrow}$ ), for which:  $\mathbf{E}[e^{\beta Y_t}] = \exp\{t\varphi(\beta)\}$  ( $\beta \in \overline{\mathbb{C}^\rightarrow}$ ,  $t \geq 0$ ).  $\varphi$  is continuous in  $\overline{\mathbb{C}^\rightarrow}$ , analytic in  $\mathbb{C}^\rightarrow$ ,  $\lim_{+\infty} \varphi|_{[0,\infty)} = +\infty$  with  $\varphi|_{[0,\infty)}$  strictly convex.
- (ii) Let  $\Phi(0) \in [0, \infty)$  be the largest root of  $\varphi$  on  $[0, \infty)$ . Then  $\varphi|_{[\Phi(0), \infty)} : [\Phi(0), \infty) \rightarrow [0, \infty)$  is an increasing bijection and we let  $\Phi := (\varphi|_{[\Phi(0), \infty)})^{-1} : [0, \infty) \rightarrow [\Phi(0), \infty)$  be its inverse.

We introduce in the next proposition two families of scale functions for  $Y$ , which play analogous roles in the solution of exit problems, as they do in the case of spectrally negative Lévy processes, see [35, Subsections 4.1-4-3]:

**Proposition 3.1** (Scale functions). *There exists a family of functions  $W^{(q)} : \mathbb{R} \rightarrow [0, \infty)$  and*

$$Z^{(q)}(x) = 1 + q \int_0^{\lfloor x/h \rfloor h} W^{(q)}(y) dy, \quad x \in \mathbb{R}$$

*defined for each  $q \geq 0$  such that for any  $q \geq 0$ , we have  $W^{(q)}(x) = 0$  for  $x < 0$  and  $W^{(q)}$  is characterised on  $[0, \infty)$  as the unique right-continuous and piecewise constant function of exponential order whose Laplace transform satisfies:*

$$\widehat{W^{(q)}}(\beta) = \frac{e^{\beta h} - 1}{\beta h(\varphi(\beta) - q)} \text{ for } \beta > \Phi(q).$$

*Remark 3.2.* (i) The functions  $W^{(q)}$  are nondecreasing and the corresponding measures  $dW^{(q)}$  are supported in  $\mathbb{Z}_h$  for each  $q \geq 0$ .

(ii) The Laplace transform of the functions  $Z^{(q)}$  is given by:

$$\widehat{Z^{(q)}}(\beta) = \frac{1}{\beta} \left( 1 + \frac{q}{\varphi(\beta) - q} \right) \text{ for } \beta > \Phi(q), q \geq 0.$$

(iii) For all  $q \geq 0$ :  $W^{(q)}(0) = 1/(h\nu(\{h\}))$ .

Finally, the following proposition, whose corollary will give rise to a method for calculating the values of the scale functions associated to  $Y$ , follows from the strong Markov property of  $Y$  (see [35, Subsection 4.4]).

**Proposition 3.2.** *Let  $P$  be the transition matrix of the jump chain of the CTMC  $Y$  and let  $q \geq 0$ ,  $\nu_q := 1 + q/\nu(\mathbb{R})$ . Assume (for ease of notation and without loss of generality) that  $h = 1$ . Then, seen as vectors,  $W^{(q)} := (W^{(q)}(k))_{k \in \mathbb{Z}}$  and  $Z^{(q)} := (Z^{(q)}(k))_{k \in \mathbb{Z}}$  satisfy, entry-by-entry:*

$$(PW^{(q)})|_{\mathbb{Z}_+} = \nu_q W^{(q)}|_{\mathbb{Z}_+} \text{ and } (PZ^{(q)})|_{\mathbb{Z}_+} = \nu_q Z^{(q)}|_{\mathbb{Z}_+},$$

*i.e. with  $\gamma := \nu(\mathbb{R})$ ,  $p := \nu(\{1\})/\gamma$  and  $q_n := \nu(\{-n\})/\gamma$  ( $n \in \mathbb{N}$ ), the recursive relations (for  $n \in \mathbb{N} \cup \{0\}$ ):*

$$pW^{(q)}(n+1) = \left(1 + \frac{q}{\gamma}\right) W^{(q)}(n) - \sum_{k=1}^n q_k W^{(q)}(n-k), \quad (3.1)$$

and

$$pZ^{(q)}(n+1) + \left(1 - p - \sum_{k=1}^{n-1} q_k\right) = \left(1 + \frac{q}{\gamma}\right) Z^{(q)}(n) - \sum_{k=1}^{n-1} q_k Z^{(q)}(n-k) \quad (3.2)$$

*hold true. Additionally  $W^{(q)}|_{\mathbb{Z}_-} = 0$  with  $W^{(q)}(0) = 1/\nu(\{1\})$ , whereas  $Z^{(q)}|_{\mathbb{Z}_-} = 1$ .*

**Corollary 3.1.** *We have for all  $n \in \mathbb{N} \cup \{0\}$ :*

$$W^{(q)}(n+1) = W^{(q)}(0) + \sum_{k=1}^{n+1} W^{(q)}(n+1-k) \frac{q + \nu(-\infty, -k]}{\nu(\{1\})}, \quad W^{(q)}(0) = 1/\nu(\{1\}), \quad (3.3)$$

and for  $\widetilde{Z}^{(q)} := Z^{(q)} - 1$ ,

$$\widetilde{Z}^{(q)}(n+1) = (n+1) \frac{q}{\nu(\{1\})} + \sum_{k=1}^n \widetilde{Z}^{(q)}(n+1-k) \frac{q + \nu(-\infty, -k]}{\nu(\{1\})}, \quad \widetilde{Z}^{(q)}(0) = 0. \quad (3.4)$$

*Remark 3.3.* Whilst we have based our algorithm on Eq. (3.3) and (3.4), nevertheless Eq. (3.1) and (3.2) should not be discounted entirely. For example, they allow to make the following observation.

Recall  $W_h$  is given by (3.3), as applied to the process  $Y = X^h/h$ , equivalently then it may be obtained from (3.1), again as applied to the process  $Y = X^h/h$ . Assume next (by scaling, without loss of generality)  $V \leq x \in \mathbb{Z}_h$ .

Suppose the Lévy measure  $\lambda$  of  $X$  is modified below (possibly including) the level  $-x$  in such a way that  $\lambda(-\infty, -x]$  is preserved, whilst  $\mathbb{1}_{(-x,0)} \cdot \lambda$ ,  $\sigma^2$  and  $\mu$  are kept fixed. Then (3.1), as applied to  $X^h/h$  for computing  $W_h$  on the interval  $[0, x]$ , remains unaffected. This is because this recursion up to level  $x$  depends solely on the probabilities of the jump-sizes with modulus at most  $x-h$  and on the total Lévy mass of the approximating Lévy chain. The latter, however, do not change by said transformation of the Lévy measure (see (2.2)-(2.3)). Consequently, since, as we shall see,  $W_h$  converges to  $W$ , as  $h \downarrow 0$ , this also means that the scale function  $W$  itself on the interval  $[0, x]$  is invariant under such a transformation. Analogously for  $W^{(q)}$  and  $Z^{(q)}$ .

## 4. CONVERGENCE OF SCALE FUNCTIONS

First we fix some notation. Pursuant to [23, Subsections 8.1 & 8.2] (respectively Section 3) we associate henceforth with  $X$  (respectively  $X^h$ ) two families of scale functions  $(W^{(q)})_{q \geq 0}$  and  $(Z^{(q)})_{q \geq 0}$  (respectively  $(W_h^{(q)})_{q \geq 0}$  and  $(Z_h^{(q)})_{q \geq 0}$ ,  $h \in (0, h_*)$ ). Note that these functions are defined on the whole of  $\mathbb{R}$ , are nondecreasing, càdlàg, with  $W^{(q)}(x) = W_h^{(q)}(x) = 0$  and  $Z^{(q)}(x) = Z_h^{(q)}(x) = 1$  for  $x \in (-\infty, 0)$ . We also let  $\Phi(0)$  (respectively  $\Phi^h(0)$ ) be the largest root of  $\psi|_{[0, \infty)}$  (respectively  $\psi^h|_{[0, \infty)}$ ) and denote by  $\Phi$  (respectively  $\Phi^h$ ) the inverse of  $\psi|_{[\Phi(0), \infty)}$  (respectively  $\psi^h|_{[\Phi^h(0), \infty)}$ ,  $h \in (0, h_*)$ ). As usual  $W$  (resp.  $W_h$ ) denotes  $W^{(0)}$  (resp.  $W_h^{(0)}$ ,  $h \in (0, h_*)$ ).

Next, for  $q \geq 0$ , recall the Laplace transforms of the functions  $W^{(q)}$  and  $Z^{(q)}$  [23, p. 214, Theorem 8.1] (for  $\beta > \Phi(q)$ ):  $\int_0^\infty e^{-\beta x} W^{(q)}(x) dx = 1/(\psi(\beta) - q)$  and  $\int_0^\infty e^{-\beta x} Z^{(q)}(x) dx = \frac{1}{\beta} \left(1 + \frac{q}{\psi(\beta) - q}\right)$  (where the latter formula follows using e.g. integration by parts). The Laplace transforms of  $W_h^{(q)}$  and  $Z_h^{(q)}$ ,  $h \in (0, h_*)$ , follow from Proposition 3.1 and Remark 3.2 (ii).

**Proposition 4.1** (Pointwise convergence). *Suppose  $\psi^h \rightarrow \psi$  and  $\Phi^h \rightarrow \Phi$  pointwise as  $h \downarrow 0$ . Then, for each  $q \geq 0$ ,  $W_h^{(q)} \rightarrow W^{(q)}$  and  $Z_h^{(q)} \rightarrow Z^{(q)}$  pointwise, as  $h \downarrow 0$ .*

*Remark 4.1.* We will see in  $(\Delta_1)$  of Subsection 5.3 that, in fact,  $\psi^h \rightarrow \psi$  locally uniformly in  $[0, \infty)$  as  $h \downarrow 0$ , which implies that  $\Phi^h \rightarrow \Phi$  pointwise as  $h \downarrow 0$ . In particular, given any  $q \geq 0$ ,  $\gamma > \Phi(q)$  implies  $\gamma > \Phi^h(q)$  for all  $h \in (0, h_0)$ , for some  $h_0 > 0$ .

*Proof.* Since  $\Phi^h(q) \rightarrow \Phi(q)$  as  $h \downarrow 0$ , it follows via integration by parts ( $\int_{[0, \infty)} e^{-\beta x} dF(x) = \beta \int_{(0, \infty)} e^{-\beta x} F(x) dx$  for any  $\beta \geq 0$  and any nondecreasing right-continuous  $F : \mathbb{R} \rightarrow \mathbb{R}$  vanishing on  $(-\infty, 0)$  [32, Chapter 0, (4.5) Proposition]) that, for some  $h_0 > 0$ , the Laplace transforms of  $dW^{(q)}$ ,  $dZ^{(q)}$ ,  $(dW_h^{(q)})_{h \in (0, h_0)}$  and  $(dZ_h^{(q)})_{h \in (0, h_0)}$ , are defined (i.e. finite). These measures are furthermore concentrated on  $[0, \infty)$  and since  $\psi^h \rightarrow \psi$  pointwise as  $h \downarrow 0$ , then  $\widehat{dW_h^{(q)}} \rightarrow \widehat{dW^{(q)}}$  and  $\widehat{dZ_h^{(q)}} \rightarrow \widehat{dZ^{(q)}}$  pointwise as  $h \downarrow 0$ . By [6, p. 110, Theorem 8.5], it follows that  $dW_{h_n}^{(q)} \rightarrow dW^{(q)}$  and  $dZ_{h_n}^{(q)} \rightarrow dZ^{(q)}$  vaguely as  $n \rightarrow \infty$ , for any sequence  $(h_n)_{n \geq 1} \downarrow 0$ . This implies that, as  $h \downarrow 0$ ,  $W_h^{(q)} \rightarrow W^{(q)}$  (respectively  $Z_h^{(q)} \rightarrow Z^{(q)}$ ) pointwise at all points of continuity of  $W^{(q)}$  (respectively  $Z^{(q)}$ ). Now, the functions  $Z^{(q)}$  are continuous everywhere, whereas  $W^{(q)}$  is continuous on  $\mathbb{R} \setminus \{0\}$  and has a jump at 0, if and only if  $X$  has sample paths of finite variation [23, p. 222, Lemma 8.6]. In the latter case, however, we necessarily have  $\sigma^2 = 0$  and  $\int_{[-1, 0)} |y| \lambda(dy) < \infty$  [33, p. 140, Theorem 21.9] and the jump size is  $W^{(q)}(0) = 1/\mu_0$  (see Section 2 for definition of  $\mu_0$ ). By Remark 3.2 (iii) and (2.3),  $W_h^{(q)}(0) = 1/(h\lambda^h(\{h\})) = 1/(\mu - \mu^h + c_0^h/h)$ . The latter quotient, however, converges to  $1/\mu_0$ , as  $h \downarrow 0$ , by the DCT.  $\square$

## 5. RATES OF CONVERGENCE

In this section we establish our main result, Theorem 1.1 from the Introduction. Subsection 5.1 describes the general method of proof and establishes some preliminary observations and notation.

Subsection 5.2 contains technical results, notationally and otherwise independent from rest of the text, which are applied time and again in the sequel. Then Subsections 5.3–5.9 establish a series of convergence results, which together imply Theorem 1.1. Finally, Subsection 5.10 contains a convergence result for the derivatives  $W^{(q)'$ .

**5.1. Method of proof, preliminary observations and notation.** The key step in the proof of Theorem 1.1 consists of a detailed analysis of the relevant differences arising in the integral representations (see Paragraph 5.1.1) of the scale functions, see Paragraph 5.1.2. A more detailed explanation of the method of proof will be given in Paragraph 5.1.3.

*Remark 5.1.* With reference to Subsection 2.2, there is of course extensive literature on numerical solutions to integro-differential equations (IDE) of the relevant (Volterra) type (viz. Eq. (2.6)). This literature will, however, typically assume at least the continuity of the kernel appearing in the integral of the IDE, to even pose the problem, and obtain rates of convergence under additional smoothness conditions thereon (and the solution to the IDE) [10, Chapters 2 and 3] [26, Chapters 7 and 11]. In our case the kernel appearing in (2.6) is of course not (necessarily) even continuous (let alone possessing higher degrees of smoothness). Further, discounting for a moment the continuity requirement on the kernel, which may appear technical, some relevant *general* results on convergence do exist, e.g. [26, p. 102, Theorem 7.2] for the case  $\sigma^2 = 0$  &  $\lambda(\mathbb{R}) < +\infty$ , but are not really (directly) applicable, since one would need to *a priori* establish (at least) a rate of convergence for the difference between the integral appearing in (2.6) and its discretization (local consistency error; see [26, p. 101, Eq. (7.12)]). This does not appear possible in general without a knowledge of the (sufficient) smoothness properties of the target function  $W^{(q)}$  (the latter not always being clear; see [12]) and indeed, it would seem, those of the tail ( $y \mapsto \lambda(-\infty, -y)$ ). Such an error analysis would be further complicated when  $\sigma^2 > 0$  (respectively  $\kappa(0) = \infty$ ), since then we are dealing with the discretization also of the derivative of  $W^{(q)}$  (respectively the integral in (2.6) cannot be split up as the difference of the integrals of each individual term of the integrand). It is then not very likely that looking at this problem from the integro-differential perspective alone would allow us to obtain, moreover sharp, rates of convergence (at least not in general).

By contrast, the method for obtaining the sharp rates of convergence that we shall use, based on the integral representations of the scale functions and their approximations, will allow us to handle all the cases within a single framework.

5.1.1. *Integral representations of scale functions.*

**Proposition 5.1.** *Let  $q \geq 0$ . For all  $\beta \in \mathbb{C}$  with  $\Re\beta > \Phi(q)$ ,  $\psi(\beta) - q \neq 0$  (respectively  $\psi^h(\beta) - q \neq 0$ ) and one has  $(\psi(\beta) - q)\widehat{W}^{(q)}(\beta) = 1$  (respectively  $\beta h(\psi^h(\beta) - q)\widehat{W}_h^{(q)}(\beta) = e^{\beta h} - 1$ ), and  $\beta\widehat{Z}^{(q)}(\beta) = 1 + \frac{q}{\psi(\beta) - q}$  (respectively  $\beta\widehat{Z}_h^{(q)}(\beta) = 1 + \frac{q}{\psi^h(\beta) - q}$ ) for the scale functions of  $X$  (respectively  $X^h$ ,  $h \in (0, h_*)$ ).*

*Proof.* The stipulated equalities extend from  $\beta > \Phi(q)$  real, to complex  $\beta$  with  $\Re\beta > \Phi(q)$ , via analytic continuation, using expressions for the Laplace transforms of the scale functions (the latter having been noted in Section 4). In particular, so extended, they then imply  $\psi(\beta) - q \neq 0$  for the range of  $\beta$  as given.  $\square$

**Corollary 5.1** (Integral representation of scale functions). *Let  $q \geq 0$ . For any  $\gamma > \Phi(q)$ , we have, for all  $x > 0$  (with  $\beta := \gamma + is$ ):*

$$W^{(q)}(x) = \frac{1}{2\pi} \lim_{T \rightarrow \infty} \int_{-T}^T \frac{e^{\beta x}}{\psi(\beta) - q} ds \quad (5.1)$$

and

$$Z^{(q)}(x) = \frac{1}{2\pi} \lim_{T \rightarrow \infty} \int_{-T}^T \frac{e^{\beta x}}{\beta} \left( 1 + \frac{q}{\psi(\beta) - q} \right) ds. \quad (5.2)$$

Likewise, for any  $h \in (0, h_*)$  and then any  $\gamma > \Phi^h(q)$ , we have, for all  $x \in \mathbb{Z}_h^+$  (again with  $\beta := \gamma + is$ ):

$$W_h^{(q)}(x) = \frac{1}{2\pi} \int_{-\pi/h}^{\pi/h} \frac{e^{\beta(x+h)}}{\psi^h(\beta) - q} ds \quad (5.3)$$

and

$$Z_h^{(q)}(x) = \frac{1}{2\pi} \int_{-\pi/h}^{\pi/h} \frac{e^{\beta x}}{\beta} \frac{\beta h}{1 - e^{-\beta h}} \left( 1 + \frac{q}{\psi^h(\beta) - q} \right) ds. \quad (5.4)$$

*Proof.* First note that  $W^{(q)}$  and  $Z^{(q)}$  (respectively  $W_h^{(q)}$  and  $Z_h^{(q)}$ ) are of  $\gamma$ -exponential order for all  $\gamma > \Phi(q)$  (respectively  $\gamma > \Phi^h(q)$ ,  $h \in (0, h_*)$ ). Then use the inverse Laplace [14, Section 3.3] (respectively  $Z$  [19, p. 11]) transform.  $\square$

5.1.2. *The differences  $\Delta_W^{(q)}$  and  $\Delta_Z^{(q)}$ .* For  $x \geq 0$ , let  $T_x$  (resp.  $T_x^h$ ) denote the first entrance time of  $X$  (resp.  $X^h$ ) to  $[x, \infty)$ , let  $\underline{X}_t := \inf\{X_s : s \in [0, t]\}$  (resp.  $\underline{X}_t^h := \inf\{X_s^h : s \in [0, t]\}$ ),  $t \geq 0$ , be the running infimum process and  $\underline{X}_\infty := \inf\{X_s : s \in [0, \infty)\}$  (resp.  $\underline{X}_\infty^h := \inf\{X_s^h : s \in [0, \infty)\}$ ) the overall infimum of  $X$  (resp.  $X^h$ ,  $h \in (0, h_*)$ ).

In the case of the spectrally negative process  $X$ , it follows from [23, Theorem 8.1 (iii)], regularity of 0 for  $(0, \infty)$  [23, p. 212], dominated convergence and continuity of  $W^{(q)}|_{(0, \infty)}$  that, for  $q \geq 0$  and  $\{x, y\} \subset \mathbb{R}^+$ :

$$\mathbb{E}[e^{-qT_y} \mathbb{1}(\underline{X}_{T_y} \geq -x)] = \frac{W^{(q)}(x)}{W^{(q)}(x+y)} \quad \text{and} \quad \mathbb{E}[e^{-qT_y} \mathbb{1}(\underline{X}_{T_y} > -x)] = \frac{W^{(q)}(x)}{W^{(q)}(x+y)}. \quad (5.5)$$

On the other hand, we find [35, Theorem 4.6] the direct analogues of these two formulae in the case of the approximating processes  $X^h$ ,  $h \in (0, h_*)$  as being ( $q \geq 0$ ,  $\{x, y\} \subset \mathbb{Z}_h^{++}$ ):

$$\mathbb{E}[e^{-qT_y^h} \mathbb{1}(\underline{X}_{T_y^h}^h \geq -x)] = \frac{W_h^{(q)}(x)}{W_h^{(q)}(x+y)} \quad \text{and} \quad \mathbb{E}[e^{-qT_y^h} \mathbb{1}(\underline{X}_{T_y^h}^h > -x)] = \frac{W^{(q)}(x-h)}{W^{(q)}(x-h+y)}. \quad (5.6)$$

We conclude by comparing (5.5) with (5.6) that there is *no a priori probabilistic reason* to favour either  $W_h^{(q)}$  or  $W_h^{(q)}(\cdot - h)$  in the choice of which of these two quantities to compare to  $W^{(q)}$ . Nevertheless, this choice is not completely arbitrary:

(a) In view of (5.1) and (5.3), the quantity  $W_h^{(q)}(\cdot - h)$  seems more favourable (cf. also the findings of Proposition 5.5, especially when  $q = |\mu| = 0$ ). In addition, when  $X$  has sample paths of infinite variation, a.s.,  $W^{(q)}(0)$  is equal to zero [22, p. 33, Lemma 3.1] and so is  $W_h^{(q)}(-h)$ , whereas  $W_h^{(q)}(0)$  is always strictly positive ( $h \in (0, h_*)$ ).

(b) On the other hand, when  $X$  has sample paths of finite variation, a.s., then  $W^{(q)}(0) = 1/\mu_0 > 0$  [22, p. 33, Lemma 3.1] and if in addition the Lévy measure is finite, then in fact also  $W_h^{(q)}(0) = 1/\mu_0$  for all  $h \in (0, h_*)$ .

*Remark 5.2.* (i) It follows from the above discussion that it is reasonable to approximate  $W^{(q)}$  by  $W_h^{(q)}(\cdot - h)$  (resp.  $W_h^{(q)}$ ), when  $X$  has sample paths of infinite (resp. finite) variation (a.s.).

(ii) When  $q = 0$  or  $x = 0$ ,  $Z^{(q)}(x) = Z_h^{(q)}(x) = 1$ ,  $h \in (0, h_*)$ . Thus, when comparing these functions, we shall always assume  $q \wedge x > 0$ , for the only interesting case.

In view of Remark 5.2 (i) we define  $\delta^0$  to be equal to 0 or 1 according as the sample paths of  $X$  are of finite or infinite variation (a.s.). Fix  $q \geq 0$ . For  $h \in (0, h_*)$  we then define the differences:

$$\Delta_W^{(q)}(x, h) := W^{(q)}(x) - W_h^{(q)}(x - \delta^0 h), \quad x \in \mathbb{Z}_h^{++} \cup \{\delta^0 h\} \quad (5.7)$$

and

$$\Delta_Z^{(q)}(x, h) := Z^{(q)}(x) - Z_h^{(q)}(x), \quad x \in \mathbb{Z}_h^{++}. \quad (5.8)$$

Fix further any  $\gamma > \Phi(q)$ . Let  $h \in (0, h_*)$  be such that also  $\gamma > \Phi^h(q)$ , Then Corollary 5.1 implies, for any  $x \in \mathbb{Z}_h^{++} \cup \{\delta^0 h\}$  (we always let, here and in the sequel,  $\beta := \gamma + is$  to shorten notation):

$$\begin{aligned} e^{-\gamma x} 2\pi \Delta_W^{(q)}(x, h) &= \underbrace{\lim_{T \rightarrow \infty} \int_{(-T, T) \setminus (-\pi/h, \pi/h)} e^{isx} \frac{ds}{\psi(\beta) - q}}_{(a)} + \underbrace{\int_{[-\pi/h, \pi/h]} e^{isx} \left[ \frac{\psi^h - \psi}{(\psi - q)(\psi^h - q)} \right] (\beta) ds}_{(b)} + \\ &\underbrace{(1 - \delta^0) \int_{[-\pi/h, \pi/h]} e^{isx} (1 - e^{ish}) \frac{ds}{\psi^h(\beta) - q}}_{(c)} \end{aligned} \quad (5.9)$$

whereas for  $x \in \mathbb{Z}_h^{++}$ :

$$\begin{aligned} e^{-\gamma x} 2\pi \Delta_Z^{(q)}(x, h) &= \underbrace{\lim_{T \rightarrow \infty} \int_{(-T, T) \setminus (-\pi/h, \pi/h)} \frac{e^{isx}}{\beta} \left( \frac{q}{\psi(\beta) - q} \right) ds}_{(a)} \\ &+ \underbrace{\int_{[-\pi/h, \pi/h]} \frac{e^{isx}}{\beta} \left( 1 - \frac{\beta h}{1 - e^{-\beta h}} \right) \left( \frac{q}{\psi^h(\beta) - q} \right) ds}_{(b)} \\ &+ \underbrace{q \int_{[-\pi/h, \pi/h]} \frac{e^{isx}}{\beta} \left[ \frac{\psi^h - \psi}{(\psi^h - q)(\psi - q)} \right] (\beta) ds}_{(c)}. \end{aligned} \quad (5.10)$$

Note that in (5.10) we have taken into account that the difference between the inverse Laplace and inverse  $Z$  transform, for the function, which is identically equal to 1, vanishes identically.

5.1.3. *Method for obtaining the rates of convergence in (5.9) and (5.10).* Apart from the Brownian motion with drift case, which is treated explicitly, the method for obtaining the rates of convergence for the differences (5.9) and (5.10) is as follows (recall  $\beta = \gamma + is$ ):

- (1) First we estimate  $|\psi^h - \psi|(\beta)$  to control the numerators. In particular, we are able to conclude  $\psi^h \rightarrow \psi$ , uniformly in bounded subsets of  $\overline{\mathbb{C}^-}$ . See Subsection 5.3.
- (2) Then we show  $|\psi^h - q|(\beta)$  is suitably bounded from below on  $s \in (-\pi/h, \pi/h)$ , uniformly in  $h \in [0, h_0)$ , for some  $h_0 > 0$ . This property, referred to as *coercivity*, controls the denominators. See Subsection 5.5.
- (3) Finally, using (1) and (2), one can estimate the integrals appearing in (5.9) and (5.10) either by a direct  $|\int \cdot ds| \leq \int |\cdot| ds$  argument, or else by first applying a combination of integrations by parts (see (5.11) below) and Fubini's Theorem. In the latter case, the estimates of  $|\frac{d(\psi(\beta) - \psi^h(\beta))}{ds}|$  and the growth in  $s$ , as  $|s| \rightarrow \infty$ , of  $\frac{d(\psi^h - q)(\beta)}{ds}$ ,  $h \in [0, h_*)$ , also become relevant, and we provide these in Subsection 5.4.

*Remark 5.3.* (i) Note that the integral representation of the scale functions is crucial for our programme to yield results. The formulae (5.9) and (5.10) suffice to give a precise rate locally uniformly in  $x \in (0, \infty)$ .

(ii) The integration by parts in (3) is applied according to the general scheme ( $f$  differentiable,  $x > 0$ ):

$$\frac{1}{x} \frac{d}{ds} (e^{isx} f(s)) = ie^{isx} f(s) + \frac{1}{x} e^{isx} f'(s), \quad (5.11)$$

and with the integral  $\int e^{isx} f(s) ds$  (over a relevant domain) in mind. Then, upon integration against  $ds$ , the left-hand side and the second term on the right-hand side of (5.11) admit for an estimate, which could not be made for  $\int e^{isx} f(s) ds$  directly, but in turn a factor of  $1/x$  emerges, implying (as we will see) that the final bound is locally uniform in  $(0, \infty)$  (in the estimates there is always also present a factor of  $e^{\gamma x}$  which from the perspective of the relative error, and in view of the growth properties of  $W^{(q)}$  and  $Z^{(q)}$  at  $+\infty$  [22, p. 129, Lemma 3.3], is perhaps not so bad). In fact, the convergence rate obtained via (1)-(3) is uniform in bounded subsets of  $(0, \infty)$ , if (3) does not involve integration by parts.

(iii) Now, it is usually the case that the estimates from (3) may be made by a direct application of the integral triangle inequality. We were not able to avoid integration by parts, however, in the case of the convergence for the functions  $W^{(q)}$ ,  $q \geq 0$ , and even then only if  $\sigma^2 = 0$  (see Subsections 5.8 and 5.9). Particularly delicate is the case when furthermore the sample paths of  $X$  are of finite variation (a.s.). In the latter case a key lemma is Lemma 5.6, which itself depends crucially on the findings of Proposition 5.2.

(iv) Even when  $\sigma^2 = 0$ , however, numerical experiments (see Section 6 and Appendix C) seem to suggest that, at least in some further subcases, one should be able to establish convergence for the functions  $W^{(q)}$ ,  $q \geq 0$ , which is uniform in bounded (rather than just compact) subsets of  $(0, \infty)$ . This remains open for future research.

*Remark 5.4.* Sharpness of the rates is obtained by constructing specific examples of Lévy processes, for which convergence is no better than stipulated (cf. the statement of Theorem 1.1). The key observation here is the following principle of *reduction by domination*:

Suppose we seek to prove that  $f \geq 0$  converges to 0 no faster than  $g > 0$ , i.e. that  $\limsup_{h \downarrow 0} f(h)/g(h) \geq C > 0$  for some  $C$ . If one can show  $f(h) \geq A(h) - B(h)$  and  $B = o(g)$ , then to show  $\limsup_{h \downarrow 0} f(h)/g(h) \geq C$ , it is sufficient to establish  $\limsup_{h \downarrow 0} A(h)/g(h) \geq C$ .

(This principle was also applied in [30] to establish sharpness of the stated rates of convergence there.)

We will use the basic, but very useful, principle of reduction by domination without explicit reference in the sequel.

5.1.4. *Further notation.* Notation-wise, we let (where  $\delta \in [0, 1]$ ):

$$\xi(\delta) := \int_{[-\delta, 0)} u^2 \lambda(du), \quad \kappa(\delta) := \int_{[-1, -\delta)} |y| \lambda(dy), \quad \zeta(\delta) := \delta \kappa(\delta) \quad \text{and} \quad \gamma(\delta) := \delta^2 \lambda([-1, -\delta))$$

and remark that, by the findings of [30, Lemma 3.8],  $\gamma(\delta) + \zeta(\delta) + \xi(\delta) \rightarrow 0$  as  $\delta \downarrow 0$ .

Finally, note that, unless otherwise indicated, we consider henceforth as having fixed:

$$X, \quad (X^h)_{h \in (0, h_*)}, \quad q \geq 0 \quad \text{and a} \quad \gamma > \Phi(q).$$

We insist that the dependence on  $x$  of the error estimates will be kept explicit throughout, whereas the dependence on the Lévy triplet,  $q$  and  $\gamma$  will be subsumed in the capital (or small)  $O$  ( $o$ ) notation. In particular, the notation  $f(x, h) = g(x, h) + l(x)O(h)$ ,  $x \in A$ , means that  $l(x) > 0$  for  $x \in A$  and:

$$\sup_{x \in A} |(f(x, h) - g(x, h))/l(x)| = O(h)$$

and analogously when  $O(h)$  is replaced by  $o(h)$  etc. Further, we shall sometimes resort to the notation:

$$A(s) := A^0(s) := \psi(\gamma + is) - q \quad (\text{for } s \in \mathbb{R}) \quad \text{and} \quad A^h(s) := \psi^h(\gamma + is) - q \quad (\text{for } s \in [-\pi/h, \pi/h]),$$

$h \in (0, h_*)$ , where reference to  $q$  and  $\gamma$  has been suppressed. We stress that in Subsections 5.3-5.10 we reserve the symbol  $\beta$  to mean  $\gamma + is$ , i.e., throughout we shall have:

$$\beta := \gamma + is.$$

The remainder of our analysis in this section will proceed as follows. First we list in Subsection 5.2, for the readers convenience, a number of auxiliary technical results. Their proofs, which are independent of the analysis in Section 5, are relegated to Appendices A and B. Then Subsection 5.3 estimates the absolute difference  $|\psi^h - \psi|$ , Subsection 5.4 analyzes the derivatives  $A^{h'}$  and the difference  $|A^{h'} - A'|$ , while in Subsection 5.5 we prove suitable coercivity of  $|\psi^h - q|$ . Subsections 5.6-5.9



deal with the various cases of convergence for the scale functions. Subsection 5.10 establishes a convergence result for the derivatives of  $W^{(q)}$  in the case when  $\sigma^2 > 0$ .

**5.2. Auxiliary technical results.** [Apart from the notation of Subsection 2.3, the contents of this subsection is notationally and otherwise independent from the remainder of the text. For proofs see Appendices A and B.]

5.2.1. *Some estimates and bounds.*

**Lemma 5.1.** *For every  $\gamma_\star \in \mathbb{R}_+$  and  $h_\star \in \mathbb{R}^+$ , there is an  $A_0 \in (0, \infty)$  such that for all  $\gamma \in [-\gamma_\star, \gamma_\star]$ ,  $h \in (0, h_\star)$  and then all  $s \in [-\pi/h, \pi/h]$  (with  $\beta = \gamma + is$ ):*

- (i)  $\left| \frac{1}{2h}(e^{\beta h} - e^{-\beta h}) \right| \leq A_0 |\beta|$ .
- (ii)  $|e^{\beta h} - 1| \leq A_0 |\beta| h$ .
- (iii)  $\left| \frac{1}{h}(e^{\beta h} - 1) - \beta \right| \leq A_0 h |\beta|^2$ .
- (iv)  $\left| \frac{1}{2h^2}(e^{\beta h} + e^{-\beta h} - 2) \right| \leq A_0 |\beta|^2$ .
- (v)  $\left| \frac{1}{2h^2}(e^{\beta h} + e^{-\beta h} - 2 - (\beta h)^2) \right| \leq A_0 h^2 |\beta|^4$ .
- (vi)  $\left| \frac{1}{2h}(e^{\beta h} - e^{-\beta h} - 2\beta h) \right| \leq A_0 h^2 |\beta|^3$ .

*Further:*

- (a) For any  $\xi \in [0, 2\pi)$ ,  $(u \mapsto (1 - \cos(u))/u^2)$  is bounded away from 0 for  $u \in [-\xi, \xi]$ .
- (b) For any  $\xi \in [0, 2\pi)$ ,  $(u \mapsto \frac{1}{u^2}(\cosh(u) - 1))$  is bounded away from 0 for  $\Im u \in [-\xi, \xi]$ .
- (c) For any  $\xi \in [0, 2\pi)$  and  $L \in \mathbb{R}$ ,  $(u \mapsto \frac{1}{u}(e^u - 1))$  is bounded away from 0 for  $\Im u \in [-\xi, \xi]$  and  $\Re u \geq L$ .

*Remark 5.5.* In (a)-(c), at  $u = 0$ , the relevant limit (which exists) is taken, in order to make the mappings well-defined in this single point.

**Lemma 5.2.** *The family of functions  $f_{\gamma_0} : [-\pi, \pi] \rightarrow \mathbb{R}$ , defined by:  $f_{\gamma_0}(s_0) := \left(1 - \frac{\gamma_0 + is_0}{1 - e^{-\gamma_0 - is_0}}\right) \frac{1}{\gamma_0 + is_0}$ ,  $s_0 \in [-\pi, \pi]$ , is uniformly bounded for  $\gamma_0$  belonging to bounded subsets of  $\mathbb{R} \setminus \{0\}$ .*

**Lemma 5.3.** (i) For any  $z \in \overline{\mathbb{C}^{\leftarrow}}$ :  $|e^z - 1| \leq |z|$ .

(ii) There exists  $C \in (0, \sqrt{5/2}]$ , such that for any  $z \in \overline{\mathbb{C}^{\leftarrow}}$ :  $|e^z - z - 1| \leq C|z|^2$ .

**Lemma 5.4.** (i) Let  $\{x, y\} \subset \mathbb{R}_-$ ,  $|x - y| \leq h/2 \leq |y|$ . Then for any  $\alpha \in \overline{\mathbb{C}^{\rightarrow}}$  we have:

$$|e^{\alpha x} - \alpha x - (e^{\alpha y} - \alpha y)| \leq 2h|\alpha|^2|y|.$$

(ii) There exists  $C \in (0, \infty)$  such that whenever  $\{x, y\} \subset \mathbb{R}_-$ ,  $\alpha \in \overline{\mathbb{C}^{\rightarrow}}$  and  $|x - y| \leq h/2 \leq |y|$ ,

$$\text{we have: } \left| e^{\alpha x} - \alpha x - \frac{\alpha^2 x^2}{2} - \left( e^{\alpha y} - \alpha y - \frac{\alpha^2 y^2}{2} \right) \right| \leq Chy^2|\alpha|^3.$$

5.2.2. *Some asymptotic properties at 0 of measures on  $\mathbb{R}$ .* Let  $\nu$  be a measure on  $\mathbb{R}$ .

**Proposition 5.2.** *If  $\nu$  is compactly supported and locally finite in  $\mathbb{R} \setminus \{0\}$ , then:  $\int |x|\nu(dx) < \infty$ , precisely when  $\int_1^\infty \frac{ds}{s^2} \int \nu(dx)(1 - \cos(sx)) < \infty$ .*

**Lemma 5.5.** *Let  $r \geq 0$ . Then:*

- (i)  $\int_{[0,r]} x\nu(dx) = \int_{[0,r]} \nu((t, r])dt.$
- (ii)  $\int_{[-r,r]} |x|\nu(dx) = \int_{[0,r]} \nu([-r, r] \setminus [-t, t])dt.$
- (iii)  $\int_{[-1,1] \setminus [-r,r]} |x|\nu(dx) = rg(r) + \int_{(r,1]} g(t)dt$  whenever  $r \leq 1$  and with  $g(r) := \nu([-1, 1] \setminus [-r, r]).$
- (iv)  $\int_{[0,r]} x^2\nu(dx) = 2 \int_{[0,r]} t\nu((t, r])dt.$
- (v)  $\int_{[-r,r]} x^2\nu(dx) = 2 \int_{[0,r]} t\nu([-r, r] \setminus [-t, t])dt.$

**Proposition 5.3.** *Suppose  $g(\delta) := \nu([-1, 1] \setminus [-\delta, \delta]) \sim 1/\delta^{1+\alpha}$  as  $\delta \downarrow 0$ , so that in particular  $g(\delta)$  is finite for all  $0 < \delta \leq 1$  and necessarily  $\alpha \geq -1$ . Then:*

- (a)  $\gamma(\delta) := \delta^2\nu([-1, 1] \setminus [-\delta, \delta]) \sim \delta^{1-\alpha}$  as  $\delta \downarrow 0$ .
- (b)  $\int_{[-1,1]} x^2\nu(dx) < \infty$ , iff  $\alpha < 1$ . If  $\alpha \in (-1, 1)$ , then  $\int_{[-\delta,\delta]} x^2\nu(dx) \sim \delta^{1-\alpha}$  as  $\delta \downarrow 0$ .
- (c)  $\nu$  is a Lévy measure, iff  $\nu(\mathbb{R} \setminus [-1, 1]) < \infty$ ,  $\nu(\{0\}) = 0$  and  $\alpha < 1$ .
- (d)  $\int_{[-1,1]} |x|\nu(dx) = \infty$ , iff  $0 \leq \alpha$ .
- (e) Finally, as  $\delta \downarrow 0$ , if  $\alpha > 0$ ,  $\int_{(\delta,1]} g(t)dt \sim \delta^{-\alpha}$  and if  $\alpha = 0$ , then  $\int_{(\delta,1]} g(t)dt \sim |\log \delta|$ . In particular,  $\zeta(\delta) := \delta \int_{[-1,1] \setminus [-\delta,\delta]} |x|\nu(dx) \sim \delta^{1-\alpha}$ , when  $\alpha > 0$ , respectively  $\zeta(\delta) \sim \delta |\log \delta|$ , when  $\alpha = 0$ .

**Proposition 5.4.** *Define  $\lambda(dx) = \mathbb{1}_{(-1,1)}(x)|x|\nu(dx)$ . Furthermore, let  $\alpha \in (0, 1)$ , with  $\limsup_{\delta \downarrow 0} \nu((-1, 1) \setminus [-\delta, \delta])\delta^{1+\alpha} < \infty$ . Then each of the quantities  $\limsup_{\delta \downarrow 0} \lambda((-1, 1) \setminus [-\delta, \delta])\delta^\alpha$ ,  $\limsup_{\delta \downarrow 0} \int_{[-\delta,\delta]} x^2\nu(dx)\delta^{\alpha-1}$  and  $\sup_{s \in \mathbb{R} \setminus \{0\}} \frac{1}{|s|^\alpha} |\int(e^{isy} - 1)\lambda(dy)|$  is finite.*

**5.3. Estimating the absolute difference  $|\psi^h - \psi|$ .** Recall that  $\beta$  is defined to be  $\gamma + is$  throughout. We establish in this subsection two key properties of the difference  $\psi^h - \psi$ :

- ( $\Delta_1$ )  $\psi^h \rightarrow \psi$  as  $h \downarrow 0$ , uniformly in bounded subsets of  $\overline{\mathbb{C}^-}$ .
- ( $\Delta_2$ ) There exists  $A_0 \in (0, \infty)$ , such that for all  $h \in (0, h_* \wedge 2)$  and then all  $s \in [-\pi/h, \pi/h]$ , the following holds (see Table 1 for values of the parameter  $V$ ):

(i) When  $\sigma^2 > 0$ :

$$|\psi^h - \psi|(\beta) \leq A_0 [h^2|\beta|^4 + h\xi(h/2)|\beta|^3 + h|\beta| + V\zeta(h/2)|\beta|^2].$$

In particular, if in addition  $\kappa(0) < \infty$ , we have:

$$|\psi^h - \psi|(\beta) \leq A_0 [h^2|\beta|^4 + h|\beta|^2].$$

If, moreover,  $\lambda(\mathbb{R}) < \infty$ , then

$$|\psi^h - \psi|(\beta) \leq A_0 [h^2|\beta|^4 + h|\beta|].$$

(ii) When  $\sigma^2 = 0$ :

$$|\psi^h - \psi|(\beta) \leq A_0 [h\xi(h/2)|\beta|^3 + (h + \zeta(h/2))|\beta|^2].$$

If in addition  $\kappa(0) < \infty$ , then:

$$|\psi^h - \psi|(\beta) \leq A_0 h |\beta|^2.$$

*Proof of  $(\Delta_1)$  and  $(\Delta_2)$ .* Indeed, suppose  $\sigma^2 > 0$  (respectively  $\sigma^2 = 0$ ), so that we are working under scheme 1 (respectively scheme 2). We decompose, referring to (2.1), (2.2) and (2.3), the difference  $\psi^h - \psi$  into terms, which allow for straightforward estimates. To wit, for any  $h_0 \in (0, 2]$  and  $\rho_0 > 0$ , there exists  $A_0 \in (0, \infty)$ , such that for all  $h \in (0, h_0)$  and then all  $s \in [-\pi/h, \pi/h]$ , as well as all  $\rho \in [0, \rho_0]$  (with  $\alpha = \rho + is$ ):

- (1)  $\left| \sigma^2 \left( \frac{e^{\alpha h} + e^{-\alpha h} - 2}{2h^2} - \frac{\alpha^2}{2} \right) \right| \leq A_0 h^2 |\alpha|^4$  by (v) of Lemma 5.1 (respectively this is void).
- (2)  $\left| c_0^h \left( \frac{e^{\alpha h} + e^{-\alpha h} - 2}{2h^2} - \frac{\alpha^2}{2} \right) \right| \leq A_0 h^2 \xi(h/2) |\alpha|^4$  by (v) of Lemma 5.1.
- (3) By a direct Taylor expansion:

$$\begin{aligned} & \left| V \int_{[-h/2, 0)} y^2 \frac{\alpha^2}{2} \lambda(dy) - \int_{[-h/2, 0)} (e^{\alpha y} - V\alpha y - 1) \lambda(dy) \right| \\ & \leq A_0 (V|\alpha|^3 h \xi(h/2) + (1 - V)|\alpha| h). \end{aligned}$$

- (4)  $\left| \mu \left( \frac{e^{\alpha h} - e^{-\alpha h}}{2h} - \alpha \right) \right| \leq A_0 h^2 |\alpha|^3$  by (vi) (respectively  $\left| \mu \left( \frac{e^{\alpha h} - 1}{h} - \alpha \right) \right| \leq A_0 h |\alpha|^2$  by (iii) of Lemma 5.1).
- (5)  $\left| (-\mu^h) \left( \frac{e^{\alpha h} - e^{-\alpha h}}{2h} - \alpha \right) \right| \leq A_0 V h^2 \kappa(h/2) |\alpha|^3$  (respectively  $\left| (-\mu^h) \left( \frac{e^{\alpha h} - 1}{h} - \alpha \right) \right| \leq A_0 V h \kappa(h/2) |\alpha|^2$ ), by the same token, since in fact:

$$-\mu^h = - \sum_{y \in \mathbb{Z}_h^-} y \int_{A_y^h} \mathbb{1}_{[-V, 0)}(z) \lambda(dz) \leq -2V \int_{[-1, -h/2)} z \lambda(dz) = 2V \kappa(h/2).$$

- (6) Finally:

$$\begin{aligned} & \left| \sum_{y \in \mathbb{Z}_h^-} c_y^h (e^{\alpha y} - 1) - \alpha \mu^h - \int_{(-\infty, -h/2]} (e^{\alpha z} - \alpha z \mathbb{1}_{[-V, 0)}(z) - 1) \lambda(dz) \right| \\ & \leq \left| \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h \cap (-\infty, -V)} (e^{\alpha y} - e^{\alpha z}) \lambda(dz) \right| + \left| \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h \cap [-V, 0)} (e^{\alpha y} - e^{\alpha z} - V\alpha(y - z)) \lambda(dz) \right| \\ & \leq A_0 (|\alpha| h + V |\alpha|^2 h \kappa(h/2)), \end{aligned}$$

by (ii) of Lemma 5.1 (since  $|e^{\alpha y} - e^{\alpha z}| \leq |1 - e^{\alpha(y-z)}|$ ) and (i) of Lemma 5.4.

From the estimates (1)-(6),  $(\Delta_1)$  follows, since any compact subset of  $\overline{\mathbb{C}^-}$  is contained in the rectangle  $[0, \rho_0] \times [-\pi/h, \pi/h]$ , for all  $h \in (0, h_0)$ , so long as  $\rho_0$  is chosen large enough, and  $h_0$  small enough. On the other hand  $(\Delta_2)$  follows by taking  $h_0 = h_* \wedge 2$  and  $\rho_0 = \rho = \gamma$ , so that  $\alpha = \beta$ .  $\square$

*Remark 5.6.* Pursuant to  $(\Delta_1)$  above and Remark 4.1, we assume henceforth that  $h_*$  has already been chosen small enough, so that in addition  $\gamma > \Phi^h(q)$  for all  $h \in (0, h_*)$

**5.4. Estimating the absolute difference  $|A^{h'} - A'|$  and growth of  $A^{h'}$  at infinity.** We establish here the following two properties pertaining to the derivatives  $A'$  and  $A^{h'}$ ,  $h \in (0, h_*)$ :

( $\Delta'_1$ ) For any finite  $h_0 \in (0, h_*]$ , there exists an  $A_0 \in (0, \infty)$  such that for all  $h \in [0, h_0)$  and then all  $s \in (-\pi/h, \pi/h)$ :

$$|A^{h'}(s)| \leq A_0 |s|^{\epsilon-1},$$

where  $\epsilon = 2$ , if  $\sigma^2 > 0$ ;  $\epsilon = 1$ , if  $\sigma^2 = 0$  and  $\kappa(0) < \infty$ ; finally, if  $\sigma^2 = 0$  and  $\kappa(0) = \infty$ , then  $\epsilon$  must satisfy (1) of Assumption 1.1 from the Introduction.

( $\Delta'_2$ ) There is an  $A_0 \in (0, \infty)$ , such that for all  $h \in (0, h_* \wedge 2)$  and then all  $s \in [-\pi/h, \pi/h]$ , the following holds:

(i) When  $\sigma^2 > 0$ :

$$|A^{h'}(s) - A'(s)| \leq A_0 (h^2 |\beta|^3 + h \xi(h/2) |\beta|^2 + (h + \zeta(h/2)) |\beta|).$$

(ii) When  $\sigma^2 = 0$ :

$$|A'(s) - A^{h'}(s)| \leq A_0 [h + \zeta(h/2) + \xi(h/2)] |\beta|.$$

If in addition  $\kappa(0) < \infty$ , then:

$$|\psi^h - \psi|(\beta) \leq A_0 h |\beta|.$$

*Proof of ( $\Delta'_1$ ) and ( $\Delta'_2$ ).* Indeed, we have, using differentiation under the integral sign, for  $s \in \mathbb{R}$ :

$$A'(s) = i\sigma^2 \beta + i\mu + i \int_{(-\infty, 0)} z \left( e^{\beta z} - \mathbb{1}_{[-V, 0)}(z) \right) \lambda(dz). \quad (5.12)$$

Suppose now first that  $\sigma^2 > 0$ . Then, for  $h \in (0, h_*)$  and  $s \in [-\pi/h, \pi/h]$ :

$$A^{h'}(s) = i\mu \frac{e^{\beta h} + e^{-\beta h}}{2} - i\mu^h \left( \frac{e^{\beta h} + e^{-\beta h}}{2} - 1 \right) + (\sigma^2 + c_0^h) i \frac{e^{\beta h} - e^{-\beta h}}{2h} + i \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h} y \left( e^{\beta y} - \mathbb{1}_{[-V, 0)}(z) \right) \lambda(dz).$$

From these expressions it follows readily, using (i) of Lemma 5.3, (i) and (iv) of Lemma 5.1,  $h|\mu^h| \leq 2h\kappa(h/2) \leq 4 \int_{[-1, 0)} y^2 \lambda(dy)$  and  $|y| \leq 2|z|$ ,  $e^{\gamma y} \leq e^{\gamma(z+h/2)}$  for  $z \in A_y^h$ ,  $y \in \mathbb{Z}_h^-$ , that  $A'$  and  $A^{h'}$  are both bounded by an affine function of  $|s|$  on  $s \in (-\pi/h, \pi/h)$ , uniformly in  $h \in (0, h_0)$  for any finite  $h_0 \in (0, h_*]$ .

On the other hand, when  $\sigma^2 = 0$ , we have for  $h \in (0, h_*)$  and then  $s \in [-\pi/h, \pi/h]$ :

$$\begin{aligned} A^{h'}(s) &= \frac{ic_0^h}{2h} \left( e^{\beta h} - e^{-\beta h} \right) + i(\mu - \mu^h) e^{\beta h} + i \sum_{y \in \mathbb{Z}_h^-} c_y^h y e^{\beta y} \\ &= \frac{ic_0^h}{2h} \left( e^{\beta h} - e^{-\beta h} \right) + i\mu e^{\beta h} - \mu^h (e^{\beta h} - 1) + i \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h} y \left( e^{\beta y} - \mathbb{1}_{[-V, 0)}(z) \right) \lambda(dz). \end{aligned} \quad (5.13)$$

Now, if  $\kappa(0) < \infty$ , it follows readily from  $c_0^h \leq h\kappa(0)$ ,  $-\mu^h \leq 2\kappa(0)$  and  $\int_{(-\infty, 0)} |y| e^{\gamma y} \lambda(dy) < \infty$  that  $A'$  and  $A^{h'}$  are bounded, uniformly in  $h \in (0, h_0)$  for any finite  $h_0 \in (0, h_*]$ . If, however,  $\kappa(0) = \infty$  and then under Assumption 1.1, the desired conclusion of ( $\Delta'_1$ ) follows from the estimates of Lemma 5.1 and Proposition 5.4 using (1) of Assumption 1.1.

Finally, from the above expressions for the derivatives  $A'$  and  $A^{h'}$ ,  $(\Delta'_2)$  follows using Lemma 5.1 and a decomposition similar to the one in Subsection 5.3, which allowed to establish  $(\Delta_2)$ . For example, when  $\sigma^2 = 0$ , we have the following decomposition of  $A^{h'}(s) - A'(s)$  into three summands, each of which is then easily estimated ( $h \in (0, h_\star \wedge 2)$ ):

- (1)  $\sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h} [y(e^{\beta y} - \mathbb{1}_{[-1,0)}(z)) - z(e^{\beta z} - \mathbb{1}_{[-1,0)}(z))] \lambda(dz)$ ;
- (2)  $i(\mu - \mu^h)(e^{\beta h} - 1)$ ;
- (3)  $i c_0^h \left[ \frac{e^{\beta h} - e^{-\beta h}}{2h} \right] - i \int_{[-h/2, 0)} y(e^{\beta y} - 1) \lambda(dy)$ .

□

*Remark 5.7.* Note that if  $\sigma^2 = 0$ ,  $\kappa(0) < \infty$ , then also  $A'(s) = i\mu_0 + i \int_{(-\infty, 0)} ye^{\beta y} \lambda(dy)$ ,  $s \in \mathbb{R}$ .

**5.5. Coercivity of  $|\psi^h - q|$ .** In this subsection we establish the following coercivity property:

- (C) There exists an  $h_0 \in (0, h_\star]$  and a  $B_0 \in (0, \infty)$ , such that for all  $h \in [0, h_0)$  and then all  $s \in (-\pi/h, \pi/h)$ , the following holds (recall  $\psi^0 = \psi$ ,  $\beta = \gamma + is$ ):

$$|\psi^h(\beta) - q| \geq B_0 |\beta|^\epsilon,$$

where  $\epsilon = 2$ , if  $\sigma^2 > 0$ ;  $\epsilon = 1$ , if  $\sigma^2 = 0$  and  $\kappa(0) < \infty$ ; finally, if  $\sigma^2 = 0$  and  $\kappa(0) = \infty$ , then  $\epsilon$  must satisfy (2) of Assumption 1.1 from the Introduction.

*Proof of (C).* (In the argument which follows, once again we refer the reader to expressions (2.1), (2.2) and (2.3).)

Suppose first  $\sigma^2 > 0$ , so that we work under scheme 1. Consider  $\psi(\beta)$ . The diffusion term is certainly quadratic in  $s$ . The drift term (viewed as a function of  $s$ ) is bounded by an affine function of  $|s|$ , and the Lévy measure integral has subquadratic growth in  $s$ , as can be seen immediately by the DCT and Lemma 5.3:

$$\lim_{R \rightarrow \infty} \sup_{\alpha \in \mathbb{C}^{\rightarrow}, |\alpha| \geq R} \frac{1}{|\alpha|^2} \left| \int_{(-\infty, 0)} (e^{\alpha y} - \alpha y \mathbb{1}_{[-V, 0)}(y) - 1) \lambda(dy) \right| = 0. \quad (5.14)$$

In addition  $(s \mapsto (\psi - q)(\beta))$  is bounded away from zero on bounded subsets of  $\mathbb{R}$ , by continuity and Proposition 5.1. This establishes the claim for  $h = 0$ .

To establish coercivity for  $\psi^h(\beta) - q$ ,  $h > 0$ , we proceed as follows. First, by (i) of Lemma 5.1, for any finite  $h_0 \in (0, h_\star]$ , there exists a  $B_0 \in (0, \infty)$ , such that for all  $h \in (0, h_0)$  and then all  $s \in (-\pi/h, \pi/h)$ :

$$\left| \frac{1}{2h} (e^{\beta h} - e^{-\beta h}) \right| \leq B_0 |\beta|.$$

This controls the term involving  $\mu$ . Next, by (vi) of Lemma 5.1, again for any finite  $h_0 \in (0, h_\star]$ , there are  $\{A_1, A_2\} \subset (0, \infty)$ , such that for all  $h \in (0, h_0)$  and then all  $s \in (-\pi/h, \pi/h)$ :

$$\left| -\mu^h \left( \frac{e^{\beta h} - e^{-\beta h}}{2h} - \beta \right) \right| \leq A_1 h^2 |\beta|^3 |\mu^h| \leq A_2 |\beta|^2 \zeta(h/2)$$

with  $\zeta(h/2) \rightarrow 0$  as  $h \downarrow 0$ . Further, just as in (5.14):

$$\lim_{R \rightarrow \infty} \sup_{\alpha \in \overline{\mathbb{C}^+}, |\alpha| \geq R} \sup_{h > 0} \frac{1}{|\alpha|^2} \left| \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h} (e^{\alpha y} - \alpha y \mathbb{1}_{[-V,0)}(z) - 1) \lambda(dz) \right| = 0,$$

where, additionally, one should note that for  $y \in \mathbb{Z}_h^-$  and  $z \in A_y^h$ ,  $|y| \leq 2|z|$ .

This, coupled with  $\sigma^2 > 0$  and (b) of Lemma 5.1, implies that there exist  $\{B_0, C_0\} \subset (0, \infty)$  and an  $h_0 \in (0, h_\star]$ , such that for all  $h \in (0, h_0)$  and then all  $s \in (-\pi/h, \pi/h) \setminus (-C_0, C_0)$ :  $|\psi^h(\beta) - q| \geq B_0 s^2$ . Finally, since, as  $h \downarrow 0$ ,  $\psi^h(\beta) - q \rightarrow \psi(\beta) - q$  uniformly in  $s$  belonging to bounded sets, and  $\psi(\beta) - q$  is bounded away from 0 on such sets, we obtain the asserted result.

Now suppose  $\sigma^2 = 0$  (so that scheme 2 is in effect) and consider first the case when  $\kappa(0) < \infty$ . With regard to  $\psi(\beta)$ , note that  $\mu_0 \beta$  is linear in  $s$ , whereas:

$$\lim_{R \rightarrow \infty} \sup_{\alpha \in \overline{\mathbb{C}^+}, |\alpha| \geq R} \frac{1}{|\alpha|} \left| \int_{(-\infty, 0)} (e^{\alpha y} - 1) \lambda(dy) \right| = 0, \quad (5.15)$$

by (i) of Lemma 5.3 and the DCT. The asserted coercivity follows immediately in the case  $h = 0$ .

To handle  $h > 0$ , it will be observed first that  $\mu - \mu^h \rightarrow \mu_0 > 0$  as  $h \downarrow 0$ , e.g. by the DCT. Also by the DCT, (i) of Lemma 5.3, and the fact that  $\kappa(0) < \infty$ :

$$\lim_{R \rightarrow \infty} \sup_{\alpha \in \overline{\mathbb{C}^+}, |\alpha| \geq R} \sup_{h > 0} \frac{1}{|\alpha|} \left| \sum_{y \in \mathbb{Z}_h^-} c_y^h (e^{\alpha y} - 1) \right| = 0.$$

Moreover, by (iv) of Lemma 5.1, for any finite  $h_0 \in (0, h_\star]$ , there exists  $A_0 \in (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $s \in (-\pi/h, \pi/h)$ ,

$$\left| c_0^h \left( \frac{e^{\beta h} + e^{-\beta h} - 2}{2h^2} \right) \right| \leq A_0 |\beta| \int_{[-h/2, 0)} |y| \lambda(dy)$$

with  $\int_{[-h/2, 0)} |y| \lambda(dy) \rightarrow 0$  as  $h \downarrow 0$ , since  $\kappa(0) < \infty$ . Coupled with (c) of Lemma 5.1, the asserted coercivity follows.

In the last instance, let  $\sigma^2 = 0$  and  $\kappa(0) = \infty$ . Necessarily,  $V = 1$  and Assumption 1.1 is in effect. We control first  $\Re \psi(\beta)$ . Clearly  $\gamma \mu$ ;  $\int_{(-\infty, -1)} (e^{\gamma y} \cos(sy) - 1) \lambda(dy)$  and  $\int_{[-1, 0)} (e^{\gamma y} - \gamma y - 1) \lambda(dy)$  are bounded in  $s$ , whereas (by (a) of Lemma 5.1 and (2) of Assumption 1.1):

$$\left| \int_{[-1, 0)} e^{\gamma y} (\cos(sy) - 1) \lambda(dy) \right| \geq B_1 s^2 \int_{[-\pi/|s|, 0)} y^2 \lambda(dy) \geq B_0 |s|^\epsilon,$$

for all  $s \in \mathbb{R}$  with  $|s| \geq K_0$ , and some  $\{K_0, B_0, B_1\} \subset (0, \infty)$ . Coercivity for  $\psi(\beta) - q$  follows.

Now we turn our attention to  $\psi^h(\beta)$  and again we control  $\Re \psi^h(\beta)$ . First observe that:

$$\underbrace{\frac{c_0^h}{2h^2} \left( e^{\gamma h} + e^{-\gamma h} - 2 \right) \cos(sh)}_{\text{bounded in } s} + \underbrace{\frac{c_0^h}{h^2} (\cos(sh) - 1)}_{=: (I) \leq 0}. \quad (5.16)$$

Next, with respect to the term involving the drift  $\mu$ , we refer to (ii) of Lemma 5.1 to obtain a linear bound in  $s$ . On the other hand we have:

$$\Re \left\{ \left( \frac{e^{\beta h} - 1}{h} - \beta \right) (-\mu^h) \right\} = \underbrace{\frac{e^{\gamma h} (\cos(sh) - 1)}{h} (-\mu^h)}_{\leq 0} + \underbrace{\frac{e^{\gamma h} - 1 - \gamma h}{h} (-\mu^h)}_{\text{bounded in } s},$$

since  $-\mu^h \leq 2\kappa(h/2)$  and  $\zeta(h/2) \rightarrow 0$  as  $h \downarrow 0$ . Finally, we consider the term:

$$\sum_{y \in \mathbb{Z}_h^{--}} c_y^h (e^{\gamma y} \cos(sy) - 1) - \sum_{y \in \mathbb{Z}_h^{--}} \gamma y \int_{A_y^h} \mathbb{1}_{[-1,0)}(z) \lambda(dz). \quad (5.17)$$

Certainly the part of (5.17) corresponding to  $\mathbb{1}_{(-\infty, -1)} \cdot \lambda$  is bounded in  $s$ . The part of (5.17) corresponding to  $\mathbb{1}_{[-1,0)} \cdot \lambda$  is:

$$\begin{aligned} & \sum_{y \in \mathbb{Z}_h^{--}} \int_{A_y^h \cap [-1,0)} \lambda(dz) [e^{\gamma y} \cos(sy) - \gamma y - 1] \\ = & \underbrace{\sum_{y \in \mathbb{Z}_h^{--}} \int_{A_y^h \cap [-1,0)} \lambda(dz) [e^{\gamma y} - \gamma y - 1]}_{\text{bounded in } s} + \underbrace{\sum_{y \in \mathbb{Z}_h^{--}} \int_{A_y^h \cap [-1,0)} \lambda(dz) e^{\gamma y} (\cos(sy) - 1)}_{=: (II) \leq 0}. \end{aligned} \quad (5.18)$$

Combining (I) and (II), we have, via (a) of Lemma 5.1, for some  $\{A_0, \beta_0, K_0, \alpha\} \subset (0, \infty)$ ,  $h_0 \in (0, h_\star]$ , for all  $h \in (0, h_0)$ , and then all  $s \in [-\pi/h, \pi/h] \setminus [-K_0, K_0]$ :

$$|\Re \psi^h(\beta)| \geq \beta_0 s^2 \left( c_0^h + \sum_{y \in \mathbb{Z}_h^{--}, -y \leq \pi/|s|} y^2 c_y^h \right) - A_0 |s|.$$

Also:

$$c_0^h + \sum_{y \in \mathbb{Z}_h^{--}, -y \leq \pi/|s|} y^2 c_y^h \geq \frac{4}{9} \int_{-\left(\left(\frac{\pi}{|s|} - \frac{h}{2}\right) \vee \frac{h}{2}\right), 0} u^2 \lambda(du) \geq \frac{4}{9} \int_{\left(-\frac{1}{2} \frac{\pi}{|s|}, 0\right)} u^2 \lambda(du),$$

since either  $\pi/|s| \geq h$ , in which case  $(\pi/|s|) - (h/2) \geq \frac{1}{2} \frac{\pi}{|s|}$ , or  $\pi/|s| \leq h$ , in which case  $\frac{h}{2} \geq \frac{1}{2} \frac{\pi}{|s|}$ .

Using now item (2) of Assumption 1.1, the required coercivity follows at once.  $\square$

**5.6. Brownian motion with drift** ( $\sigma^2 > 0 = \lambda(\mathbb{R})$ ). The scale functions can be calculated explicitly here, by using the recursive relations of Proposition 3.2. Then the following two proposition follow readily (essentially by Taylor expansions; recall also the notation from (5.7) and (5.8)):

**Proposition 5.5** ( $\sigma^2 > 0 = \lambda(\mathbb{R})$  ( $W^{(q)}$  convergence)). *Suppose  $\sigma^2 > 0 = \lambda(\mathbb{R})$  and let  $q \geq 0$ . If  $q \vee |\mu| = 0$ , then for all  $h \in (0, h_\star)$  and all  $x \in \mathbb{Z}_h^{++}$ :  $\Delta_W^{(q)}(x, h) = 0$ . If, however,  $q \vee |\mu| > 0$ , then:*

- (i) *There exist  $\{A_0, h_0\} \subset (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $x \in \mathbb{Z}_h^{++}$  with  $xh^2 \leq 1$ :*

$$\left| \Delta_W^{(q)}(x, h) \right| \leq A_0 h^2 (1+x) e^{\alpha x}.$$

(ii) For any nested sequence  $(h_n)_{n \geq 1} \downarrow 0$  and then any  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$ :

$$\lim_{n \rightarrow \infty} \frac{\Delta_W^{(q)}(x, h_n)}{h_n^2} = \frac{q^2}{2(\mu^2 + 2\sigma^2 q)} W^{(q)}(x) + \frac{x}{\sqrt{\mu^2 + 2\sigma^2 q}} (e^{\alpha+x\theta_+} - e^{\alpha-x\theta_-}).$$

(In particular, when  $q = 0$ , this limit is  $-\frac{2}{3} \frac{\mu^2 x}{(\sigma^2)^3} e^{-2\mu x/\sigma^2}$ .)

Here:

$$\begin{aligned} \alpha_{\pm} &:= \frac{-\mu \pm \sqrt{\mu^2 + 2q\sigma^2}}{\sigma^2} \\ \theta_{\pm} &:= \frac{\mu^3 \sqrt{2q\sigma^2 + \mu^2} \pm (\frac{1}{2}q^2(\sigma^2)^2 - \mu^4 - \mu^2\sigma^2 q)}{3(\sigma^2)^3 \sqrt{2q\sigma^2 + \mu^2}}. \end{aligned}$$

*Remark 5.8.* We note that for all  $x \geq 0$ :

$$W^{(q)}(x) = \frac{1}{\sqrt{\mu^2 + 2\sigma^2 q}} (e^{\alpha+x} - e^{\alpha-x})$$

(when  $q \wedge |\mu| > 0$ ) and  $W^{(q)}(x) = x$  (otherwise). Observe also that, unless  $q = 0$ ,  $\alpha_{\pm} \in \pm(0, \infty)$ .

**Proposition 5.6** ( $\sigma^2 > 0 = \lambda(\mathbb{R})$  ( $Z^{(q)}$  convergence)). *Suppose  $\sigma^2 > 0 = \lambda(\mathbb{R})$ , let  $q > 0$ .*

(i) *There exist  $\{A_0, h_0\} \subset (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $x \in \mathbb{Z}_h^{++}$  with  $xh^2 \leq 1$ :*

$$\left| \Delta_Z^{(q)}(x, h) \right| \leq A_0 [h^2(1+x)e^{\alpha+x} + h(e^{\alpha+x} - e^{\alpha-x})].$$

(ii) *For any nested sequence  $(h_n)_{n \geq 1} \downarrow 0$  and then any  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$ :*

$$\lim_{n \rightarrow \infty} \frac{\Delta_Z^{(q)}(x, h_n)}{h_n} = -\frac{1}{2} \frac{q}{\sqrt{\mu^2 + 2\sigma^2 q}} (e^{\alpha+x} - e^{\alpha-x}).$$

**5.7. Non-trivial diffusion component.** We consider the convergence when  $\sigma^2 > 0$ . This case is relatively straightforward, as coercivity is very strong (namely, quadratic). Note that  $\delta^0 = 1$  and we work under scheme 1.

**Proposition 5.7** ( $\sigma^2 > 0$  ( $W^{(q)}$  convergence)). *Suppose  $\sigma^2 > 0$  and let  $q \geq 0$ .*

(i) *For any  $\gamma > \Phi(q)$ , there are  $\{A_0, h_0\} \subset (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $x \in \mathbb{Z}_h^{++}$ :*

$$\left| \Delta_W^{(q)}(x, h) \right| \leq A_0 (h + \zeta(h/2) + \xi(h/2)h \log(1/h)) e^{\gamma x}.$$

*In particular, if  $\kappa(0) < \infty$ , then  $\left| \Delta_W^{(q)}(x, h) \right| \leq A_0 h e^{\gamma x}$  and under Assumption 1.1,*

$$\left| \Delta_W^{(q)}(x, h) \right| \leq A_0 h^{2-\epsilon} e^{\gamma x}.$$

(ii) *There exist:*

(a) *a Lévy triplet  $(\sigma^2, \lambda, \mu)$  with  $\sigma^2 \neq 0$  and  $0 < \kappa(0) < \infty$ ;*

(b) *for each  $\epsilon \in (1, 2)$  a Lévy triplet  $(\sigma^2, \lambda, \mu)$  with  $\sigma^2 \neq 0$  and  $\lambda(-1, -\delta) \sim 1/\delta^\epsilon$  as  $\delta \downarrow 0$ ;*



and then in each of the cases (a)-(b) a nested sequence  $(h_n)_{n \geq 1} \downarrow 0$  such that for each  $q \geq 0$  there is an  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$  with:

$$\liminf_{n \rightarrow \infty} \frac{|\Delta_W^{(q)}(x, h_n)|}{h_n \vee \zeta(h_n)} > 0,$$

where  $h_n \vee \zeta(h_n) \sim h_n$ , if  $\kappa(0) < \infty$  and  $\sim h_n^{2-\epsilon}$ , if  $\kappa(0) = \infty$  as  $n \rightarrow \infty$ .

*Remark 5.9.* Note that if  $\lambda(-1, \delta) \sim 1/\delta^\epsilon$  as  $\delta \downarrow 0$ , with  $\epsilon \in (1, 2)$ , then (as  $h \downarrow 0$ )  $\xi(h/2) \sim h^{2-\epsilon}$ , and  $\zeta(h/2) \sim h^{2-\epsilon}$ , so that  $h\xi(h/2) \log(1/h) = o(h\kappa(h/2))$ . See Proposition 5.3. More generally, Assumption 1.1 is fulfilled if  $\lambda(-1, -\delta) \sim \delta^{-\epsilon} l(\delta)$  where  $0 < \liminf_{0+} l < \limsup_{0+} l < +\infty$  (see Lemma 5.5).

*Remark 5.10.* Under Assumption 1.1, it follows that  $\zeta(h/2) + \xi(h/2) = O(h^{2-\epsilon})$  as  $h \downarrow 0$  (see again Lemma 5.5).

*Proof.* First, with respect to (i), we have as follows. (a) of (5.9) is seen immediately to be of order  $O(h)$  by coercivity (C); whereas (b) of (5.9) is of order  $O(h + h\xi(h/2) \log(1/h) + V\zeta(h/2))$  by coercivity (C) and the estimate of the absolute difference  $|\psi^h - \psi|$  ( $\Delta_2$ ). Since  $\delta^0 = 1$ , (c) of (5.9) is void.

Second we prove (ii).

- We consider first (a). Take  $\lambda = \delta_{-1/2}$ ,  $h_n = 1/3^n$  ( $n \geq 1$ ),  $\mu = 0$ ,  $\sigma^2 = 1$ ,  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$  ( $x$  is now fixed!). The goal is to establish no better than linear convergence in this case.

Now, (a) of (5.9) is actually of order  $O(h^2)$ . Indeed, in the difference to replacing  $\psi(\beta) - q$  by  $-\frac{1}{2}\sigma^2 s^2$ , this is seen immediately to be even of order  $O(h^3)$ , by coercivity (C) and a simple  $|\int \cdot| \leq \int |\cdot|$  argument. On the other hand:

$$\lim_{T \rightarrow \infty} \int_{(-T, T) \setminus (-\pi/h, \pi/h)} \frac{e^{isx}}{s^2} ds = O(h^2).$$

This is so by an integration by parts argument, writing:

$$\frac{d}{ds} \left( \frac{e^{isx}}{s^2} \right) = \frac{ixe^{isx}}{s^2} - \frac{2e^{isx}}{s^3}.$$

We can thus focus on (b) of (5.9). Consider there the difference:

$$(\psi^h - \psi)(\beta) = \overbrace{\frac{\sigma^2}{2h^2} (e^{\beta h} + e^{-\beta h} - 2) - \frac{1}{2}\sigma^2 \beta^2}^{(b.1)} + \overbrace{\int_{(-\infty, 0)} (e^{\beta y} - 1) \lambda(dy) - \sum_{y \in \mathbb{Z}_h^{--}} c_y^h (e^{\beta y} - 1)}^{(b.2)}. \quad (5.19)$$

The part of (b) in (5.9) corresponding to (b.1) is, in the difference to the analogous term for Brownian motion without drift, bounded (up to a non-zero multiplicative constant) by:

$$\int_{[-\pi/h, \pi/h]} ds \frac{h^2 |\beta|^4 |\beta|^2}{|\beta|^8} = O(h^2)$$

(this follows by (iv) and (v) of Lemma 5.1, the fact that  $e^{\beta y} - 1$  is uniformly bounded by 2, and by coercivity (C)). Since the term corresponding to just Brownian motion is shown to be of order  $O(h^2)$  itself (see Proposition 5.5), we can thus focus on (b.2). The latter is  $e^{-\beta/2} - e^{-(1-h_n)\beta/2} = e^{-\beta/2}(1 - e^{\beta h_n/2})$ . In the difference to replacing  $1 - e^{\beta h_n/2}$  by  $-\beta h_n/2$ , a term of order  $O(h_n^2)$  emerges in  $\Delta_W^{(q)}(x, h_n)$ , this by (iii) of Lemma 5.1, and coercivity (C). Hence it is sufficient to study:

$$\frac{1}{2\pi i} \int_{[-\pi/h_n, \pi/h_n]} e^{\beta x} \frac{e^{-\beta/2} \beta}{(\psi - q)(\beta)(\psi^{h_n} - q)(\beta)} ds$$

which we would like bounded away from 0, as  $n \rightarrow \infty$ . Now, by coercivity (C), and the DCT, this expression in fact converges to:

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{\beta(x-1/2)} \beta}{(\psi - q)^2(\beta)} ds =: g(x).$$

Note that  $g$  is continuous in its parameter  $x \in [0, \infty)$  by the DCT. Moreover,  $g$  cannot vanish identically on  $\cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$ , since then it would do so on  $\mathbb{R}_+$  by continuity. But this cannot be. Naively, since in  $g$  we are looking at the inverse Laplace transform of a non-vanishing function  $T$ . Formally, one performs a Laplace transform of  $g$ , and concludes, via Fubini and Cauchy's Residue Theorem (recalling the quadratic behaviour of  $\psi(\beta)$  as  $|\beta| \rightarrow \infty$  over  $\overline{\mathbb{C}^-}$ , see (5.14)) that  $T = \hat{g}$ , where  $T(\alpha) := \frac{e^{-\alpha/2} \alpha}{(\psi - q)^2(\alpha)}$  ( $\alpha > \gamma$ ). Then  $g$  vanishing implies the same of  $T$ , which is a clear contradiction.

- Consider now (b). We are seeking to establish strictly worse than linear convergence here, since  $\kappa(0) = \infty$ .

For sure (a) in (5.9) is of order  $O(h)$ . When it comes to (b), consider its decomposition, in the numerator of the integrand according to items (1)-(6) from Subsection 5.3. Now, (1) thus yields in (b) a term of order  $O(h)$ ; (2) one of order  $o(h)$ ; with respect to (3) we will choose a  $\lambda$  which falls under the scope of Remark 5.9 and hence this will contribute a term of order  $o(\zeta(h/2))$ ; (4) gives a term of order  $o(h)$ ; (5) contributes as  $o(h\kappa(h/2))$ ; whereas finally (6) yields a term of order  $O(h)$  in the part corresponding to  $\mathbb{1}_{(-\infty, 1)} \cdot \lambda$  and the part corresponding to  $\mathbb{1}_{[-1, 0)} \cdot \lambda$  is where we will get sharpness of the rate from.

So we take  $\sigma^2 = 1$ ,  $\mu = 0$ ,  $h_n = 1/3^n$  ( $n \geq 1$ ),  $\lambda = \sum_{k=1}^{\infty} w_k \delta_{-x_k}$ ,  $x_k = \frac{3}{2} h_k$  and  $w_k = 1/x_k^\epsilon$  ( $k \geq 1$ ). Clearly  $\kappa(0) = \infty$  and by checking it on the decreasing sequence  $(h_n)_{n \geq 1}$  it is clear that  $\lambda(-1, -\delta) \sim \delta^{-\epsilon}$  as  $\delta \downarrow 0$ . Moreover,

$$\left| \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h \cap [-1, 0)} \left( e^{\beta y} - \beta y - (e^{\beta z} - \beta z) - \frac{1}{2} \beta^2 (y^2 - z^2) \right) \lambda(dz) \right|$$

yields in (5.9), by (ii) of Lemma 5.4 and coercivity (C), a term of order  $O(h \log(1/h)) = o(h^{2-\epsilon})$ . Therefore it is sufficient to study:

$$\sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h \cap [-1, 0)} \beta^2(y^2 - z^2) \lambda(dz) = \sigma_2 - \sigma_1,$$

where:

$$\sigma_1 := \int_{[-1, -h_n/2)} u^2 \lambda(du) = \sum_{k=1}^n x_k^2 w_k, \quad \sigma_2 := \sum_{k=1}^n (x_k - h_n/2)^2 w_k$$

and hence  $\sigma_1 - \sigma_2 = 2\zeta(h_n/2) - \gamma(h_n/2) \geq \zeta(h_n/2)$ . Moreover,

$$\int_{[-\pi/h_n, \pi/h_n]} e^{isx} \frac{\beta^2}{[(\psi - q)(\psi^{h_n} - q)](\beta)} ds \rightarrow \int_{\mathbb{R}} e^{isx} \frac{\beta^2}{(\psi - q)^2(\beta)} ds$$

as  $n \rightarrow \infty$  by the DCT. By the usual arguments, this integral does not vanish simultaneously in all  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$ , whence tightness obtains.  $\square$

**Proposition 5.8** ( $\sigma^2 > 0$  ( $Z^{(q)}$  convergence)). *Suppose  $\sigma^2 > 0$ , let  $q > 0$ .*

(i) *For any  $\gamma > \Phi(q)$ , there are  $\{A_0, h_0\} \subset (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $x \in \mathbb{Z}_h^{++}$ :*

$$\left| \Delta_Z^{(q)}(x, h) \right| \leq A_0 (h + \zeta(h/2)) e^{\gamma x}.$$

*In particular, if  $\kappa(0) < \infty$ , then  $\left| \Delta_Z^{(q)}(x, h) \right| \leq A_0 h e^{\gamma x}$  and otherwise  $\left| \Delta_Z^{(q)}(x, h) \right| \leq A_0 \zeta(h/2) e^{\gamma x}$ .*

(ii) (a) *There exists a nested sequence  $(h_n)_{n \geq 1} \downarrow 0$ , such that for any  $q > 0$  and any  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$ , there exists a Lévy triplet  $(\sigma^2, \mu, \lambda)$  with  $\sigma^2 > 0$  and  $0 < \kappa(0) < \infty$ , and such that:*

$$\liminf_{n \rightarrow \infty} \frac{\left| \Delta_Z^{(q)}(x, h_n) \right|}{h_n} > 0.$$

(b) *There exists for each  $\epsilon \in (1, 2)$  a Lévy triplet  $(\sigma^2, \lambda, \mu)$  with  $\sigma^2 > 0$  and  $\lambda(-1, \delta) \sim 1/\delta^\epsilon$  as  $\delta \downarrow 0$ , and a nested sequence  $(h_n)_{n \geq 1} \downarrow 0$  such that for each  $q > 0$ , there is an  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$  with:*

$$\liminf_{n \rightarrow \infty} \frac{\left| \Delta_Z^{(q)}(x, h_n) \right|}{\zeta(h_n)} > 0.$$

*Proof.* With respect to (i), we have as follows. First, (a) of (5.10) is  $O(h^2)$  and (b)  $O(h)$  by coercivity (C) (and by Lemma 5.2 in the case of (b)). Second, (c) is  $O(h + \zeta(h/2))$  by coercivity (C) and the estimate  $(\Delta_2)$ .

Next we show (ii).

- We consider first (a). Take  $\sigma^2 = 1$ ,  $\mu = 0$  and  $\lambda = \alpha \delta_{-1}$ , where we fix  $\alpha > 0$ . The idea is to note that convergence is ‘tightly linear’ in the Brownian motion case (see Proposition 5.6), and then to show that by taking  $\alpha$  small enough, we do not spoil this.

Now, remark that:

- As  $\alpha \downarrow 0$ ,  $\psi$  is nondecreasing, hence  $\Phi(q)$  is nonincreasing, and so a  $\gamma > \Phi(q)$  can be chosen, uniformly in all  $\alpha$  bounded.
- Moreover, the presence of  $\alpha$  does not affect coercivity, which is in addition uniform in all  $\alpha$  small enough. Indeed, just take  $\gamma > \Phi^{BM}(q)$ , where  $\Phi^{BM}$  corresponds to the Brownian motion part of this Lévy process. Then  $|\psi_{BM}^h(\beta) - q| \geq B_0|\beta|^2$ , for all  $s \in (-\pi/h, \pi/h)$ , for all  $h \in [0, h_0)$ , for some  $\{B_0, h_0\} \subset (0, \infty)$ . The part of  $\psi^h(\beta)$  corresponding to the CP term is bounded uniformly in  $s$  and all  $h$  small enough (including 0), and moreover, scales with  $\alpha$ . Hence there are  $\{B_0, h_0, \alpha_0\} \subset (0, \infty)$  such that for all  $\alpha \in (0, \alpha_0)$ :  $|\psi^h(\beta) - q| \geq B_0|\beta|^2$ , for all  $s \in (-\pi/h, \pi/h)$ , for all  $h \in [0, h_0)$ .

Take now  $h_n = 1/2^n$  ( $n \geq 1$ ), and a fixed  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++}$ . We show that (5.10), when looked in the difference to the analogous expression for the Brownian motion part, is of order  $o(h) + \alpha O(h)$  (i.e. bounded in absolute value by terms either decaying faster than linear, or else with a coefficient that scales with  $\alpha$ ). Indeed, the difference of (b)s in (5.10) follows readily as being  $\alpha O(h)$ . In addition, (a) of (5.10) is of order  $O(h^2)$ . Also (c) is of order  $o(h)$  except in the part corresponding to the CP term, which is itself  $\alpha O(h)$ . Thus, choosing  $\alpha$  small enough, the desired sharpness obtains. However, at least in principle, the choice of  $\alpha$  depends on  $q$  and  $x$ , hence note the formulation of the proposition.

- Consider now (b). Clearly here the same example works as for the functions  $W^{(q)}$ : the presence of the extra  $1/\beta$  in the integrand of (c) of (5.10) is of no consequence (if anything, beneficial), and the rest of the terms are of order  $O(h)$  anyhow.

□

5.8.  $\sigma^2 = 0$ , **finite variation paths**. In this subsection we study the convergence when  $\sigma^2 = 0$  and  $\kappa(0) < \infty$ . Note that in this case necessarily  $\mu_0 > 0$ , whereas  $\delta^0 = 0$ , and we work under scheme 2. The estimates are delicate here, since coercivity is weak (namely, linear).

We make the following key observation. By decomposing:

$$A(s) = \psi(\beta) - q = \underbrace{\mu_0\gamma + \int (e^{\gamma y} \cos(sy) - 1)\lambda(dy)}_{=:A_e(s)} - q + \underbrace{is\mu_0 + i \int e^{\gamma y} \sin(sy)\lambda(dy)}_{=:A_o(s)} \quad (5.20)$$

into its even and odd part, it can be shown that, crucially,

$$\int_1^\infty \frac{ds}{s^2} |A_e(s)| < \infty. \quad (5.21)$$

See Proposition 5.2. On the other hand, a similar argument to (5.15), shows that  $|A_o(s)| \geq B_0|s|$  for all  $s \notin [a_0, a_0]$ , for some  $\{a_0, B_0\} \subset \mathbb{R}$ . We shall refer to the latter property as “coercivity of  $A_o$ ”.

Note also that by the DCT,  $\mu - \mu^h \rightarrow \mu_0$  as  $h \downarrow 0$ .

We next prove a key lemma. While (1) and (3) thereof will both be used explicitly in the sequel, the same cannot be said of (2). Nevertheless, the proof of the latter is instructive of the techniques which we will be using, and so (also for the sake of completeness) we choose to keep it.

**Lemma 5.6.** *Suppose  $\sigma^2 = 0$ ,  $\kappa(0) < \infty$ . Let  $\{l, a, b, M\} \subset \mathbb{N}_0$  and let  $h_0 \in (0, h_\star]$  be given by the coercivity condition (C).*

(1) *If  $a + b + l \geq M + 1$ , then:*

$$\sup_{(h,z) \in (0, h_0) \times \mathbb{R}} \left| \int_{[-\frac{\pi}{h}, \frac{\pi}{h}]} e^{isz} \frac{(h \wedge 1)^{l-l \wedge (M+1)} h^{l \wedge (M+1)} s^M}{A(s)^a A^h(s)^b} ds \right| < \infty.$$

(2) *If only  $a + b + l \geq M$ , then:*

$$\sup_{(h,z) \in (0, h_0 \wedge K) \times \mathbb{R}} \left| z \int_{[-\frac{\pi}{h}, \frac{\pi}{h}]} e^{isz} \frac{h^l s^M}{A(s)^a A^h(s)^b} ds \right| < \infty$$

for any  $K \in (0, \infty)$ .

(3) *If even  $a + b + l \geq M + 2$ , then:*

$$\sup_{(h,x,z) \in (0, h_0) \times \mathbb{R} \times (\mathbb{R} \setminus \{0\})} \left| \frac{1}{z} \int_{[-\frac{\pi}{h}, \frac{\pi}{h}]} e^{isx} \frac{(e^{isz} - 1)(h \wedge 1)^{l-l \wedge (M+2)} h^{l \wedge (M+2)} s^M}{A(s)^a A^h(s)^b} ds \right| < \infty.$$

*Remark 5.11.* Suppose  $l = b = 0$  and  $a = M + 1$  (respectively  $a = M$ ,  $a = M + 2$ ) for simplicity (indeed the proof of (1) and (3) will be reduced to this case using (C) and  $(\Delta_2)$ , whereas (2) will (essentially) follow by an application of (1)). Then for large  $|s|$ , the integrand in (1) (respectively (2), (3)) behaves as  $\sim e^{isz}/s$  (respectively  $e^{isz}$ ,  $e^{isx}(e^{isz} - 1)/s^2$ ) in the variable  $s$ . It is then not surprising that the proof of the claims is essentially a modification of the argument implying that (in the sense of Cauchy's principal values, as appropriate)  $\int_{[-\pi/h, \pi/h]} (e^{isz}/s) ds$ ,  $\int_{[-\pi/h, \pi/h]} z e^{isz} ds$  and  $\int_{[-\pi/h, \pi/h]} (e^{isx}(e^{isz} - 1)/s^2) ds/z$  are bounded in the relevant suprema (as they are).

*Proof.* We use in the proof, without explicit reference, as has already often been the case, the observation that  $|sh| \leq \pi$  within the integration domain and the basic  $|\int \cdot ds| \leq \int |\cdot| ds$ -argument; it will be clear whenever these are being applied. Further, we sometimes employ, but do not always refer to, some elementary trigonometric inequalities, specifically  $|\sin(v)| \leq |v|$ ,  $1 - \cos(v) \leq v^2/2$  ( $v \in \mathbb{R}$ ), as well as (i) of Lemma 5.3. Finally, note that the integrands in the formulation of this lemma are certainly locally bounded, by coercivity (C), and (hence) the integrals well-defined. Indeed, by the same token, it is only non-trivial to show the finiteness of the suprema in (1) and (3) for  $h \in (0, h_0 \wedge 1)$ , a restriction which we therefore make outright. In this respect, note that in (2) the restriction to bounded  $h$  is made a priori.

Consider now first (1). By coercivity (C), it is assumed without loss of generality that  $l \leq M$ , then  $a + b = M - l + 1$  and finally,  $l = 0$ .

Next, it will be sufficient to consider the case when  $b = 0$ , since  $A(s)$  can then be successively replaced in the denominator by  $A^h(s)$  modulo a quantity, which, using coercivity (C) and the estimates  $(\Delta_2)$ , remains bounded (in the supremum over  $(h, z) \in (0, h_0 \wedge 1) \times \mathbb{R}$ ).

Recall now the decomposition  $A = A_e + A_o$  of (5.20). With  $a_0$  as above, it will furthermore be sufficient to establish (1) with the integration region  $[-\pi/h, \pi/h] \setminus [-a_0, a_0]$  in place of  $[-\pi/h, \pi/h]$ , the integrand being locally bounded in the supremum over  $(h, z) \in (0, h_0 \wedge 1) \times \mathbb{R}$  (again by (C)). Moreover, one can then successively replace  $A(s)$  by  $A_o(s)$ , using coercivity of  $A_o$  and (C), as well as (5.21). Again this is done modulo a term which remains bounded in the supremum over  $(h, z) \in (0, h_0 \wedge 1) \times \mathbb{R}$ . Hence we need only establish the finiteness of the quantity:

$$\sup_{(h,z) \in (0, h_0 \wedge 1) \times \mathbb{R}} \left| \int_{[-\frac{\pi}{h}, \frac{\pi}{h}] \setminus [-a_0, a_0]} e^{isz} \frac{s^M}{A_o(s)^{M+1}} ds \right|.$$

Owing to the fact that the quotient in the integrand is odd in  $s$ , we may clearly restrict the supremum to  $z \in \mathbb{R} \setminus \{0\}$ , replacing also  $e^{isz}$  by  $\sin(sz)$  therein. A change of variables  $u = s|z|$  then leads us to consider:

$$\int_{[-\frac{\pi}{h}|z|, \frac{\pi}{h}|z|] \setminus [-a_0|z|, a_0|z|]} \sin(u) \frac{u^M}{(|z|A_o(u/|z|))^{M+1}} du, \quad (5.22)$$

whose finiteness in the supremum over  $(h, z) \in (0, h_0 \wedge 1) \times \mathbb{R} \setminus \{0\}$  we seek to establish. Let  $\mathcal{A} := [-\frac{\pi}{h}|z|, \frac{\pi}{h}|z|] \setminus [-a_0|z|, a_0|z|]$ . By coercivity of  $A_o$ , and since  $|\sin(u)| \leq |u|$  ( $u \in \mathbb{R}$ ), we do indeed get a finite quantity for the integral (in (5.22)) over  $\mathcal{A} \cap [-1, 1]$ . On the other hand, to handle the rest of the domain,  $\mathcal{A} \setminus [-1, 1]$ , we resort to integration by parts;

$$\begin{aligned} \frac{d}{du} \left( \cos(u) \frac{u^M}{(|z|A_o(u/|z|))^{M+1}} \right) &= -\sin(u) \frac{u^M}{(|z|A_o(u/|z|))^{M+1}} + \\ &+ \cos(u) \left( \frac{Mu^{M-1}}{(|z|A_o(u/|z|))^{M+1}} - \frac{u^M(M+1)A'_o(u/|z|)}{(|z|A_o(u/|z|))^{M+2}} \right). \end{aligned}$$

Now, once integration over  $\mathcal{A} \setminus [-1, 1]$  has been performed, on the left-hand side, a bounded quantity obtains, by coercivity of  $A_o$ . On the right-hand side we obtain from the first term the desired quantity (modulo the sign), whereas what emerges from the second term is bounded by coercivity of  $A_o$ , and the boundedness of  $A'_o$  (see  $(\Delta'_1)$  in Subsection 5.4).

We next consider (2). Here an integration by parts must be done outright, thus:

$$\frac{d}{ds} \left( e^{isz} \frac{h^l s^M}{A(s)^a A^h(s)^b} \right) = iz e^{isz} \frac{h^l s^M}{A(s)^a A^h(s)^b} + e^{isz} \left( \frac{h^l M s^{M-1}}{A(s)^a A^h(s)^b} \right) - e^{isz} \left( \frac{h^l s^M a A'(s)}{A(s)^{a+1} A^h(s)^b} + \frac{h^l s^M b A^h(s)}{A(s)^a A^h(s)^{b+1}} \right).$$

Further, once integration over  $[-\pi/h, \pi/h]$  has been performed in this last equality, on the left-hand side a bounded quantity obtains by coercivity (C). On the right-hand side, the first term yields the desired quantity (modulo a non-zero multiplicative constant), and the second is bounded by part (1). Now, using (5.12) and (5.13), via Fubini's Theorem, part (1) again, and by elementary estimates such as  $e^{\gamma h}$  being bounded for  $h$  bounded,  $|\sum_{y \in \mathbb{Z}_h^-} y c_y^h e^{\beta y}| \leq 2e^{\gamma h/2} \int |y| e^{\gamma y} \lambda(dy)$ ,  $c_0^h = O(h)$  and (i) of Lemma 5.1, the claim obtains.

Finally, we are left to consider (3). Again by coercivity (C), it is assumed without loss of generality that  $l \leq M$ , then  $a + b = M - l + 2$  and finally,  $l = 0$ . Moreover, by the same argument

as for (1), we may further insist on  $b = 0$ , replace the integration region by  $[-\pi/h, \pi/h] \setminus [-a_0, a_0]$  and finally  $A$  by  $A_o$ . Thus we are left to analyse:

$$\frac{1}{z} \int_{[-\frac{\pi}{h}, \frac{\pi}{h}] \setminus [-a_0, a_0]} e^{isx} \frac{(\cos(sz) - 1)s^M}{A_o(s)^{M+2}} ds \quad \text{and} \quad \frac{1}{z} \int_{[-\frac{\pi}{h}, \frac{\pi}{h}] \setminus [-a_0, a_0]} e^{isx} \frac{\sin(sz)s^M}{A_o(s)^{M+2}} ds, \quad (5.23)$$

which we require both to be bounded in the relevant supremum.

In the first integral of (5.23) make the substitution  $v = s|z|$  to obtain:

$$\text{sgn}(z) \int_{[-\frac{\pi}{h}|z|, \frac{\pi}{h}|z|] \setminus [-a_0|z|, a_0|z|]} e^{ivx/|z|} \frac{(\cos(v) - 1)v^M}{(|z|A_o(v/|z|))^{M+2}} dv.$$

Letting, as usual,  $\mathcal{A} := [-\frac{\pi}{h}|z|, \frac{\pi}{h}|z|] \setminus [-a_0|z|, a_0|z|]$ , the integral over  $\mathcal{A} \cap [-1, 1]$  (respectively  $\mathcal{A} \cap [-1, 1]$ ) is bounded by coercivity of  $A_o$  (respectively the latter and since  $1 - \cos(v) \leq v^2/2$ ).

On the other hand, in the second integral of (5.23), note that by the oddness of  $A_o$  only  $\sin(sx)$  makes a non-zero contribution. Then we may assume  $x \neq 0$  and make the substitution  $u = s|x|$  to arrive at:

$$\frac{x}{z} \int_{[-\frac{\pi}{h}|x|, \frac{\pi}{h}|x|] \setminus [-a_0|x|, a_0|x|]} \frac{\sin(u) \sin(uz/|x|)u^M}{(|x|A_o(u/|x|))^{M+2}} du.$$

Let again  $\mathcal{A} := [-\frac{\pi}{h}|x|, \frac{\pi}{h}|x|] \setminus [-a_0|x|, a_0|x|]$  be the domain of integration. It is clear that the integral over  $\mathcal{A} \cap [-1, 1]$  is finite, using coercivity of  $A_o$  and twice  $|\sin(w)| \leq |w|$  ( $w \in \mathbb{R}$ ). To handle the remainder of the domain,  $\mathcal{A} \setminus [-1, 1]$ , we use one last time integration by parts, thus;

$$\begin{aligned} \frac{x}{z} \frac{d}{du} \left( \frac{\cos(u) \sin(uz/|x|)u^M}{(|x|A_o(u/|x|))^{M+2}} \right) &= -\frac{x}{z} \frac{\sin(u) \sin(uz/|x|)u^M}{(|x|A_o(u/|x|))^{M+2}} + \text{sgn}(x) \frac{\cos(u) \cos(uz/|x|)u^M}{(|x|A_o(u/|x|))^{M+2}} + \\ &\quad \frac{x}{z} \frac{\cos(u) \sin(uz/|x|)Mu^{M-1}}{(|x|A_o(u/|x|))^{M+2}} - \frac{x}{z} \frac{\cos(u) \sin(uz/|x|)u^M (M+2)A'_o(u/|x|)}{(|x|A_o(u/|x|))^{M+3}}. \end{aligned}$$

The claim now obtains by coercivity of  $A_o$ , boundedness of  $A'_o$  ( $\Delta'_1$ ) and by using the elementary estimate  $|\sin(w)| \leq |w|$  ( $w \in \mathbb{R}$ ), as appropriate.  $\square$

**Proposition 5.9** ( $\sigma^2 = 0$ ,  $\kappa(0) < \infty$  ( $W^{(q)}$  convergence)). *Suppose  $\sigma^2 = 0$  and  $\kappa(0) < \infty$ . Let  $q \geq 0$ .*

(i) *For any  $\gamma > \Phi(q)$  there are  $\{A_0, h_0\} \subset (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $x \in \mathbb{Z}_h^+$ :*

$$\left| \Delta_W^{(q)}(x, h) \right| \leq A_0 \frac{h}{x} e^{\gamma x}.$$

(ii) *For the Lévy triplet  $(0, \delta_{-1}, 1)$  and the nested sequence  $(h_n := 1/2^n)_{n \geq 1} \downarrow 0$ , for each  $q \geq 0$  and any  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^+ \cap [0, 1)$ :*

$$\lim_{n \rightarrow \infty} \frac{\Delta_W^{(q)}(x, h_n)}{h_n} = e^{x(1+q)} \frac{1}{2} (1+q)^2 x.$$

*Proof.* With respect to (i), we estimate the three terms appearing on the right-hand side of (5.9) one by one.

First, (a) in (5.9) is easily seen to be of order  $\frac{1}{x}O(h)$  by an obvious integration by parts argument, using coercivity and the fact that  $A'$  is bounded.

Second, when it comes to (b) in (5.9), an integration by parts is also performed immediately:

$$\frac{d}{ds} \left( e^{isx} \left( \frac{1}{A(s)} - \frac{1}{A^h(s)} \right) \right) = ix e^{isx} \left( \frac{1}{A(s)} - \frac{1}{A^h(s)} \right) + e^{isx} \left( -\frac{A'(s)}{A(s)^2} + \frac{A^{h'}(s)}{A^h(s)^2} \right).$$

Upon integration on  $[-\pi/h, \pi/h]$ , by coercivity, the left-hand side is of order  $O(h)$  and hence will contribute  $\frac{1}{x}O(h)$  to the right-hand side of (5.9). Write:

$$-\frac{A'(s)}{A(s)^2} + \frac{A^{h'}(s)}{A^h(s)^2} = \underbrace{\frac{A^{h'}(s) - A'(s)}{A(s)^2}}_{(1)} + \underbrace{A^{h'}(s) \frac{(A(s) - A^h(s))(A(s) + A^h(s))}{A(s)^2 A^h(s)^2}}_{(2)}. \quad (5.24)$$

We focus on each term one at a time. For (1), there corresponds to it, modulo non-zero multiplicative constants, and by Fubini (using (5.12) and (5.13)):

$$\begin{aligned} & (\mu - \mu^h) \int_{[-\pi/h, \pi/h]} \frac{e^{\beta h} - 1}{A(s)^2} e^{isx} ds + c_0^h \int_{[-\pi/h, \pi/h]} \left( \frac{e^{\beta h} - e^{-\beta h}}{2h} \right) \frac{1}{A(s)^2} e^{isx} ds \\ & + \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h} \lambda(dz) \int_{[-\pi/h, \pi/h]} \frac{y(e^{\beta y} - \mathbb{1}_{[-V,0)}(z)) - z(e^{\beta z} - \mathbb{1}_{[-V,0)}(z))}{A(s)^2} e^{isx} ds. \end{aligned} \quad (5.25)$$

There are three summands in (5.25). The first is  $O(h)$  by employing the decomposition  $e^{\beta h} - 1 = (e^{\beta h} - \beta h - 1) + \beta h$  and then using (iii) of Lemma 5.1 and coercivity (C) (respectively (1) of Lemma 5.6) for the first (respectively second) term. The same is true of the second summand, noting that, for sure,  $c_0^h = O(h)$ , employing the decomposition  $\frac{e^{\beta h} - e^{-\beta h}}{2h} = \left( \frac{e^{\beta h} - e^{-\beta h}}{2h} - \beta \right) + \beta$  and then using (vi) of Lemma 5.1 and again coercivity (C) (respectively (1) of Lemma 5.6) for the first (respectively second) term. As for the third summand of (5.25), write when  $z \notin [-V, 0)$ :

$$ye^{\beta y} - ze^{\beta z} = \overbrace{(y-z)e^{\beta y}}^{(I)} + ze^{\beta z} \overbrace{(e^{\beta(y-z)} - \beta(y-z) - 1)}^{(II)} + \overbrace{\beta(y-z)}^{(III)}.$$

By the findings of Lemma 5.1, the fact that  $\lambda(-\infty, -V) < \infty$  and coercivity (C), it is clear that (I) and (II) will contribute a term of order  $\frac{1}{x}O(h)$  to the right-hand side of (5.9). On the other hand, for (III), the same follows by (1) of Lemma 5.6. Next, when  $z \in [-V, 0)$  we write:

$$ye^{\beta y} - ze^{\beta z} - (y-z) = \overbrace{e^{\beta z} y (e^{\beta(y-z)} - 1)}^{(I)} + \overbrace{(y-z)(e^{\beta z} - 1)}^{(II)}. \quad (5.26)$$

When it comes to (I), it is dealt with precisely as it was for  $z \notin [-V, 0)$  (but note also that  $|y| \leq 2|z|$ ). With regard to (II), apply (3) of Lemma 5.6.

To handle (2) of (5.24), i.e.:

$$A^{h'}(s) \frac{(A(s) - A^h(s))(A(s) + A^h(s))}{A(s)^2 A^h(s)^2},$$

notice that in the difference to replacing this with

$$2A'(s) \frac{(A(s) - A^h(s))}{A(s)^3}$$



a term of  $\frac{1}{x}O(h)$  is contributed to (5.9) (just make successive replacements  $A^h \rightarrow A$  and study the difference by taking advantage of coercivity (C),  $(\Delta_2)$  and  $(\Delta'_2)$ ). So it is in fact sufficient to study:

$$\int_{[-\pi/h, \pi/h]} e^{isx} A'(s) \frac{(A(s) - A^h(s))}{A(s)^3} ds.$$

Now we do first Fubini for  $A'$  (via Remark 5.7), and get, beyond a factor of  $i$ :

$$\mu_0 \int_{[-\pi/h, \pi/h]} e^{isx} \frac{(A(s) - A^h(s))}{A(s)^3} ds + \int \lambda(dy) y e^{\gamma y} \int_{[-\pi/h, \pi/h]} e^{is(x+y)} \frac{(A(s) - A^h(s))}{A(s)^3} ds.$$

So what we would really like, is to show that:

$$\begin{aligned} & \int_{[-\pi/h, \pi/h]} e^{isz} \frac{(A^h(s) - A(s))}{A(s)^3} ds \\ &= (\mu - \mu^h) \int_{[-\pi/h, \pi/h]} e^{isz} \frac{e^{\beta h} - 1}{h} - \beta \frac{1}{A(s)^3} ds + c_0^h \int_{[-\pi/h, \pi/h]} e^{isz} \left( \frac{e^{\beta h} + e^{-\beta h} - 2}{2h^2} \right) \frac{1}{A(s)^3} ds \\ &+ \sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h} \lambda(du) \int_{[-\pi/h, \pi/h]} e^{isz} \frac{e^{\beta y} - \beta y \mathbb{1}_{[-v, 0)}(u) - (e^{\beta u} - \beta u \mathbb{1}_{[-v, 0)}(u))}{A(s)^3} ds \end{aligned} \quad (5.27)$$

is bounded by a (constant times  $h$ ), uniformly in  $z \in \mathbb{R}$  (it will then follow immediately that a term of  $\frac{1}{x}O(h)$  is being contributed to (5.9)).

- Now, the part corresponding to  $\mathbb{1}_{(-\infty, -V)} \cdot \lambda$ , namely:

$$\sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h \cap (-\infty, -V)} \lambda(du) \int_{[-\pi/h, \pi/h]} e^{isz} \frac{e^{\beta y} - e^{\beta u}}{A(s)^3} ds,$$

is clearly so.

- With respect to the term involving  $c_0^h = O(h)$ , make the decomposition:

$$\frac{e^{\beta h} + e^{-\beta h} - 2}{2h^2} = \left( \frac{e^{\beta h} + e^{-\beta h} - 2}{2h^2} - \frac{\beta^2}{2} \right) + \frac{\beta^2}{2}.$$

Then use coercivity (C) and (v) of Lemma 5.1 (respectively (1) of Lemma 5.6) for the first (respectively second) term.

- As regards:

$$(\mu - \mu^h) \int_{[-\pi/h, \pi/h]} e^{isz} \frac{\frac{1}{h}(e^{\beta h} - 1) - \beta}{A(s)^3} ds$$

write:

$$\frac{1}{h} (e^{\beta h} - 1) - \beta = \frac{1}{h} \left( \overbrace{e^{\beta h} - \frac{\beta^2 h^2}{2} - \beta h - 1}^{(I)} + \overbrace{\frac{\beta^2 h^2}{2}}^{(II)} \right). \quad (5.28)$$

By an expansion into a series, which converges absolutely and locally uniformly, and coercivity (C), it is clear that (I) has the desired property, whereas (1) of Lemma 5.6 may be applied to (II).

- Finally it will be sufficient to consider:

$$\sum_{y \in \mathbb{Z}_h^-} \int_{A_y^h \cap [-V, 0]} \lambda(du) \int_{[-\pi/h, \pi/h]} e^{isz} \frac{e^{\beta u} - \beta u - (e^{\beta y} - \beta y)}{A(s)^3} ds,$$

which we need bounded by a (constant times  $h$ ) uniformly in  $z \in \mathbb{R}$ . For this to work it is sufficient that the innermost integral produces ( $|u|$  times  $h$  times a constant). Moreover, it is enough to produce  $|y|$  (or, a fortiori,  $|y - u|$ ) in place of  $|u|$ . Now write:

$$\begin{aligned} e^{\beta u} - \beta u - (e^{\beta y} - \beta y) &= (e^{\beta y} - 1) \left( (e^{\beta(u-y)} - \beta(u-y) - 1) + \beta(u-y) \right) + \\ &\quad \left( e^{\beta(u-y)} - \frac{\beta^2(u-y)^2}{2} - \beta(u-y) - 1 \right) + \frac{\beta^2(u-y)^2}{2}. \end{aligned}$$

These terms can now be dealt with in part straightforwardly and in part by employing (3) and (1) of Lemma 5.6.

Third, with respect to (c) of (5.9), again an integration by parts is made outright, thus:

$$\frac{d}{ds} \left( e^{isx} \frac{1 - e^{ish}}{A^h(s)} \right) = ix e^{isx} \frac{1 - e^{ish}}{A^h(s)} + e^{isx} \frac{-ih e^{ish}}{A^h(s)} - e^{isx} \frac{(1 - e^{ish}) A^{h'}(s)}{A^h(s)^2}.$$

Now the left-hand side is handled using coercivity (C). On the right-hand side, we apply (1) of Lemma 5.6 to the second term. Finally, in the third term on the right-hand side we may replace  $A^{h'}(s)$  by  $A'(s)$ , followed by Fubini for  $A'$  and an application of (3) of Lemma 5.6. All in all, a term of order  $\frac{1}{x}O(h)$  thus emerges on the right-hand side of (5.9).

Part (ii) can be obtained by explicit computation, and is elementary.  $\square$

**Proposition 5.10** ( $\sigma^2 = 0$ ,  $\kappa(0) < \infty$  ( $Z^{(q)}$  convergence)). *Suppose  $\sigma^2 = 0$  and  $\kappa(0) < \infty$ . Let  $q > 0$ .*

- (i) *For any  $\gamma > \Phi(q)$ , there are  $\{A_0, h_0\} \subset (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $x \in \mathbb{Z}_h^{++}$ :*

$$\left| \Delta_Z^{(q)}(x, h) \right| \leq A_0 h e^{\gamma x}.$$

- (ii) *For the Lévy triplet  $(0, \delta_{-1}, 1)$  and the nested sequence  $(h_n := 1/2^n)_{n \geq 1} \downarrow 0$ , for each  $q > 0$  and any  $x \in \cup_{n \geq 1} \mathbb{Z}_{h_n}^{++} \cap (0, 1)$ :*

$$\lim_{n \rightarrow \infty} \frac{\Delta_Z^{(q)}(x, h_n)}{h_n} = \frac{1}{2} q (1 + q) x e^{x(1+q)}.$$

*Proof.* With respect to (i), we have as follows. First, (a) of (5.10) is of order  $O(h)$  by coercivity. Second, in (b), we employ the decomposition:

$$1 - \frac{\beta h}{1 - e^{-\beta h}} = \left( 1 - \frac{\beta h}{1 - e^{-\beta h}} + \frac{\beta h}{2} \right) - \frac{\beta h}{2}.$$

Then the first term may be estimated via (c) of Lemma 5.1 (for the denominator), a Taylor expansions into absolutely and locally uniformly convergent series (for the numerator) and coercivity

(C); while to the second term we apply (1) of Lemma 5.6. It follows that (b) of (5.10) is  $O(h)$ . Finally, when it comes to (c) of (5.10), we have (beyond a non-zero multiplicative constant):

$$\int_{[-\pi/h, \pi/h]} e^{isx} \frac{(A(s) - A^h(s))}{A(s)A^h(s)\beta} ds.$$

This can now be seen to be  $O(h)$  in the same manner as (5.27) was seen to be so (indeed, one can simply follow, word-for-word, the argument pursuant to (5.27), and recognize that the substitution of  $A(s)A^h(s)\beta$  in place of  $A(s)^3$  results in no material change).

Part (ii) follows by a direct computation.  $\square$

**5.9.  $\sigma^2 = 0$ , infinite variation paths.** Finally we consider the case when  $\sigma^2 = 0$  and  $\kappa(0) = \infty$ . We assume here that Assumption 1.1 is in effect. Note also that  $\delta^0 = 1$  and we work under scheme 2. We do not establish sharpness of the rates.

**Proposition 5.11** ( $\sigma^2 = 0$  &  $\kappa(0) = \infty$ ). *Assume  $\sigma^2 = 0$  and Assumption 1.1, let  $q \geq 0$ ,  $\gamma > \Phi(q)$ . Then there are  $\{A_0, h_0\} \subset (0, \infty)$  such that for all  $h \in (0, h_0)$  and then all  $x \in \mathbb{Z}_h^{++}$ :*

$$\begin{aligned} (i) \quad & \left| \Delta_W^{(q)}(x, h) \right| \leq A_0 \frac{h^{2-\epsilon}}{x} e^{\gamma x}. \\ (ii) \quad & \left| \Delta_Z^{(q)}(x, h) \right| \leq A_0 h^{2-\epsilon} e^{\gamma x}. \end{aligned}$$

*Proof.* With respect to (i), we have as follows. Note that (a) in (5.9) is  $\frac{1}{x}O(h^\epsilon)$  by an integration by parts argument and  $(\Delta'_1)$  from Subsection 5.4. We do the same for (b):

$$\begin{aligned} & \frac{d}{ds} \left( e^{isx} \left( \frac{1}{A(s)} - \frac{1}{A^h(s)} \right) \right) = ix e^{isx} \left( \frac{1}{A(s)} - \frac{1}{A^h(s)} \right) + \\ & + e^{isx} \left( \frac{A^2(s)(A^{h'}(s) - A'(s)) + A'(s)(A(s) - A^h(s))(A(s) + A^h(s))}{A^2(s)A^h(s)^2} \right). \end{aligned}$$

Upon integration, one gets on the left-hand side a contribution of  $\frac{1}{x}O(h^\epsilon)$  to (5.9), by coercivity (C). With regard to the rightmost quotient on the right-hand side, we obtain a contribution of order  $\frac{1}{x}O(h^{2-\epsilon})$ , as follows again by coercivity (C), Remark 5.10,  $(\Delta'_1)$  and the estimates  $(\Delta_2)$  and  $(\Delta'_2)$ . Remark that  $\epsilon > 2 - \epsilon$ .

With respect to (ii), we have as follows. (a) of (5.10) is of order  $O(h^\epsilon)$  by coercivity (C). Also, (b) is of order  $O(h)$ . Finally (c) is of order  $O(h^{2-\epsilon})$  immediately, with no need for an integration by parts.  $\square$

**5.10. A convergence result for the derivatives of  $W^{(q)}$  ( $\sigma^2 > 0$ ).**

**Proposition 5.12.** *Let  $q \geq 0$ ,  $\sigma^2 > 0$ . Note that  $W^{(q)}$  is then differentiable on  $(0, \infty)$  [22, Lemma 2.4]. Moreover, for any  $\gamma > \Phi(q)$ , there exist  $\{A_0, h_0\} \subset (0, \infty)$ , such that for all  $x \in \mathbb{Z}_h^{++} \setminus \{h\}$ :*

$$\left| W^{(q)'}(x) - \frac{W_h^{(q)}(x) - W_h^{(q)}(x - 2h)}{2h} \right| \leq A_0 \frac{e^{\gamma x}}{x} (h + \zeta(h/2) + \xi(h/2)h \log(1/h)).$$

*Remark 5.12.* The case  $\sigma^2 = 0$  appears much more difficult to analyze, since the balance between coercivity and the estimates of the differences of Laplace exponents (and their derivatives) worsens.

*Proof.* First, integration by parts, monotone convergence and the fact that  $W^{(q)}(0) = 0$  [23, p. 222, Lemma 8.6] yield (for  $\beta > \Phi(q)$ ):  $\widehat{W^{(q)'}}(\beta) = \beta \widehat{W^{(q)}}(\beta) = \beta/(\psi(\beta) - q)$ . Then analytic continuation, Laplace inversion and dominated convergence allow to conclude, for any  $x > 0$  that:

$$W^{(q)'}(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{\beta e^{\beta x}}{\psi(\beta) - q} ds.$$

On the other hand, it follows directly from Corollary 5.1, that for  $h \in (0, h_*)$  and then  $x \in \mathbb{Z}_h^{++}$ :

$$\frac{W_h^{(q)}(x) - W^{(q)}(x - 2h)}{2h} = \frac{1}{2\pi} \int_{-\pi/h}^{\pi/h} e^{\beta x} \left( \frac{e^{\beta h} - e^{-\beta h}}{2h} \right) \frac{ds}{\psi^h(\beta) - q}.$$

Now, it will be sufficient to estimate the following integrals:

$$\int_{(-\infty, \infty) \setminus [-\pi/h, \pi/h]} \beta e^{\beta x} A(s)^{-1} ds; \int_{-\pi/h}^{\pi/h} e^{\beta x} \left( \frac{e^{\beta h} - e^{-\beta h}}{2h} - \beta \right) A^h(s)^{-1} ds; \int_{-\pi/h}^{\pi/h} \beta e^{\beta x} \left( A(s)^{-1} - A^h(s)^{-1} \right) ds.$$

An integration by parts, coupled with coercivity (C) and the boundedness in linear growth of  $A'$  ( $\Delta'_1$ ), establishes the first of these two integrals as being of order  $\frac{1}{x}O(h)$ . The same emerges as being true of the second integral, this time using the boundedness in linear growth of  $A^{h'}$  instead, but also (iv) and (vi) of Lemma 5.1. Finally, with respect to the third integral, again one performs integration by parts, and then uses ( $\Delta_2$ ), coercivity (C), the decomposition (5.24) and ( $\Delta'_2$ ). The claim follows.  $\square$

## 6. NUMERICAL ILLUSTRATIONS AND CONCLUDING REMARKS

**6.1. Numerical examples.** We illustrate our algorithm for computing  $W$ , described in Eq. (1.1) of the Introduction, in two concrete examples, applying it to determine some relevant quantities arising in applied probability. The examples are chosen with two criteria in mind:

- (1) They are natural from the modeling perspective (computation of (Example 6.1) ruin parameters in the classical Cramér-Lundberg model with log-normal jumps and (Example 6.2) the Lévy-Kintchine triplet of the limit law of a CBI process).
- (2) They do not possess a closed form formula for the Laplace exponent of the spectrally negative Lévy processes. Such examples arise often in practice, making it difficult to apply the standard algorithms for scale functions based on Laplace inversion. Our algorithm is well-suited for such applications.

*Example 6.1.* A popular choice for the claim-size modeling in the Cramér-Lundberg surplus process is the log-normal distribution [3, Paragraph I.2.b, Example 2.8]. Fixing the values of the various parameters, consider the spectrally negative Lévy process  $X$  having  $\sigma^2 = 0$ ;  $\lambda(dy) = \mathbb{1}_{(-\infty, 0)}(y) \exp(-(\log(-y))^2/2)/(\sqrt{2\pi}(-y))dy$ ; and (with  $V = 0$ )  $\mu = 5$  (this satisfies

the security loading condition [24, Section 1.2]). Remark that the log-normal density has fat tails and is not completely monotone.

We complement the computation of  $W$  by applying it to the calculation of the density  $k$  of the deficit at ruin, on the event that  $X$  goes strictly above the level  $a = 5$ , before venturing strictly below 0, conditioned on  $X_0 = x = 2$ :

$$\mathbb{E}_x[-X_{\tau_0^-} \in dy, \tau_0^- < \tau_a^+] = k(y)dy$$

( $\tau_0^-$ , respectively  $\tau_a^+$ , being the first entrance time of  $X$  to  $(-\infty, 0)$ , respectively  $(a, \infty)$ ). Indeed,  $k(y)$  may be expressed as [24, Theorem 5.5]  $k(y) = \int_0^a f(z+y) \frac{W(x)W(a-z) - W(a)W(x-z)}{W(a)} dz$ , where  $f(y) := \exp(-(\log(y))^2/2)/(\sqrt{2\pi}y)$ ,  $y \in (0, +\infty)$ . We approximate the integral  $k$  by the discrete sum  $k_h$ , given for  $y \in (0, \infty)$  as follows:

$$k_h(y) := h \left[ f(y+a) \frac{W_h(x)W_h(0)}{2W_h(a)} + \sum_{k=1}^{a/h-1} f(kh+y)W_h(a-kh) \frac{W_h(x)}{W_h(a)} - \sum_{k=1}^{x/h-1} W_h(x-kh)f(kh+y) - \frac{W_h(0)f(x+y)}{2} \right].$$

Results are reported in Figure 1.

*Example 6.2.* We take  $\sigma^2 = 0$ ; the Lévy measure  $\lambda = \lambda_a + \lambda_c$  has atomic part  $\lambda_a = \frac{1}{2}(\delta_{-1} + \delta_{-2})$ , whilst the density of its absolutely continuous part  $\lambda_c(dy) = l(y)dy$  is given by:

$$l(y) = \frac{3}{2(-y)^{5/2}} \mathbb{1}_{[-1,0)}(y) + \frac{1}{2(-y-1)^{1/2}} \mathbb{1}_{[-2,-1)}(y) + \left( \frac{e^{\cos(y)}(3+y\sin(y))}{(-y)^4} + \frac{e}{(-y)^3} \right) \mathbb{1}_{(-\infty,-1)}(y), \quad y \in \mathbb{R};$$

and (with  $V = 1$ )  $\mu = 15$ . Remark the case is extreme: the Lévy measure has fat tails at 0 and  $-\infty$ , a discontinuity (indeed, a pole) in the Lévy density of its absolutely continuous part (which, in particular, is not completely monotone), and two atoms. Furthermore, there is no Gaussian component, and the sample paths of the process have infinite variation.

We compute  $W$  for the Lévy process having the above characteristic triplet, and complement this with the following application. Let furthermore  $X^F$  be an independent Lévy subordinator, given by  $X_t^F = t + Z_t$ ,  $t \in [0, \infty)$ , where  $Z$  is a compound Poisson process with Lévy measure  $m(dy) = e^{-y} \mathbb{1}_{(0, \infty)}(y)dy$ . Denote the dual  $-X$  of  $X$  by  $X^R$ . To the pair  $(X^F, X^R)$  there is associated, in a canonical way, (the law of) a (conservative) CBI process [20]. The latter process converges to a limit distribution  $L$ , as time goes to infinity, since  $\psi'(0+) > 0$  [21, Theorem 2.6(c)] and since further the log-moment of  $m$  away from zero is finite [21, Corollary 2.8]. Moreover, the limit  $L$  is infinitely divisible, and:

$$-\log \int_0^\infty e^{-ux} dL(x) = u\gamma - \int_{(0, \infty)} (e^{-ux} - 1) \frac{k(x)}{x} dx,$$

where  $\gamma = bW(0)$  vanishes, whilst:

$$k(x) = bW'(x+) + \int_{(0, \infty)} [W(x) - W(x-\xi)]m(d\xi),$$




FIGURE 1. The scale function  $W$  and the density  $k$  of the deficit at ruin on the event  $\{\tau_0^- < \tau_5^+\}$ , for the log-normal Cramér-Lundberg process. The relative errors are consistent with the linear order of convergence predicted by Theorem 1.1. See Example 6.1 for details.

where  $b = 1$  and  $m$  are the drift, respectively the Lévy measure, of  $X^F$  [21, Theorem 3.1]. We compute  $k$  via approximating, for  $x \in \mathbb{Z}_h^{++}$ ,  $k(x)$  by  $k_h(x)$ :

$$k_h(x) := b \frac{W_h(x) - W_h(x-h)}{h} + W_h(x-h)m[h/2, \infty) - \sum_{k=1}^{x/h-1} W_h(x-kh-h)m[kh-h/2, kh+h/2).$$

Results are reported in Figure 2.

Let us also mention that we have tested our algorithm on very simple processes with completely monotone Lévy densities [2] (Brownian motion with drift; positive drift minus a compound Poisson subordinator with exponential jumps; spectrally negative stable Lévy process — see Appendix C), and the results were in nice agreement with the explicit formulae which are available for the scale functions in the latter cases.




FIGURE 2. The scale function  $W$  for the spectrally negative Lévy process  $X$ , as described in Example 6.2; followed by the  $k$ -function  $k$  of the weak limit (as time goes to infinity) of the CBI process, whose spectrally positive component is the dual  $X^R = -X$  of  $X$ , whilst the Lévy subordinator part is the sum of a unit drift and a compound Poisson process of unit intensity and mean one exponential jumps. The relative errors are consistent with the  $O(\sqrt{h})$  order of convergence predicted by Theorem 1.1. See Example 6.2 for details.

- 6.2. Concluding remarks.** (1) Computational cost. To compute  $W_h^{(q)}(x)$  or  $Z_h^{(q)}(x)$  for some  $x \in \mathbb{Z}_h$  one effects recursions (3.3) and (3.4) (as applied to  $Y = X^h/h$ ,  $h \in (0, h_*)$ ), at a cost of  $O((x/h)^2)$  operations (assuming given the parameters of  $X^h$ ).
- (2) Quantity  $Z_h^{(q)}(x)$  may be obtained from the values of  $W_h^{(q)}$  on  $[0, x] \cap \mathbb{Z}_h$  at a cost of order  $O(x/h)$  operations by a nonnegative summation (see Proposition 3.1).
- (3) The computation of the functions  $W^{(q)}$ ,  $q \geq 0$ , can be reduced, under an exponential change of measure, to the computation of  $W$  [23, p. 222, Lemma 8.4] for a process having  $\Phi(0) = 0$  [34].

Under such an exponential tilting  $W_h(x)$  will have a temperate growth [35, Proposition 4.8(ii)], since  $\Phi^h(0) \rightarrow \Phi(0)$ .

Finally, in comparison to the Laplace inversion methods discussed in [22, Section 5.6], we note: (1) Regarding *only* the efficiency of our algorithm (i.e. how costly it is, to achieve a given precision): Firstly, that Filon’s method (with Fast Fourier Transform) appears to outperform ours when an explicit formula for the Laplace exponent  $\psi$  is known. Secondly, that our method is largely insensitive to the degree of smoothness of the target scale function – and can match or outperform Euler’s, the Gaver-Stehfest and the fixed Talbot’s method in regimes when the scale function is less smooth, even as  $\psi$  remains readily available (such, at least, was the case for Sets 3 and 4 of [22, pp. 177-178] — see Appendix C). Thirdly, that when  $\psi$  is not given in terms of elementary/special function, any Laplace inversion algorithm, by its very nature, must resort to further numerical evaluations of  $\psi$  (at complex values of its argument), which hinders its efficiency and makes it hard to control the error. Indeed, such evaluations of  $\psi$  appear disadvantageous, as compared to the more innocuous operations required to compute the coefficients present in our recursion.

(2) In our method there is only one spatial discretization parameter  $h$  to vary. On the other hand, Filon’s method (which, when coupled with Fast Fourier Transform, appears the most efficient of the Laplace inversion techniques), has additionally a cutoff parameter in the (complex) Bromwich integral.

## REFERENCES

1. J. Abate and W. Whitt. A Unified Framework for Numerically Inverting Laplace Transforms. *INFORMS Journal on Computing*, 18(4):408–421, 2006.
2. H. Albrecher, F. Avram, and D. Kortschak. On the efficient evaluation of ruin probabilities for completely monotone claim distributions. *Journal of Computational and Applied Mathematics*, 233(10):2724 – 2736, 2010.
3. S. Asmussen and H. Albrecher. *Ruin Probabilities*. Advanced series on statistical science and applied probability. World Scientific, 2010.
4. F. Avram, A. E. Kyprianou, and M. R. Pistorius. Exit Problems for Spectrally Negative Lévy Processes and Applications to (Canadized) Russian Options. *Annals of Applied Probability*, 14(1):215–238, 2004.
5. J. Bertoin. *Lévy Processes*. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1996.
6. R. N. Bhattacharya and E. C. Waymire. *A Basic Course in Probability Theory*. Universitext - Springer-Verlag. Springer, 2007.
7. E. Biffis and A. E. Kyprianou. A note on scale functions and the time value of ruin for Lévy insurance risk processes. *Insurance: Mathematics and Economics*, 46:85–91, 2010.
8. N. H. Bingham. Continuous branching processes and spectral positivity. *Stochastic Processes and their Applications*, 4(3):217 – 242, 1976.
9. A. N. Borodin and P. Salminen. *Handbook of Brownian motion—facts and formulae*. Probability and its Applications. Birkhäuser Verlag, Basel, second edition, 2002.
10. H. Brunner. *Collocation Methods for Volterra Integral and Related Functional Differential Equations*. Cambridge Monographs on Applied and Computational Mathematics. Cambridge University Press, 2004.
11. R. M. R. Cardoso and H. R. Waters. Recursive calculation of finite time ruin probabilities under interest force. *Insurance: Mathematics and Economics*, 33(3):659 – 676, 2003.



12. T. Chan, A. E. Kyprianou, and M. Savov. Smoothness of scale functions for spectrally negative Lévy processes. *Probability Theory and Related Fields*, 150(3-4):691–708, 2011.
13. A. M. Cohen. *Numerical methods for Laplace transform inversion*, volume 5 of *Numerical Methods and Algorithms*. Springer, New York, 2007.
14. B. Davies. *Integral Transforms and Their Applications*. Texts in Applied Mathematics. Springer, Third edition, 2002.
15. D. C. M. Dickson and J. R. Gray. Approximations to ruin probability in the presence of an upper absorbing barrier. *Scandinavian Actuarial Journal*, 1984(2):105–115, 1984.
16. D. C. M. Dickson and H. R. Waters. Recursive calculation of survival probabilities. *ASTIN Bulletin*, 21(2):199–221, 1991.
17. J. Cl. Evard and F. Jafari. A Complex Rolle’s Theorem. *The American Mathematical Monthly*, 99(9):858–861, 1992.
18. F. Hubalek and A. E. Kyprianou. Old and New Examples of Scale Functions for Spectrally Negative Lévy Processes. In R. Dalang, M. Dozzi, and F. Russo, editors, *Seminar on Stochastic Analysis, Random Fields and Applications VI*, Progress in Probability, pages 119–145. Springer, 2011.
19. E. I. Jury. *Theory and application of the z-transform method*. Robert E. Krieger Publishing Co., Huntington, New York, 1964.
20. K. Kawazu and S. Watanabe. Branching Processes with Immigration and Related Limit Theorems. *Theory of Probability and its Applications*, 16(1):36–54, 1971.
21. M. Keller-Ressel and A. Mijatovic. On the limit distributions of continuous-state branching processes with immigration. *Stochastic Processes and Their Applications*, 122:2329–2345, 2012.
22. A. Kuznetsov, A. E. Kyprianou, and V. Rivero. The theory of scale functions for spectrally negative Lévy processes. In *Lévy Matters II*, Lecture Notes in Mathematics, pages 97–186. Springer Berlin Heidelberg, 2013.
23. A. E. Kyprianou. *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer-Verlag, Berlin Heidelberg, 2006.
24. A. E. Kyprianou. *Gerber-Shiu Risk Theory*. EEA Series. Springer, 2013.
25. A. Lambert. Species abundance distributions in neutral models with immigration or mutation and general lifetimes. *Journal of Mathematical Biology*, 63(1):57–72, 2011.
26. P. Linz. *Analytical and Numerical Methods for Volterra Equations*. Studies in Applied Mathematics. Society for Industrial and Applied Mathematics, 1985.
27. R. L. Loeffen. On optimality of the barrier strategy in de Finetti’s dividend problem for spectrally negative Lévy processes. *Annals of Applied Probability*, 18(5):1669–1680, 2008.
28. A. Mijatović and M. R. Pistorius. On the drawdown of completely asymmetric Lévy processes. *Stochastic Processes and their Applications*, 122(11):3812–3836, 2012.
29. A. Mijatović, M. Vidmar, and S. Jacka. Matlab code for the scale function algorithm in Equation (1.1) above, 2013. Available at [http://www.ma.ic.ac.uk/~amijatov/Abstracts/ScaleFun\\_Levy.html](http://www.ma.ic.ac.uk/~amijatov/Abstracts/ScaleFun_Levy.html).
30. A. Mijatović, M. Vidmar, and S. Jacka. Markov chain approximations for transition densities of Lévy processes. *Electronic Journal of Probability*, 19(7):1–37, 2014.
31. H. H. Panjer and S. Wang. On the stability of recursive formulas. *ASTIN Bulletin*, 23(2):227–258, 1993.
32. D. Revuz and M. Yor. *Continuous Martingales and Brownian Motion*. Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin Heidelberg, 1999.
33. K. I. Sato. *Lévy Processes and Infinitely Divisible Distributions*. Cambridge studies in advanced mathematics. Cambridge University Press, Cambridge, 1999.

34. B. A. Surya. Evaluating Scale Functions of Spectrally Negative Lévy Processes. *Journal of Applied Probability*, 45(1):135–149, 2008.
35. M. Vidmar. Fluctuation theory for upwards skip-free Lévy chains. arXiv:1309.5328 [math.PR], 2013.
36. F. De Vylder and M. J. Goovaerts. Recursive calculation of finite-time ruin probabilities. *Insurance: Mathematics and Economics*, 7(1):1–7, 1988.

## APPENDIX A. PROOFS OF LEMMAS FROM PARAGRAPH 5.2.1

*Proof.* (Of Lemma 5.1.) Assertions (i)-(vi) obtain at once by expansion into Taylor series which converge absolutely and locally uniformly. (a) follows from the equality  $1 - \cos(u) = 2 \sin^2(u/2)$  and the concavity of  $\sin|_{[0, \xi/2]}$ . Then (b) is got from (letting  $u = \gamma_0 + is_0$ ,  $\{\gamma_0, s_0\} \subset \mathbb{R}$ ):

$$\left| \frac{1}{u^2} (\cosh u - 1) \right|^2 = \frac{[(\cosh \gamma_0 - 1) + (1 - \cos s_0)]^2}{(\gamma_0^2 + s_0^2)^2} \geq \frac{(\gamma_0^2/2 + 1 - \cos s_0)^2}{(\gamma_0^2 + s_0^2)^2},$$

since  $\cosh|_{\mathbb{R}}$ , upon expansion into a power series, is a nondecreasing limit of its partial sums. Finally, (c) obtains from (again  $u = \gamma_0 + is_0$ ,  $\{\gamma_0, s_0\} \subset \mathbb{R}$ ):

$$\left| \frac{1}{u} (e^u - 1) \right|^2 = \frac{(e^{\gamma_0} - 1)^2 + 2e^{\gamma_0}(1 - \cos s_0)}{\gamma_0^2 + s_0^2},$$

noting that  $|e^{\gamma_0} - 1| \geq B|\gamma_0|$  for all  $\gamma_0 \geq L$ , for some  $B > 0$ , whereas  $\exp|_{[L, \infty)}$  is itself bounded away from zero.  $\square$

*Proof.* (Of Lemma 5.2.) Expressing  $f_{\gamma_0}(u) := \frac{1 - e^{-\gamma_0 - is_0} - (\gamma_0 + is_0)}{(1 - e^{-\gamma_0 - is_0})(\gamma_0 + is_0)}$ , use (c) of Lemma 5.1 in the denominator, and expansion into Taylor series which converge absolutely and locally uniformly in the numerator.  $\square$

*Proof.* (Of Lemma 5.3.) The first inequality follows from (writing  $z = \gamma_0 + is_0$ ,  $\{\gamma_0, s_0\} \subset \mathbb{R}$ ):

$$|e^z - 1|^2 = (e^{\gamma_0} - 1)^2 + 2e^{\gamma_0}(1 - \cos s_0).$$

Then, since  $\gamma_0 \leq 0$ ,  $e^{\gamma_0} \leq 1$  and  $1 - e^{\gamma_0} \leq -\gamma_0$  (by comparing derivatives). Finally, use  $1 - \cos s_0 \leq s_0^2/2$ .

The second inequality obtains from the relation (for  $\{\gamma_0, s_0\} \subset \mathbb{R}$ ):

$$e^{\gamma_0 + is_0} - (\gamma_0 + is_0) - 1 = e^{\gamma_0}(\cos(s_0) - 1) + (e^{\gamma_0} - \gamma_0 - 1) + i(e^{\gamma_0} - 1)\sin(s_0) + i(\sin(s_0) - s_0),$$

noting in addition, that  $e^{\gamma_0} - \gamma_0 - 1 \leq \gamma_0^2/2$  for  $\gamma_0 \leq 0$  (compare derivatives),  $|\sin(s_0)| \leq |s_0|$  and finally  $\text{sgn}(s_0)(s_0 - \sin(s_0)) \leq (|s_0|^3/6) \wedge (2|s_0|) \leq 2s_0^2$ .  $\square$

*Proof.* (Of Lemma 5.4.) Apply the complex Mean Value Theorem [17, p. 859, Theorem 2.2].  $\square$

## APPENDIX B. PROOFS FOR PARAGRAPH 5.2.2

*Proof.* (Of Proposition 5.2.) It is Fubini's Theorem that:

$$I := \int_1^\infty \frac{ds}{s^2} \int \nu(dx)(1 - \cos(sx)) = \int \nu(dx) \int_1^\infty \frac{ds}{s^2}(1 - \cos(sx)).$$

Next, for each  $x \in \mathbb{R}$ , do integration by parts for the integral  $\int_1^\infty \frac{ds}{s^2}(1 - \cos(sx))$ ; first,

$$\frac{d}{ds} \left( \frac{1 - \cos(sx)}{s} \right) = -\frac{1 - \cos(sx)}{s^2} + x \frac{\sin(sx)}{s};$$

second, integrate from 1 to  $N$  against  $ds$ ; third, let  $N \rightarrow \infty$  and use the Monotone Convergence Theorem. We obtain:

$$I = \int \nu(dx) \left( x \int_1^\infty \frac{\sin(sx)}{s} ds + (1 - \cos(x)) \right).$$

The inner integral is, of course, in the improper Riemann sense; now change variables in the latter to get:

$$I = \int \nu(dx) \left( |x| \int_{|x|}^\infty \frac{\sin(u)}{u} du + (1 - \cos(x)) \right).$$

Since the sine integral is bounded,  $1 - \cos(x) \leq x^2/2$  and  $\nu$  is compactly supported, it follows that  $I < \infty$  whenever  $\int |x|\nu(dx) < \infty$ . Conversely, if  $\int |x|\nu(dx) = \infty$ , since  $\int_{|x|}^\infty \frac{\sin(u)}{u} du \rightarrow \int_0^\infty \frac{\sin(u)}{u} du \in (0, \infty)$  as  $x \rightarrow 0$ , we deduce  $I = \infty$  by the local finiteness of  $\nu$  in  $\mathbb{R} \setminus \{0\}$ .  $\square$

*Proof.* (Of Lemma 5.5.) Fubini's Theorem.  $\square$

*Proof.* (Of Proposition 5.3.) Apply Lemma 5.5.  $\square$

*Proof.* (Of Proposition 5.4.) The first assertion follows at once from (iii) of Lemma 5.5 (applied to the measure  $\nu$ ). The second follows from the first and (ii) of Lemma 5.5 applied to the measure  $\lambda$ . Now we consider  $\int (e^{isy} - 1)\lambda(dy)$ . It is assumed without loss of generality that  $\nu$  is supported by the positive half-line and  $s > 0$ . Furthermore, there is an  $A \in (0, \infty)$ , with  $\hat{\lambda}(u) := \lambda(u, 1) \leq A/u^\alpha$  for all  $u \in (0, 1)$ .

Consider the imaginary part of  $\int (e^{isy} - 1)\lambda(dy)$  first. Fubini and dominated convergence yield:

$$\int \lambda(dy) \sin(sy) = s \int \lambda(dy) \int_0^y \cos(su) du = s \int_0^1 du \cos(su) \hat{\lambda}(u) = s \sum_{k=0}^\infty \int_{(0,1) \cap [k\pi/2, k\pi + \pi/2)} du \cos(su) \hat{\lambda}(u) =: I(s).$$

We thus obtain an alternating series, whence:  $|I(s)| \leq s \int_0^{3\pi/(2s)} du \hat{\lambda}(u)$ . This is crucial. Indeed, taking into account that  $\hat{\lambda}(u) \leq A/u^\alpha$  throughout the integration region, we now obtain immediately  $\sup_{s>0} |I(s)|/s^\alpha < \infty$ .

The real part is treated in a similar vein. First, Fubini yields:

$$\int \lambda(dy)(1 - \cos(sy)) = s \int \lambda(dy) \int_0^y \sin(su) du = s \int_0^1 du \sin(su) \hat{\lambda}(u)$$

and the remaining steps are very similar and hence omitted.  $\square$

## APPENDIX C. MORE NUMERICAL EXAMPLES




FIGURE 3. Left: Brownian motion with drift ( $\sigma^2 = 1$ , (with  $V = 0$ )  $\mu = 1$ ,  $\lambda = 0$ ). Right: a stable process ( $\sigma^2 = 0$ , (with  $V = 1$ )  $\mu = 1/(\beta - 1)$ ,  $\lambda(dy) = \frac{1}{(-y)^{1+\beta}} \mathbb{1}_{(-\infty, 0)}(y) dy$ ,  $\beta = 3/2$ ). Note the quadratic, respectively ‘square-root’, order of convergence.

*Example C.1.* This example is taken from [22, pp. 177-178; Eq. (137), Table 4 (Sets 3 and 4)]; with two *caveats*: (i) parameter  $c_4$  is set equal to  $e$  in Set 3, since there appears to be a typo in the last term of the expression for the Laplace exponent [22, p. 178, Equation (138)] and presumably the form of the Laplace exponent as given (*sic*), is the one which was used in the actual computations of [22]; (ii) our parameter  $\mu$  (with  $V = 1$ ) takes on such values as to ensure equality of Laplace exponents (i.e. laws of the processes) between us and [22] for each of the two sets. We perform the computation for the function  $W^{(1/2)}$  on the decreasing sequence  $(h_n = 1/(20 \cdot n))_{n \in \{5, 20, 80, 320\}}$ .  $W_{h_{2000}}^{(1/2)}(\cdot - \delta_0 h_{2000})$  is taken as a benchmark, and this gives  $\max \text{rel}_n :=$

$\max_{i \in [100]} \frac{|W_{h_n}^{(1/2)}(x_i - \delta_0 h_n) - W_{h_{2000}}^{(1/2)}(x_i - \delta_0 h_{2000})|}{W_{h_{2000}}^{(1/2)}(x_i - \delta_0 h_{2000})}$ , where  $x_i = i/20$ ,  $i \in [100]$ . Note the linear order of convergence.

$n$	5	20	80	320
max rel $_n$ (Set 3)	0.0011	$3.4086 \cdot 10^{-5}$	$6.6513 \cdot 10^{-6}$	$1.2391 \cdot 10^{-6}$
max rel $_n$ (Set 4)	0.0121	0.0031	$7.5203 \cdot 10^{-4}$	$1.6512 \cdot 10^{-4}$




FIGURE 4. Exponential jumps model ( $\lambda(dy) = a\rho e^{\rho y} \mathbb{1}_{(-\infty, 0)}(y)dy$ ,  $a = \rho = 1$ ,  $\sigma^2 = 0$  and (with  $V = 0$ )  $\mu \in \{2, 1\}$ ). Note the linear order of convergence.

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, UK

*E-mail address:* a.mijatovic@imperial.ac.uk

DEPARTMENT OF STATISTICS, UNIVERSITY OF WARWICK, UK

*E-mail address:* m.vidmar@warwick.ac.uk

DEPARTMENT OF STATISTICS, UNIVERSITY OF WARWICK, UK

*E-mail address:* s.d.jacka@warwick.ac.uk