



Quotients of \mathbb{R}^8 & Reduced Holonomy

Simon Salamon

In honour of Dmitri Alekseevsky

9 September 2020

...contained in this talk include

- Riemannian holonomy groups
- Einstein metrics
- Quaternionic geometry
- Twistor spaces

- [ABS] B. Acharya, R. Bryant, S. Salamon: in DGA 2020
- [KL] S. Karigiannis, J. Lotay: arXiv:2002.06444
- [FHN] L. Foscolo, M. Haskins, J. Nordstrom: arXiv:1805.02612
- [K] K. Kawai: in Comm. Anal. Geom. 2018
- [B] O. Bogoyavlenskaya: in Sibirsk. Mat. Zh. 2013
- [AS] V. Apostolov, S. Salamon: in Comm. Math. Phys. 2004
- [AyW] M. Atiyah, E. Witten: in Adv. Theor. Math. Phys. 2002
- [AW] B. Acharya, E. Witten: arXiv:hep-th/0109152

If N^6 is nearly Kähler then the cone $\mathbb{R}^+ \times N$ has a Ricci-flat metric with holonomy in G_2 .

We shall take $N = \mathbb{C}\mathbb{P}^3$ with its NK structure and non-integrable almost complex structure J_2 arising from the fibration $\mathbb{C}\mathbb{P}^3 \rightarrow S^4$. Set $\mathcal{C} = \mathbb{R}^+ \times \mathbb{C}\mathbb{P}^3$. We shall

- construct the resulting G_2 metric h on \mathcal{C} starting from $\mathbb{C}^4 = \mathbb{R}^8$,
- investigate the geometry arising from an action of $SO(2)$ rotating S^4 ,
- explain that the quotient $\mathcal{C}/SO(2)$ is essentially \mathbb{R}^6 ,
- describe the induced $SU(3)$ structure (σ, g, \mathbb{J}) on \mathbb{R}^6 in the spirit of [AS].

The metric h can be smoothed into a complete asymptotically conical (AC) metric on the total space of $\Lambda_-^2 T^*S^4$ [BS]. There are analogous AC metrics formed from the NK spaces $SU(3)/T^2$ and $S^3 \times S^3$, though the last one is the most amenable for study (next slides).

The AC metric on the spin bundle over S^3 with isometry group $SU(2)^3$ represents a bifurcation in a one-parameter family of G_2 metrics with a cohomogeneous-one action by $SU(2)^2 \times U(1)$, in two different ways giving a G_2 flop [FHN].


The AC metric is a limit of asymptotically locally conical (ALC) metrics, each of which has a circle of fixed radius r at infinity. These ALC metrics first appeared in the physics literature [BGGG, CGLP, ...] with the names \mathbb{B}_7 and \mathbb{D}_7 and the existence of one was proved by [B]. In the collapsed limit as $r \rightarrow 0$, one obtains an AC Calabi-Yau space.

Circle bundles over singular Calabi-Yau spaces can be used to construct G_2 metrics [FHN']. There is an infinite family of complete AC G_2 metrics on circle bundles $M_{m,n} \rightarrow K_{\mathbb{C}P^1 \times \mathbb{C}P^1}$ that are asymptotic to cones over finite quotients of $S^3 \times S^3$.

G_2 metrics of cohomogeneity one

5/20

...based on one-parameter families of half-flat $SU(3)$ structures on $S^3 \times S^3$ invariant by $SU(2)^2 \times U(1)$:



By analogy, the $\mathrm{SO}(5)$ -invariant G_2 metric on $\Lambda_-^2 T^*S^4$ arises as a collapsed limit of metrics with holonomy $\mathrm{Spin}(7)$ on the spin bundle over S^4 .

Dirac monopole: $\mathrm{U}(1)$ acts on the left on \mathbb{H} with quotient $\mathbb{R}^4/\mathrm{U}(1) \cong \Lambda_-^2$

Let's return to $\mathcal{C} = \mathbb{R}^+ \times \mathbb{C}\mathbb{P}^3$ with its conical G_2 metric h . There is no obvious way to associate ALC metrics to this set-up because of the absence of free circle and group actions. Nonetheless, there is an action of $\mathrm{SO}(2)$ on

$$S^4 \subset \mathbb{R}^2 \oplus \mathbb{R}^3$$

that lifts to $\mathbb{C}\mathbb{P}^3$ and fixes two 2-spheres. Then $\mathcal{C}/\mathrm{SO}(2)$ has singular locus $\mathbb{R}^3 \cup \mathbb{R}^3$ (minus the origin). M-theory formulated on \mathcal{C} is dual to Type IIA superstring theory on \mathbb{R}^6 , and fixed points of $\mathrm{SO}(2)$ on \mathcal{C} are identified with D-branes of the quotient [AyW].

Start with $\mathbb{H}^2 = \mathbb{C}^4 = \mathbb{R}^8$. Its QK structure corresponds to $\mathrm{Sp}(2)_\ell \times \mathrm{Sp}(1)_r$ modulo \mathbb{Z}_2 . Consider the subgroups

$$\mathrm{U}(1)_\ell \times \mathrm{SU}(2) \subset \mathrm{Sp}(2)_\ell, \quad \mathrm{Sp}(1)_r \supset \mathrm{U}(1)_r.$$

The 2-torus $\mathrm{U}(1)_\ell \times \mathrm{U}(1)_r$ acts on $\mathbb{H}^2 = \mathbb{C}^4$ as

$$\begin{aligned} (q_0, q_1) &\mapsto e^{i\theta}(q_0, q_2)e^{i\phi} \\ (\mathfrak{z}_0, \mathfrak{z}_1, \mathfrak{z}_2, \mathfrak{z}_3) &\mapsto (e^{i(\theta+\phi)}\mathfrak{z}_0, e^{i(\theta-\phi)}\mathfrak{z}_1, e^{i(\theta+\phi)}\mathfrak{z}_2, e^{i(\theta-\phi)}\mathfrak{z}_3). \end{aligned}$$

It splits $\mathbb{R}^8 = \mathbb{R}_{0145}^4 \oplus \mathbb{R}_{2367}^4$ with a ‘transposed’ hyperkähler structure associated to

$$\Lambda_-^2(\mathbb{R}_{0145}^4) \oplus \Lambda_-^2(\mathbb{R}_{2367}^4) \cong \mathfrak{su}(2) \oplus \mathfrak{su}(2) \cong \mathbb{R}^6.$$

This space is T^2 invariant, and is the target of an associated moment map.

Define $\mathbb{C}\mathbb{P}^3 = \mathcal{S}^7/U(1)_r$ and set $\mathcal{C} = \mathbb{R}^+ \times \mathbb{C}\mathbb{P}^3$. Then the moment map induces

$$\begin{aligned} \mu: \quad \mathcal{C} &\longrightarrow \mathbb{R}^6 \\ [\mathfrak{z}_0, \mathfrak{z}_1, \mathfrak{z}_2, \mathfrak{z}_3] &\longmapsto (\mathbf{u}, \mathbf{v}), \end{aligned}$$

whose fibres are orbits of $U(1)_\ell/\mathbb{Z}_2 = \text{SO}(2)$ à la Gibbons-Hawking, and

$$\begin{cases} u_1 = |\mathfrak{z}_0|^2 - |\mathfrak{z}_2|^2 \\ u_2 = 2\text{Re}(\mathfrak{z}_0\bar{\mathfrak{z}}_2) \\ u_3 = -2\text{Im}(\mathfrak{z}_0\bar{\mathfrak{z}}_2), \end{cases} \quad \begin{cases} v_1 = |\mathfrak{z}_1|^2 - |\mathfrak{z}_3|^2 \\ v_2 = 2\text{Re}(\mathfrak{z}_1\bar{\mathfrak{z}}_3) \\ v_3 = 2\text{Im}(\mathfrak{z}_1\bar{\mathfrak{z}}_3). \end{cases}$$

$$R = \sum_{i=0}^3 |\mathfrak{z}_i|^2 \text{ equals } u + v, \text{ where } u = |\mathbf{u}| \text{ and } v = |\mathbf{v}|.$$

The action of T^2 on \mathbb{R}^8 commutes with $\text{SU}(2)$ that acts as $\text{SO}(3)$ diagonally on \mathbb{R}^6 .

Let $\{\alpha_1, \alpha_2, \alpha_3\}$ be a HK triple of 2-forms on \mathbb{R}^8 trivializing the action of $\mathrm{Sp}(1)_r$, and suppose that α_1^\sharp generates $\mathrm{U}(1)_r$.

Proposition. The G_2 3-form φ on \mathcal{C} equals $-d(R\tau)$, where

$$\tau = dR \wedge \alpha_1 - \alpha_2 \wedge \alpha_3.$$

To smooth the vertex ($r = 0$) of the cone, replace the coefficient R of τ by $(R^4 + 1)^{1/4}$. The resulting complete AC metric is

$$(R^4 + 1)^{-1/2} g_{\mathrm{ver}} + (R^4 + 1)^{1/2} g_{\mathrm{hor}}.$$

It has convergence rate -4 (since R is Euclidean radius squared) and is rigid as an AC metric [KL'].

We are now considering the quotient of the G_2 manifold $\mathcal{C} = \mathbb{R}^+ \times \mathbb{C}\mathbb{P}^3$ by $\text{SO}(2)$, which is $\mathbb{R}^6 \setminus \mathbf{0}$. The $\text{SO}(3)$ orbit of a bivector $(\mathbf{u}, \mathbf{v}) \in \mathbb{R}^6$ has dimension 3 unless $\mathbf{u} \wedge \mathbf{v} = 0$.

Definition. Set $\mathcal{F}_\pm = \{(\mathbf{u}, \mathbf{v}) \in \mathbb{R}^6 : v\mathbf{u} = \pm u\mathbf{v}\}$.

The equation $uv = 0$ defines the singular locus $\mathbb{R}^3 \cup \mathbb{R}^3$ of \mathbb{R}^6 where the circle fibres of μ collapse. If $uv \neq 0$ then $(\mathbf{u}, \mathbf{v}) \in \mathcal{F}_+$ (resp. \mathcal{F}_-) iff \mathbf{u}, \mathbf{v} are aligned (resp. anti-aligned).

We can interpret these sets in terms of the fibration $\mathbb{C}\mathbb{P}^3 \rightarrow \mathcal{S}^4$ (next slide):

Theorem. $\mu^{-1}(\mathcal{F}_\pm)/\mathbb{R}^* = Q_\pm \subset \mathbb{C}\mathbb{P}^3$ where

$Q_+ = \{[z_0, z_1, z_2, z_3] : z_0\bar{z}_3 - z_1\bar{z}_2 = 0\}$ consists of points where $U(1)_\ell$ acts vertically,

$Q_- = \{[z_0, z_1, z_2, z_3] : z_0z_1 + z_2z_3 = 0\}$ consists of points where $U(1)_\ell$ acts horizontally.

Rather than using the Hopf map $\mathbb{C}P^3 \rightarrow \mathbb{H}P^1$, one can pass directly to the 4-sphere:

$$\begin{array}{ccc}
 \boxed{\begin{array}{c} \mathbb{C}^4 \\ \downarrow \\ \Lambda_0^2(\mathbb{C}^4) \end{array}} & \cong \mathbb{R}^5 & \supset \mathcal{S}^4 \\
 & & \pi \downarrow \\
 & & \mathbb{C}P^3 \times \mathbb{R}^+ = \mathcal{C}
 \end{array}$$

The action of $U(1)_\ell$ on \mathbb{R}^8 covers a rotation of \mathcal{S}^4 :

$$\begin{array}{ccccccc}
 U(1)_\ell & \subset & U(2) & \subset & Sp(2)_\ell & & \\
 & & & & \downarrow & & \\
 SO(2) & \subset & SO(2) \times SO(3) & \subset & SO(5) & &
 \end{array}$$

Let $\mathcal{S}^1 = \mathcal{S}^4 \cap \mathbb{R}^2$ denote the fixed point set for the action of $SO(3)$ ($s = 0$ next),

Let $\mathcal{S}^2 = \mathcal{S}^4 \cap \mathbb{R}^3$ denote the fixed point set for the action of $SO(2)$ ($s = 1$ next).

The non-holomorphic quadric Q_+ is simply $\pi^{-1}(\mathbb{S}^2) \cong \mathcal{S}^2 \times \mathcal{S}^2$.

By contrast, Q_- contains $\pi^{-1}(\mathbb{S}^1)$, away from which it is a double covering of

$$\mathcal{S}^4 \setminus \mathbb{S}^1 \cong \mathbb{H} \setminus \mathbb{R} \cong \mathcal{S}^2 \times \mathcal{H}^2.$$

It encodes the conformally Kähler metric [Pont,SV] and the orthogonal complex structure on $\mathbb{H} \setminus \mathbb{R}$ that can be used to define quaternion power series [GSS].

If X is the Killing field generated by $\text{SO}(2)$, then

$$X^\flat = (1 - s^2)dt,$$

where $t: \mathcal{S}^4 \setminus \mathbb{S}^2 \rightarrow [0, 2\pi)$ and $s: \mathcal{S}^4 \rightarrow [0, 1]$.

The 'dual pair' $SO(2) \times SO(3)$ (arising from $U(2) \subset Sp(2)_\ell$) acts on $\mathbb{C}P^3$ and \mathcal{C} . We have already parametrized the orbits of $SO(2)$, and will deal with those of $SO(3)$ shortly.

One could instead focus on

$$U(1) \times Sp(1) \subset Sp(2)_\ell$$

that acts as $U(2)$ on \mathbb{R}^4 fixing two poles of S^4 . Or work with arbitrary weights for the action of a circle subgroup of $U(2)$ on \mathbb{C}^2 .

Backtracking, we could replace $U(1)_r$ by $U(1)$ with weights (p, q) on \mathbb{H}^2 giving rise to weighted projective space $\mathbb{W}CP^3_{p,p,q,q}$ with a circle action again fixing two projective lines. This space is conjectured to carry a NK metric [AW].

Let G be a compact Lie group, for instance $SO(5)$.

Key fact. Each conjugacy class of subalgebras $\mathfrak{su}(2) \subset \mathfrak{g}$ gives rise to a complex nilpotent orbit $\mathcal{N} \subset \mathfrak{g}_{\mathbb{C}}$ with a HK metric [Kr], and a (typically incomplete) QK metric on the total space M^{4n} of a vector bundle over $L = G/N(\mathfrak{su}(2))$ [Sw].

There are 3 such classes for $\mathfrak{so}(5)$:

- the minimal $\mathfrak{su}(2)$ with normalizer $SO(4)$, so $L = M = S^4$ and $n = 1$
- our $\mathfrak{so}(3) = \mathfrak{su}(2)$ with $L = SO(5)/SO(2) \times SO(3) \cong Q^3$ and $n = 2$
- the principal $\mathfrak{su}(2)$ with $L = SO(5)/SO(3)$ and $n = 3$.

In the last case, C has a nearly-parallel G_2 structure, so $\mathbb{R}^+ \times C$ has holonomy $Spin(7)$.

Work inside the 7-manifold $\mathcal{C} = \mathbb{R}^+ \times \mathbb{C}\mathbb{P}^3$, with its 3-form φ defining the G_2 metric h .

Let $i: \mathcal{U} \rightarrow \mathcal{C}$ be a 3-dimensional orbit of some subgroup $SU(2)$ or $SO(3)$ of $SO(5)$. Since the latter preserves φ , $i^*\varphi$ must be a constant multiple of the volume form on \mathcal{U} . But $[i^*\varphi] = i^*[-d(R\tau)] = 0$, so $i^*\varphi \equiv 0$.

Lemma. A 4-dimensional submanifold \mathcal{V} of a G_2 manifold is coassociative iff $i^*\varphi \equiv 0$.

We expect each orbit \mathcal{U} to be contained in a unique such a submanifold, $T_x\mathcal{U} \subset T_x\mathcal{V}$.

In favourable circumstances, there will be a foliation of \mathcal{C} by coassociatives of codim one [K, KL]. (\mathcal{C} has a more elementary foliation by Eguchi-Hanson spaces $T^*\mathcal{S}^2$, each with $\mathbf{u} - \mathbf{v}$ constant, but this is not really related to G_2 .)

For the diagonal action of $SO(3)$ on \mathbb{R}^6 , invariant functions are obviously u , v and $\mathbf{u} \cdot \mathbf{v}$. We can recover these on $\mathbb{C}P^3$ from the function $z_0 \bar{z}_3 - \bar{z}_1 z_2 = \sqrt{a/2} e^{it}$ defining Q_+ :


Lemma. $2a = uv - \mathbf{u} \cdot \mathbf{v} = R^2(1 - s^2)$, so $R = \sqrt{\frac{2a}{1 - s^2}}$.

An $SO(3)$ orbit \mathcal{U} will intersect a twistor fibre S^2 of radius R over $p \in S^4 \setminus S^1$ in a circle at 'height' $h = (u - v)/(Rs) \in [-1, 1]$ relative to poles defined by Q_- .

Define $b = u^2 - v^2 = RHs$, so

$$h = (b/a^2) \frac{1 - s^2}{s}.$$

Theorem [ABS]. Setting a, b, t constant defines a coassociative submanifold of \mathcal{C} diffeomorphic to T^*S^2 unless $a = b = 0$.



This is induced from the G_2 structure on \mathcal{C} . Let X be the Killing vector field generating the $SO(2)$ fibres of $\mu: \mathcal{C} \rightarrow \mathbb{R}^6$. Then

$$\sigma = X \lrcorner \varphi = X \lrcorner d(R\tau) = -d(RX \lrcorner \tau).$$

Set $\mathbf{p} = \mathbf{u} + \mathbf{v}$ and $\mathbf{q} = R(\mathbf{u} - \mathbf{v})$, where $R = u + v = |\mathbf{u}| + |\mathbf{v}|$.

Unexplained theorem [ABS]. The components of \mathbf{p}, \mathbf{q} are Darboux coordinates:

$$\sigma = -\frac{1}{2} \sum_{i=1}^3 dp_i \wedge dq_i.$$

Note that σ extends to $\mathbb{R}^3 \cup \mathbb{R}^3$ and is non-degenerate on $\mathbb{R}^6 \setminus \mathbf{0}$. The projections $(\mathbf{u}, \mathbf{v}) \mapsto R^{1/2}\mathbf{u}$ and $(\mathbf{u}, \mathbf{v}) \mapsto R^{1/2}\mathbf{v}$ also have Lagrangian fibres.

Recall that h is the conical metric on \mathcal{C} with holonomy G_2 . We seek the metric g induced on $\mathbb{R}^6 \setminus (\mathbb{R}^3 \cup \mathbb{R}^3)$ by setting

$$h = \mu^*g + \frac{1}{4}N\Theta^2,$$

where $\Theta = 2(X \lrcorner h)/N$ is the connection 1-form, and $N = h(X, X) = 6uv - 2\mathbf{u} \cdot \mathbf{v}$ measures the size of the circle fibres. This makes μ a Riemannian submersion.

Computational theorem [ABS].

$$g = \frac{1}{2}dR^2 + \frac{1}{2}|d\mathbf{u} + d\mathbf{v}|^2 + \frac{2}{N}|u d\mathbf{v} - v d\mathbf{u}|^2 + \frac{1}{2N}\Gamma_+^2 - \frac{1}{4N}\Gamma_-^2,$$

where the 1-forms

$$\Gamma_+ = \mathbf{u} \cdot d\mathbf{v} + \mathbf{v} \cdot d\mathbf{u} - u dv - v du,$$

$$\Gamma_- = \mathbf{u} \cdot d\mathbf{v} - \mathbf{v} \cdot d\mathbf{u} + u dv - v du.$$

vanishes on \mathcal{F}_+ , \mathcal{F}_- respectively.

We can recover the conical nature of g by the change of variables

$$u = R \cos^2(\phi/2), \quad v = R \sin^2(\phi/2).$$

Then $\mathcal{F}_{\pm} \cong \mathbb{R}^+ \times [0, \pi] \times S^2$, with coordinates R, ϕ, σ .

Corollary 1. The restriction of g to \mathcal{F}_{\pm} equals $dR^2 + R^2 \hat{g}$ where

$$\hat{g} = \begin{cases} \frac{1}{2}|d\sigma|^2 + \frac{1}{4}d\phi^2 & \text{on } \mathcal{F}_+ \\ \frac{1}{8}(3 + \cos 2\phi)|d\sigma|^2 + \frac{1}{2}d\phi^2 & \text{on } \mathcal{F}_-. \end{cases}$$

Corollary 2. Relative to g , vectors in the respective singular \mathbb{R}^3 axes meet at an angle of

$$\frac{1}{2}\pi \leq \pi \sqrt{\frac{3}{8} - \frac{1}{8} \cos \theta} \leq \frac{1}{\sqrt{2}}\pi \sim 127^\circ.$$

This is determined by g and the closed 3-form $\psi^+ = X \lrcorner (*\varphi)$.

Proposition.
$$\begin{aligned} \delta uv\psi^+ &= \frac{1}{3}v(2v^2 + 3uv - \mathbf{u} \cdot \mathbf{v})\{d\mathbf{u}, d\mathbf{u}, d\mathbf{u}\} \\ &-v(4u^2 + 3uv + \mathbf{u} \cdot \mathbf{v})\{d\mathbf{v}, d\mathbf{u}, d\mathbf{u}\} + ((u + 2v)\mathbf{v} \cdot d\mathbf{v} + v\mathbf{u} \cdot d\mathbf{v}) \wedge \{\mathbf{u}, d\mathbf{u}, d\mathbf{u}\} \\ &+(v\mathbf{u} \cdot d\mathbf{v} - uv \cdot d\mathbf{v}) \wedge \{\mathbf{v}, d\mathbf{u}, d\mathbf{u}\} + \text{terms interchanging } \mathbf{u} \text{ and } \mathbf{v}. \end{aligned}$$

Let $\mathbf{n} \in S^2$ and set $\mathcal{M}_{\mathbf{n}} = \{(\mathbf{u}, \mathbf{v}) : \mathbf{u} \cdot \mathbf{n} = 0 = \mathbf{v} \cdot \mathbf{n}, uv \neq 0\}$, essentially an \mathbb{R}^4 .

Note that $\mathcal{M}_{\mathbf{n}} = \mathcal{M}_{\mathbf{n}'}$ iff $\mathbf{n} = \pm\mathbf{n}'$, otherwise the intersection is 2-dimensional.

Corollary 3. $\mathcal{M}_{\mathbf{n}}$ is a non-integrable \mathbb{J} -holomorphic subvariety of \mathbb{R}^6 .

An open subset of \mathbb{R}^6 is exhausted by a family of \mathbb{J} -holomorphic surfaces parametrized by $\mathbb{R}\mathbb{P}^2$. The intersection of any two is a 'superminimal' \mathbb{J} -holomorphic curve.

- S.-T. Yau, Simons Foundation lecture, tonight!
- T. Madsen, Multisymplectic Geometry Leuven, in 24 hours.
- K. Dixon, SCSHGAP workshop, in 7 days.

