


Tests of Lorentz and CPT Violation with Neutrino Oscillation Experiments



Teppei Katori
Queen Mary University of London
Cavendish HEP seminar, Univ. Cambridge, Cambridge, UK, Nov. 11, 2014

Tests of Lorentz and CPT Violation with Neutrino Oscillation Experiments



Teppei Katori

Queen Mary University of London

Cavendish HEP seminar, Univ. Cambridge, Cambridge, UK, Nov. 11, 2014

Tests of Lorentz and CPT Violation with Neutrino Oscillation Experiments

outline

1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
4. Lorentz violating neutrino oscillations
5. Test for Lorentz violation with MiniBooNE data
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit analysis
8. Extra-terrestrial neutrinos
9. Conclusion

Teppei Katori

Queen Mary University of London

Cavendish HEP seminar, Univ. Cambridge, Cambridge, UK, Nov. 11, 2014

- 1. Spontaneous Lorentz symmetry breaking**
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
4. Lorentz violating neutrino oscillation
5. Test for Lorentz violation with MiniBooNE data
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos
9. Conclusion

1. Spontaneous Lorentz symmetry breaking (SLSB)

Every fundamental symmetry needs to be tested, including Lorentz symmetry.

After the recognition of theoretical processes that create Lorentz violation, testing Lorentz invariance becomes very exciting

Lorentz and CPT violation has been shown to occur in Planck scale theories, including:

- string theory
- noncommutative field theory
- quantum loop gravity
- extra dimensions
- etc

However, it is very difficult to build a self-consistent theory with Lorentz violation...

1. Spontaneous Lorentz symmetry breaking (SLSB)

Every fundamental symmetry needs to be tested, including Lorentz symmetry.

After the recognition of theoretical processes that create Lorentz violation, testing Lorentz invariance becomes very exciting

Lorentz and CPT violation has been shown to occur in Planck scale theories, including:

- string theory
- noncommutative field theory
- quantum loop gravity
- extra dimensions
- etc

However, it is very difficult to build a self-consistent theory with Lorentz violation...

Spontaneous
Symmetry Breaking
(SSB)!



Y. Nambu
(Nobel prize winner 2008),
picture taken from CPT04 at
Bloomington, IN



1. Spontaneous Lorentz symmetry breaking (SLSB)

$$\text{vacuum Lagrangian for fermion } \mathcal{L} = i\bar{\Psi}\gamma_{\mu}\partial^{\mu}\Psi$$

e.g.) SSB of scalar field in Standard Model (SM)
- If the scalar field has Mexican hat potential

$$L = \frac{1}{2}(\partial_{\mu}\varphi)^2 - \frac{1}{2}\mu^2(\varphi^*\varphi) - \frac{1}{4}\lambda(\varphi^*\varphi)^2$$

$$M(\varphi) = \mu^2 < 0$$





1. Spontaneous Lorentz symmetry breaking (SLSB)

$$\text{vacuum Lagrangian for fermion } \mathcal{L} = i\bar{\Psi}\gamma_{\mu}\partial^{\mu}\Psi - m\bar{\Psi}\Psi$$

e.g.) SSB of scalar field in Standard Model (SM)
- If the scalar field has Mexican hat potential

$$L = \frac{1}{2}(\partial_{\mu}\varphi)^2 - \frac{1}{2}\mu^2(\varphi^*\varphi) - \frac{1}{4}\lambda(\varphi^*\varphi)^2$$

$$M(\varphi) = \mu^2 < 0$$



Particle acquires mass term!

1. Spontaneous Lorentz symmetry breaking (SLSB)

vacuum Lagrangian for fermion $L = i\bar{\Psi}\gamma_\mu\partial^\mu\Psi - m\bar{\Psi}\Psi$

e.g.) SSB of scalar field in Standard Model (SM)

- If the scalar field has Mexican hat potential

$$L = \frac{1}{2}(\partial_\mu\varphi)^2 - \frac{1}{2}\mu^2(\varphi^*\varphi) - \frac{1}{4}\lambda(\varphi^*\varphi)^2$$



$$M(\varphi) = \mu^2 < 0$$

e.g.) SLSB in string field theory

- There are many Lorentz vector fields

- If any of vector field has Mexican hat potential

$$M(a^\mu) = \mu^2 < 0$$



1. Spontaneous Lorentz symmetry breaking (SLSB)

vacuum Lagrangian for fermion $L = i\bar{\Psi}\gamma_\mu\partial^\mu\Psi - m\bar{\Psi}\Psi + \bar{\Psi}\gamma_\mu a^\mu\Psi$

e.g.) SSB of scalar field in Standard Model (SM)

- If the scalar field has Mexican hat potential

$$L = \frac{1}{2}(\partial_\mu\varphi)^2 - \frac{1}{2}\mu^2(\varphi^*\varphi) - \frac{1}{4}\lambda(\varphi^*\varphi)^2$$


$$M(\varphi) = \mu^2 < 0$$

e.g.) SLSB in string field theory


- There are many Lorentz vector fields

- If any of vector field has Mexican hat potential

$$M(a^\mu) = \mu^2 < 0$$



Lorentz symmetry
is spontaneously
broken!



1. Spontaneous Lorentz symmetry breaking

Test of Lorentz violation is to find the coupling of these background fields and ordinary fields (electrons, muons, neutrinos, etc); then **the physical quantities may depend on the rotation of the earth (sidereal time dependence)**.



vacuum Lagrangian for fermion

$$L = i\bar{\Psi}\gamma_{\mu}\partial^{\mu}\Psi - m\bar{\Psi}\Psi + \bar{\Psi}\gamma_{\mu}a^{\mu}\Psi + \bar{\Psi}\gamma_{\mu}c^{\mu\nu}\partial_{\nu}\Psi \dots$$

background fields of the universe



Scientific American (Sept. 2004)



1. Spontaneous Lorentz symmetry breaking

Test of Lorentz violation is to find the coupling of these background fields and ordinary fields (electrons, muons, neutrinos, etc); then **the physical quantities may depend on the rotation of the earth (sidereal time dependence)**.

vacuum Lagrangian for fermion

$$L = i\bar{\Psi}\gamma_{\mu}\partial^{\mu}\Psi - m\bar{\Psi}\Psi + \bar{\Psi}\gamma_{\mu}a^{\mu}\Psi + \bar{\Psi}\gamma_{\mu}c^{\mu\nu}\partial_{\nu}\Psi \dots$$

background fields of the universe

Sidereal time dependence

The smoking gun of Lorentz violation is the **sidereal time dependence** of the observables.

Solar time: 24h 00m 00.0s
sidereal time: 23h 56m 04.1s

Sidereal time dependent physics is often smeared out in solar time distribution
→ Maybe we have some evidence of Lorentz violation but we just didn't notice?!



Target scale

Since it is Planck scale physics, either $>10^{19}\text{GeV}$ or $<10^{-19}\text{GeV}$ is the interesting region.
 $>10^{19}\text{GeV}$ is not possible (LHC is 10^4GeV), but $<10^{-19}\text{GeV}$ is possible.

1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
4. Lorentz violating neutrino oscillation
5. Test for Lorentz violation with MiniBooNE data
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos
9. Conclusion

2. What is Lorentz violation?

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)$$




2. What is Lorentz violation?


$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)$$

hypothetical background vector field

moving particle





Einstein (observer)



2. What is Lorentz violation?

Under the **particle** Lorentz transformation:

$$U \bar{\Psi}(x) \gamma_{\mu} a^{\mu} \Psi(x) U^{-1}$$





2. What is Lorentz violation?

Under the **particle** Lorentz transformation:

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x) \rightarrow U[\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)]U^{-1}$$
$$\neq \bar{\Psi}(\Lambda x)\gamma_{\mu}a^{\mu}\Psi(\Lambda x)$$

Lorentz violation is observable when a particle is moving in the fixed coordinate space




2. What is Lorentz violation?

Under the **particle** Lorentz transformation:

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x) \rightarrow U[\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)]U^{-1}$$



$$\neq \bar{\Psi}(\Lambda x)\gamma_{\mu}a^{\mu}\Psi(\Lambda x)$$

Lorentz violation is observable when a particle is moving in the fixed coordinate space



Under the **observer** Lorentz transformation:

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)$$




2. What is Lorentz violation?

Under the **particle** Lorentz transformation:

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x) \rightarrow U[\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)]U^{-1}$$

$$\neq \bar{\Psi}(\Lambda x)\gamma_{\mu}a^{\mu}\Psi(\Lambda x)$$



Lorentz violation is observable when a particle is moving in the fixed coordinate space



Under the **observer** Lorentz transformation:

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)$$

$$x \rightarrow \Lambda^{-1}x$$




2. What is Lorentz violation?

Under the **particle** Lorentz transformation:

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x) \rightarrow U[\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x)]U^{-1}$$

$$\neq \bar{\Psi}(\Lambda x)\gamma_{\mu}a^{\mu}\Psi(\Lambda x)$$

Lorentz violation is observable when a particle is moving in the fixed coordinate space





Under the **observer** Lorentz transformation:

$$\bar{\Psi}(x)\gamma_{\mu}a^{\mu}\Psi(x) \xrightarrow{\Lambda^{-1}} \bar{\Psi}(\Lambda^{-1}x)\gamma_{\mu}a^{\mu}\Psi(\Lambda^{-1}x)$$

Lorentz violation cannot be generated by observers motion (coordinate transformation is unbroken)

all observers agree for all observations




2. What is CPT violation?


CPT symmetry is the invariance under the CPT transformation

$$L \xrightarrow{\text{CPT}} \Theta L \Theta^{-1} = L' = L, \quad \Theta = \text{CPT}$$


C: charge conjugation



P: parity transformation



T: time reversal





2. What is CPT violation?

CPT phase = $(-1)^n$
 $n = \#$ Lorentz indices

CPT symmetry is the invariance under the CPT transformation

$$L \xrightarrow{\text{CPT}} \Theta L \Theta^{-1} = L' = L, \quad \Theta = \text{CPT}$$

CPT is the perfect symmetry of the Standard Model, due to **CPT theorem**




2. What is CPT violation?

CPT phase = $(-1)^n$
 $n = \# \text{ Lorentz indices}$

CPT symmetry is the invariance under the CPT transformation

$$L \xrightarrow{\text{CPT}} \Theta L \Theta^{-1} = L' = L, \quad \Theta = \text{CPT}$$

CPT is the perfect symmetry of the Standard Model, due to **CPT theorem**





CPT-odd Lorentz violating coefficients (odd number Lorentz indices, e.g., a^{μ} , $g^{\lambda\mu\nu}$)
 CPT-even Lorentz violating coefficients (even number Lorentz indices, e.g., $c^{\mu\nu}$, $\kappa^{\alpha\beta\mu\nu}$)



2. CPT violation implies Lorentz violation

Lorentz invariance \longrightarrow CPT \longrightarrow Lorentz invariance of quantum field theory

CPT violation implies Lorentz violation in interactive quantum field theory.



1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
- 3. Modern test of Lorentz violation**
4. Lorentz violating neutrino oscillation
5. Test for Lorentz violation with MiniBooNE data
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos
9. Conclusion

3. Test of Lorentz violation

Lorentz violation is realized as a coupling of particle fields and background fields, so the basic strategy to find Lorentz violation is:



- (1) choose the coordinate system
- (2) write down the Lagrangian, including Lorentz-violating terms under the formalism
- (3) write down the observables using this Lagrangian

3. Test of Lorentz violation


Lorentz violation is realized as a coupling of particle fields and background fields, so the basic strategy to find Lorentz violation is:

- (1) choose the coordinate system
- (2) write down the Lagrangian, including Lorentz-violating terms under the formalism
- (3) write down the observables using this Lagrangian

- Neutrino beamline is described in Sun-centred coordinates



MiniBooNE beamline



3. Test of Lorentz violation

Lorentz violation is realized as a coupling of particle fields and background fields, so the basic strategy to find Lorentz violation is:

- (1) choose the coordinate system
- (2) write down the Lagrangian, including Lorentz-violating terms under the formalism
- (3) write down the observables using this Lagrangian

Standard Model Extension (SME) is the standard formalism for the general search for Lorentz violation. SME is a minimum extension of QFT with Particle Lorentz violation

SME Lagrangian in neutrino sector

$$L = \frac{1}{2} i \bar{\psi}_A \Gamma_{AB}^{\nu} \partial_{\nu} \psi_B - M_{AB} \bar{\psi}_A \psi_B + h.c.$$

SME coefficients

$$\Gamma_{AB}^{\nu} = \gamma^{\nu} \delta_{AB} + c_{AB}^{\mu\nu} \gamma_{\mu} + d_{AB}^{\mu\nu} \gamma_{\mu} \gamma_5 + e_{AB}^{\nu} + i f_{AB}^{\nu} \gamma_5 + \frac{1}{2} g_{AB}^{\lambda\mu\nu} \sigma_{\lambda\mu} \dots$$

$$M_{AB} = m_{AB} + i m_{5AB} \gamma_5 + a_{AB}^{\mu} \gamma_{\mu} + b_{AB}^{\mu} \gamma_5 \gamma_{\mu} + \frac{1}{2} H_{AB}^{\mu\nu} \sigma_{\mu\nu} \dots$$

3. Test of Lorentz violation

Lorentz violation is realized as a coupling of particle fields and background fields, so the basic strategy to find Lorentz violation is:

- (1) choose the coordinate system
- (2) write down the Lagrangian, including Lorentz-violating terms under the formalism
- (3) write down the observables using this Lagrangian

Standard Model Extension (SME) is the standard formalism for the general search for Lorentz violation. SME is a minimum extension of QFT with Particle Lorentz violation

SME Lagrangian in neutrino sector

$$L = \frac{1}{2} i \bar{\psi}_A \Gamma_{AB}^\nu \partial_\nu \psi_B - M_{AB} \bar{\psi}_A \psi_B + h.c.$$

SME coefficients

$$\Gamma_{AB}^\nu = \gamma^\nu \delta_{AB} + c_{AB}^{\mu\nu} \gamma_\mu + d_{AB}^{\mu\nu} \gamma_\mu \gamma_5 + e_{AB}^\nu + i f_{AB}^\nu \gamma_5 + \frac{1}{2} g_{AB}^{\lambda\mu\nu} \sigma_{\lambda\mu} \dots$$

$$M_{AB} = m_{AB} + i m_{5AB} \gamma_5 + a_{AB}^\mu \gamma_\mu + b_{AB}^\mu \gamma_5 \gamma_\mu + \frac{1}{2} H_{AB}^{\mu\nu} \sigma_{\mu\nu} \dots$$

CPT odd

CPT even

3. Test of Lorentz violation

Lorentz violation is realized as a coupling of particle fields and background fields, so the basic strategy to find Lorentz violation is:

- (1) choose the coordinate system
- (2) write down the Lagrangian, including Lorentz-violating terms under the formalism
- (3) write down the observables using this Lagrangian

Various physics are predicted under SME, but among them, the smoking gun of Lorentz violation is the **sidereal time dependence** of the observables

solar time: 24h 00m 00.0s
sidereal time: 23h 56m 04.1s

$$\begin{array}{l} \text{sidereal frequency } \omega_{\oplus} = \frac{2\pi}{23h56m4.1s} \\ \text{sidereal time } T_{\oplus} \end{array}$$

Lorentz-violating neutrino oscillation probability for short-baseline experiments

$$P_{\nu_{\mu} \rightarrow \nu_e} = \left(\frac{L}{\hbar c} \right)^2 \left| (C)_{e\mu} + (A_s)_{e\mu} \sin \omega_{\oplus} T_{\oplus} + (A_c)_{e\mu} \cos \omega_{\oplus} T_{\oplus} + (B_s)_{e\mu} \sin 2\omega_{\oplus} T_{\oplus} + (B_c)_{e\mu} \cos 2\omega_{\oplus} T_{\oplus} \right|^2$$

3. Test of Lorentz violation

Lorentz violation is realized as a coupling of particle fields and background fields, so the basic strategy to find Lorentz violation is:

- (1) choose the coordinate system
- (2) write down the Lagrangian, including Lorentz-violating terms under the formalism
- (3) write down the observables using this Lagrangian

Various physics are predicted under SME, but among them, the smoking gun of Lorentz violation is the **sidereal time dependence** of the observables

solar time: 24h 00m 00.0s
sidereal time: 23h 56m 04.1s

$$\text{sidereal frequency } \omega_{\oplus} = \frac{2\pi}{23h56m4.1s}$$

$$\text{sidereal time } T_{\oplus}$$

Lorentz-violating neutrino oscillation probability for short-baseline experiments

$$P_{\nu_{\mu} \rightarrow \nu_e} = \left(\frac{L}{\hbar c} \right)^2 \left| \underbrace{(C)_{e\mu}}_{\text{time independent amplitude}} + \underbrace{(A_s)_{e\mu}}_{\text{sidereal time dependent amplitude}} \sin \omega_{\oplus} T_{\oplus} + \underbrace{(A_c)_{e\mu}}_{\text{sidereal time dependent amplitude}} \cos \omega_{\oplus} T_{\oplus} + \underbrace{(B_s)_{e\mu}}_{\text{sidereal time dependent amplitude}} \sin 2\omega_{\oplus} T_{\oplus} + \underbrace{(B_c)_{e\mu}}_{\text{sidereal time dependent amplitude}} \cos 2\omega_{\oplus} T_{\oplus} \right|^2$$


Sidereal variation analysis for short baseline neutrino oscillation is 5-parameter fitting problem

3. Modern tests of Lorentz violation

The latest meeting was in June 2013

<http://www.physics.indiana.edu/~kostelec/faq.html>

CPT'13



MEETING LINKS

[Meeting Home](#)
[Registration](#)
[Program](#)
[Proceedings](#)
[Travel](#)
[Accommodations](#)

LOCAL LINKS

[IUCSS](#)
[IU Physics](#)
[IU Astronomy](#)
[IU Bloomington](#)
[Bloomington area](#)

Sixth Meeting on **CPT AND LORENTZ SYMMETRY**

June 17-21, 2013

Indiana University, Bloomington

The *Sixth Meeting on CPT and Lorentz Symmetry* will be held in the [Physics Department, Indiana University](#) in [Bloomington](#), Indiana, U.S.A. on June 17-21, 2013. The meeting will focus on tests of these fundamental symmetries and on related theoretical issues, including scenarios for possible violations.

Topics include:


- experimental and observational searches for CPT and Lorentz violation involving
 - accelerator and collider experiments
 - atomic, nuclear, and particle decays
 - birefringence, dispersion, and anisotropy in cosmological sources
 - clock-comparison measurements
 - CMB polarization
 - electromagnetic resonant cavities and lasers
 - tests of the equivalence principle
 - gauge and Higgs particles
 - high-energy astrophysical observations
 - laboratory and gravimetric tests of gravity
 - matter interferometry
 - neutrino oscillations and propagation, neutrino-antineutrino mixing

Atomic Interferometer
 $(a,c)^{n,p,e} < 10^{-6}$



PRL106(2011)151102

CERN Antiproton Decelerator
 $(M_p - \bar{M}_p)/M_p < 10^{-8}$



Nature419(2002)456

Spin torsion pendulum
 $b_e < 10^{-30}$ GeV



PRL97(2006)

Tevatron and LEP
 $-5.8 \times 10^{-12} < \kappa_{tr} - 4/3 c_e^{00} < 1.2 \times 10^{-11}$



PRL102(2009)170402

KTev/KLOE (strange)
 $\Delta a_K < 10^{-22}$ GeV
 FOCUS (charm)
 $\Delta a_D < 10^{-16}$ GeV
 BaBar/Belle (bottom)
 $\Delta m_B/m_B < 10^{-14}$

- ★ clock-comparison measurements
- ★ CMB polarization
- ★ electrodynamic constants
- ★ tests of gauge invariance
- ★ high precision laboratory measurements
- ★ matter interferometry
- ★ neutrino oscillations and propagation, neutrino-antineutrino oscillations and decays of K, B, D mesons
- ★ particle-antiparticle comparisons

Limits from all these experiments >50 page tables!
 Rev.Mod.Phys.83(2011)11
 ArXiv:0801.0287v6

Neutrino TOF
 $(v-c)/c < 10^{-5}$



PRD76(2007)072005
 JHEP11(2012)049


vacuum birefringence
 $\kappa_{e+}, \kappa_{o-} < 10^{-37}$



PRD76(2006)140401

Test of Lorentz invariance with neutrino oscillation is very interesting, because neutrinos are the least known standard model particles!

stable gas maser (station) < 10⁻³³ GeV
 (boost) < 10⁻²⁷ GeV



<p>LSND</p> <p>PRD72(2005)076004</p>	<p>MINOS ND</p> <p>PRL101(2008)151601</p>	<p>MINOS FD</p> <p>PRL105(2010)151601</p>	<p>IceCube</p> <p>PRD82(2010)112003</p>	<p>MiniBooNE</p> <p>PLB718(2013)1303</p>	<p>Double Chooz</p> <p>PRD86(2013)112009</p>	<p>Super-Kamiokande</p> <p>ArXiv:1410.4267</p>
--------------------------------------	---	---	---	--	--	--

PLB556(2003)7
 PRL100(2008)131802

classical and quantum field theory
 mathematical foundations, Finn Toppe Katori
 PRL99(2007)050401


uar
 PRL105(2010)151604
 PRL107(2010)171604

1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
- 4. Lorentz violating neutrino oscillation**
5. Test for Lorentz violation with MiniBooNE data
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos
9. Conclusion

4. Neutrinos


Neutrinos in the standard model

The standard model describes 6 quarks and 6 leptons and 3 types of force carriers.



Neutrinos are special because,


1. they only interact with weak nuclear force.



2. interaction eigenstate is not Hamiltonian eigenstate (propagation eigenstate). Thus propagation of neutrinos changes their species, called **neutrino oscillation**.

4. Neutrino oscillations, natural interferometers


Neutrino oscillation is an interference experiment (cf. double slit experiment)



For double slit experiment, if path ν_1 and path ν_2 have different length, they have different phase rotations and it causes interference.

4. Neutrino oscillations, natural interferometers


Neutrino oscillation is an interference experiment (cf. double slit experiment)



If 2 neutrino Hamiltonian eigenstates, ν_1 and ν_2 , have different phase rotation, they cause quantum interference.

4. Neutrino oscillations, natural interferometers

Neutrino oscillation is an interference experiment (cf. double slit experiment)




If 2 neutrino Hamiltonian eigenstates, ν_1 and ν_2 , have different phase rotation, they cause quantum interference.

If ν_1 and ν_2 , have different mass, they have different velocity, so thus different phase rotation.

4. Neutrino oscillations, natural interferometers

Neutrino oscillation is an interference experiment (cf. double slit experiment)




If 2 neutrino Hamiltonian eigenstates, ν_1 and ν_2 , have different phase rotation, they cause quantum interference.

If ν_1 and ν_2 , have different mass, they have different velocity, so thus different phase rotation.

The detection may be different flavor (neutrino oscillations).

4. Lorentz violation with neutrino oscillation


Neutrino oscillation is an interference experiment (cf. double slit experiment)



If 2 neutrino Hamiltonian eigenstates, ν_1 and ν_2 , have different phase rotation, they cause quantum interference.

4. Lorentz violation with neutrino oscillation

Neutrino oscillation is an interference experiment (cf. double slit experiment)




If 2 neutrino Hamiltonian eigenstates, ν_1 and ν_2 , have different phase rotation, they cause quantum interference.

If ν_1 and ν_2 , have different coupling with Lorentz violating field, neutrinos also oscillate. The sensitivity of neutrino oscillation is comparable the target scale of Lorentz violation ($<10^{-19}\text{GeV}$).

4. Lorentz violation with neutrino oscillation

Neutrino oscillation is an interference experiment (cf. double slit experiment)



If 2 neutrino Hamiltonian eigenstates, ν_1 and ν_2 , have different phase rotation, they cause quantum interference.

If ν_1 and ν_2 , have different coupling with Lorentz violating field, neutrinos also oscillate. The sensitivity of neutrino oscillation is comparable the target scale of Lorentz violation ($<10^{-19}\text{GeV}$).

If neutrino oscillation is caused by Lorentz violation, interference pattern (oscillation probability) may have sidereal time dependence.

1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
4. Lorentz violating neutrino oscillation
- 5. Test for Lorentz violation with MiniBooNE data**
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos
9. Conclusion

5. MiniBooNE experiment

MiniBooNE is a short-baseline neutrino oscillation experiment at Fermilab.

$$\nu_{\mu} \xrightarrow{\text{oscillation}} \nu_e + n \rightarrow e^{-} + p$$


$$\bar{\nu}_{\mu} \xrightarrow{\text{oscillation}} \bar{\nu}_e + p \rightarrow e^{+} + n$$

Booster Neutrino Beamline (BNB) creates $\sim 800(600)$ MeV neutrino(anti-neutrino) by pion decay-in-flight. Cherenkov radiation from the charged leptons are observed by MiniBooNE Cherenkov detector to reconstruct neutrino energy.

FNAL Booster




Magnetic focusing horn



$\sim 520\text{m}$
→

MiniBooNE detector



primary beam
→
(8 GeV protons)

secondary beam
→
(1-2 GeV pions)

tertiary beam
→
(800 MeV ν_{μ} , 600 MeV anti- ν_{μ})



1280 of 8" PMT

5. MiniBooNE experiment

MiniBooNE is a short-baseline neutrino oscillation experiment at Fermilab.


$$\nu_{\mu} \xrightarrow{\text{oscillation}} \nu_e + n \rightarrow e^{-} + p$$

$$\bar{\nu}_{\mu} \xrightarrow{\text{oscillation}} \bar{\nu}_e + p \rightarrow e^{+} + n$$


Neutrino mode analysis: MiniBooNE saw the 3.0σ excess at **low energy region**

Antineutrino mode analysis: MiniBooNE saw the 1.4σ excess at **low and high energy region**

MiniBooNE low E ν_e excess



MiniBooNE anti- ν_e excess



These excesses are not predicted by neutrino Standard Model (ν SM), therefore it might sterile neutrino or other new physics, such as Lorentz violation

→ Oscillation candidate events may have sidereal time dependence!

5. Lorentz violation with MiniBooNE neutrino data

MiniBooNE is a short-baseline neutrino oscillation experiment at Fermilab.


$$\nu_{\mu} \xrightarrow{\text{oscillation}} \nu_e + n \rightarrow e^{-} + p$$

$$\bar{\nu}_{\mu} \xrightarrow{\text{oscillation}} \bar{\nu}_e + p \rightarrow e^{+} + n$$

Neutrino mode analysis: MiniBooNE saw the 3.0σ excess at low energy region

Antineutrino mode analysis: MiniBooNE saw the 1.4σ excess at low and high energy region

Electron neutrino candidate data prefer **sidereal time independent solution (flat)**



5. Lorentz violation with MiniBooNE neutrino data

MiniBooNE is a short-baseline neutrino oscillation experiment at Fermilab.


$$\nu_{\mu} \xrightarrow{\text{oscillation}} \nu_e + n \rightarrow e^{-} + p$$

$$\bar{\nu}_{\mu} \xrightarrow{\text{oscillation}} \bar{\nu}_e + p \rightarrow e^{+} + n$$


Neutrino mode analysis: MiniBooNE saw the 3.0σ excess at low energy region

Antineutrino mode analysis: MiniBooNE saw the 1.4σ excess at low and high energy region

Electron neutrino candidate data prefer **sidereal time independent solution (flat)**



Electron antineutrino candidate data prefer **sidereal time dependent solution**, however statistical significance is marginal



We find no evidence of Lorentz violation

5. Lorentz violation with MiniBooNE neutrino data

MiniBooNE is a short-baseline neutrino oscillation experiment at Fermilab.

$$\nu_{\mu} \xrightarrow{\text{oscillation}} \nu_e + n \rightarrow e^{-} + p$$

$$\bar{\nu}_{\mu} \xrightarrow{\text{oscillation}} \bar{\nu}_e + p \rightarrow e^{+} + n$$

Neutrino mode analysis: MiniBooNE saw the 3.0σ excess at low energy region

Antineutrino mode analysis: MiniBooNE saw the 1.4σ excess at low and high energy region

Since we find no evidence of Lorentz violation, we set limits on the combination SME coefficients.

	ν -mode BF	2σ limit	$\bar{\nu}$ -mode BF	2σ limit
$ (C)_{e\mu} $	$3.1 \pm 0.6 \pm 0.9$	< 4.2	$0.1 \pm 0.8 \pm 0.1$	< 2.6
$ (\mathcal{A}_S)_{e\mu} $	$0.6 \pm 0.9 \pm 0.3$	< 3.3	$2.4 \pm 1.3 \pm 0.5$	< 3.9
$ (\mathcal{A}_C)_{e\mu} $	$0.4 \pm 0.9 \pm 0.4$	< 4.0	$2.1 \pm 1.2 \pm 0.4$	< 3.7



SME coefficients combination (unit 10^{-20} GeV)

$ (C)_{e\mu} $	$\pm[(a_L)_{e\mu}^T + 0.75(a_L)_{e\mu}^Z] - \langle E \rangle [1.22(c_L)_{e\mu}^{TT} + 1.50(c_L)_{e\mu}^{TZ} + 0.34(c_L)_{e\mu}^{ZZ}]$
$ (\mathcal{A}_S)_{e\mu} $	$\pm[0.66(a_L)_{e\mu}^Y] - \langle E \rangle [1.33(c_L)_{e\mu}^{TY} + 0.99(c_L)_{e\mu}^{YZ}]$
$ (\mathcal{A}_C)_{e\mu} $	$\pm[0.66(a_L)_{e\mu}^X] - \langle E \rangle [1.33(c_L)_{e\mu}^{TX} + 0.99(c_L)_{e\mu}^{XZ}]$

5. Summary of results

LSND experiment



LSND is a short-baseline neutrino oscillation experiment at Los Alamos.



LSND saw the 3.8σ excess of electron antineutrinos from muon antineutrino beam; since this excess is not understood by neutrino Standard Model, it might be new physics

Data is consistent with flat solution, but sidereal time solution is not excluded.

LSND oscillation candidate sidereal time distribution



$\sim 10^{-19}$ GeV CPT-odd or $\sim 10^{-17}$ GeV CPT-even Lorentz violation could be the solution of LSND excess

5. Summary of results

Since we find no evidence of Lorentz violation from MiniBooNE analysis, we set limits on the SME coefficients.

These limits exclude SME values to explain LSND data, **therefore there is no simple Lorentz violation motivated scenario to accommodate LSND and MiniBooNE results simultaneously**

Coefficient	$e\mu$ (ν mode low energy region)	$e\mu$ ($\bar{\nu}$ mode combined region)
$\text{Re}(a_L)^T$ or $\text{Im}(a_L)^T$	4.2×10^{-20} GeV	2.6×10^{-20} GeV
$\text{Re}(a_L)^X$ or $\text{Im}(a_L)^X$	6.0×10^{-20} GeV	5.6×10^{-20} GeV
$\text{Re}(a_L)^Y$ or $\text{Im}(a_L)^Y$	5.0×10^{-20} GeV	5.9×10^{-20} GeV
$\text{Re}(a_L)^Z$ or $\text{Im}(a_L)^Z$	5.6×10^{-20} GeV	3.5×10^{-20} GeV
$\text{Re}(c_L)^{XY}$ or $\text{Im}(c_L)^{XY}$	—	—
$\text{Re}(c_L)^{XZ}$ or $\text{Im}(c_L)^{XZ}$	1.1×10^{-19}	6.2×10^{-20}
$\text{Re}(c_L)^{YZ}$ or $\text{Im}(c_L)^{YZ}$	9.2×10^{-20}	6.5×10^{-20}
$\text{Re}(c_L)^{XX}$ or $\text{Im}(c_L)^{XX}$	—	—
$\text{Re}(c_L)^{YY}$ or $\text{Im}(c_L)^{YY}$	—	—
$\text{Re}(c_L)^{ZZ}$ or $\text{Im}(c_L)^{ZZ}$	3.4×10^{-19}	1.3×10^{-19}
$\text{Re}(c_L)^{TT}$ or $\text{Im}(c_L)^{TT}$	9.6×10^{-20}	3.6×10^{-20}
$\text{Re}(c_L)^{TX}$ or $\text{Im}(c_L)^{TX}$	8.4×10^{-20}	4.6×10^{-20}
$\text{Re}(c_L)^{TY}$ or $\text{Im}(c_L)^{TY}$	6.9×10^{-20}	4.9×10^{-20}
$\text{Re}(c_L)^{TZ}$ or $\text{Im}(c_L)^{TZ}$	7.8×10^{-20}	2.9×10^{-20}




1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
4. Lorentz violating neutrino oscillation
5. Test for Lorentz violation with MiniBooNE data
- 6. Test for Lorentz violation with Double Chooz data**
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos
9. Conclusion

6. Double Chooz experiment

Reactor electron antineutrino disappearance

- Double Chooz, DayaBay and RENO experiments observed disappearance signals


Double Chooz reactor neutrino candidate




6. Double Chooz experiment

Reactor electron antineutrino disappearance


- Double Chooz, DayaBay and RENO experiments observed disappearance



Double Chooz reactor neutrino candidate



The Big Bang Theory (CBS)




6. Double Chooz experiment



Reactor electron antineutrino disappearance

- Double Chooz, DayaBay and RENO experiments observed disappearance

This small disappearance may have sidereal time dependence?



Double Chooz reactor neutrino candidate




Leonard: What do you think about the latest Double Chooz result?
 Sheldon: I think this is Lorentz violation..., check sidereal time dependence

6. Double Chooz experiment

So far, we have set limits on


- 1. $\nu_e \leftrightarrow \nu_\mu$ channel: LSND, MiniBooNE, MINOS ($<10^{-20}$ GeV)
- 2. $\nu_\mu \leftrightarrow \nu_\tau$ channel: MINOS, IceCube ($<10^{-23}$ GeV)

The last untested channel is $\nu_e \leftrightarrow \nu_\tau$



It is possible to limit $\nu_e \leftrightarrow \nu_\tau$ channel from reactor ν_e disappearance experiment

$$P(\nu_e \leftrightarrow \nu_e) = 1 - P(\nu_e \leftrightarrow \nu_\mu) - P(\nu_e \leftrightarrow \nu_\tau) \sim 1 - P(\nu_e \leftrightarrow \nu_\tau)$$



The Big Bang Theory (CBS)

Leonard: What do you think about the latest Double Chooz result?
Sheldon: I think this is Lorentz violation..., check sidereal time dependence

6. Double Chooz experiment

So far, we have set limits on

1. $\nu_e \leftrightarrow \nu_\mu$ channel: LSND, MiniBooNE, MINOS ($<10^{-20}$ GeV)
2. $\nu_\mu \leftrightarrow \nu_\tau$ channel: MINOS, IceCube ($<10^{-23}$ GeV)


The last untested channel is $\nu_e \leftrightarrow \nu_\tau$

It is possible to limit $\nu_e \leftrightarrow \nu_\tau$ channel from reactor ν_e disappearance experiment

$$P(\nu_e \leftrightarrow \nu_e) = 1 - P(\nu_e \leftrightarrow \nu_\mu) - P(\nu_e \leftrightarrow \nu_\tau) \sim 1 - P(\nu_e \leftrightarrow \nu_\tau)$$

Small disappearance signal
prefers **sidereal time independent**
solution (flat)

We set limits in the e- τ sector for
the first time; $\nu_e \leftrightarrow \nu_\tau$ ($<10^{-20}$ GeV)



6. Double Chooz experiment

By this work, Lorentz violation is tested with all neutrino channels

Chance to see the Lorentz violation in terrestrial neutrino experiments will be very small

MiniBooNE
MINOS ND

Double Chooz

IceCube
MINOS FD

$d = 3$	Coefficient	$e\mu$	$e\tau$	$\mu\tau$
	$\text{Re}(a_L)^T$	10^{-20} GeV	10^{-19} GeV	–
	$\text{Re}(a_L)^X$	10^{-20} GeV	10^{-19} GeV	10^{-23} GeV
	$\text{Re}(a_L)^Y$	10^{-21} GeV	10^{-19} GeV	10^{-23} GeV
	$\text{Re}(a_L)^Z$	10^{-19} GeV	10^{-19} GeV	–
$d = 4$	Coefficient	$e\mu$	$e\tau$	$\mu\tau$
	$\text{Re}(c_L)^{XY}$	10^{-21}	10^{-17}	10^{-23}
	$\text{Re}(c_L)^{XZ}$	10^{-21}	10^{-17}	10^{-23}
	$\text{Re}(c_L)^{YZ}$	10^{-21}	10^{-16}	10^{-23}
	$\text{Re}(c_L)^{XX}$	10^{-21}	10^{-16}	10^{-23}
	$\text{Re}(c_L)^{YY}$	10^{-21}	10^{-16}	10^{-23}
	$\text{Re}(c_L)^{ZZ}$	10^{-19}	10^{-16}	–
	$\text{Re}(c_L)^{TT}$	10^{-19}	10^{-17}	–
	$\text{Re}(c_L)^{TX}$	10^{-22}	10^{-17}	10^{-27}
	$\text{Re}(c_L)^{TY}$	10^{-22}	10^{-17}	10^{-27}
	$\text{Re}(c_L)^{TZ}$	10^{-20}	10^{-16}	–

Recently, Super-Kamiokande collaboration published significantly better limits
 arXiv:1410.4267

1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
4. Lorentz violating neutrino oscillation
5. Test for Lorentz violation with MiniBooNE data
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos
9. Conclusion

7. Massive Lorentz-violating model

Double Chooz oscillation signal has no sidereal time dependence.

By assuming main source of neutrino oscillation is neutrino mass, we can study perturbation terms to find secondary effect to cause oscillations.

massive neutrino oscillation Lorentz violating neutrino oscillation

$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) = P^0(\bar{\nu}_e \rightarrow \bar{\nu}_e) + P^1(\bar{\nu}_e \rightarrow \bar{\nu}_e) + P^2(\bar{\nu}_e \rightarrow \bar{\nu}_e) + \dots$$

In this way, we can access to different types of Lorentz violation

7. Anomalous energy spectrum

Sidereal variation is one of many predicted phenomena of Lorentz violating neutrino oscillations.

Lorentz violation predicts unexpected energy dependence of neutrino oscillations from standard neutrino mass oscillations.

Effective Hamiltonian for neutrino oscillation

massive neutrino oscillation

Lorentz violating neutrino oscillation

$$h_{eff} = \frac{m^2}{2E} + a + cE + \dots$$

This is very useful to differentiate 2 effects:

- massive neutrino oscillation
- sidereal time independent Lorentz violating neutrino oscillation


Double Chooz released its energy spectrum (with full error matrix). We use this to test time independent Lorentz violating neutrino oscillation.

7. Double Chooz spectrum fit

Neutrino-Antineutrino oscillation

- Most of neutrino-neutrino oscillation channels are constraint from past analyses
- Here, we focus to test neutrino-antineutrino oscillation

ex) anti- $\nu_e \rightarrow \nu_e$ oscillation fit with Double Chooz data



These fits provide first limits on neutrino-antineutrino time independent Lorentz violating coefficients

1. Spontaneous Lorentz symmetry breaking
2. What is Lorentz and CPT violation?
3. Modern test of Lorentz violation
4. Lorentz violating neutrino oscillation
5. Test for Lorentz violation with MiniBooNE data
6. Test for Lorentz violation with Double Chooz data
7. Double Chooz spectrum fit
8. Extra-terrestrial neutrinos

9. Conclusion


8. Neutrino standard Model (ν SM)

This is the world data of neutrino oscillation

It looks majority of region is either accepted (positive signals) or excluded


But this is model dependent diagram, because it assumes **neutrino mass as phase**, and **mass mixing matrix elements as amplitude of neutrino oscillations**

What is model independent diagram look like?




8. Lorentz violation with neutrino oscillation

Model independent neutrino oscillation data is the function of neutrino energy and baseline.




8. Lorentz violation with neutrino oscillation

Model independent neutrino oscillation data is the function of neutrino energy and baseline.




8. Lorentz violation with neutrino oscillation

Model independent neutrino oscillation data is the function of neutrino energy and baseline.



8. Lorentz violation with neutrino oscillation

Model independent neutrino oscillation data is the function of neutrino energy and baseline.




8. Lorentz violation with neutrino oscillation

extra galactic
neutrino potential

→
1Mpc(~Andromeda)

?
?





8. Lorentz violation with neutrino oscillation

extra galactic neutrino potential

→
1Mpc(~Andromeda)

?
?

IceCube collaboration
PRL111(2013)021103




TeV neutrino potential

↑
1TeV



8. Lorentz violation with extra-terrestrial neutrinos

Combination of longer baseline and higher energy makes extra-terrestrial neutrino to be the most sensitive source of fundamental physics.



cf) The most sensitive test of Planck scale physics is vacuum birefringence of polarized GRB

Conclusion

Lorentz and CPT violation has been shown to occur in Planck-scale theories.

There is a world wide effort to test Lorentz violation with various state-of-the-art technologies.

LSND and MiniBooNE data suggest Lorentz violation is an interesting solution to neutrino oscillation.

MiniBooNE sets limits on Lorentz violation on $\nu_{\mu} \rightarrow \nu_e$ oscillation coefficients. These limits together with MINOS exclude simple Lorentz violation motivated scenario to explain LSND anomaly.

MiniBooNE, LSND, MINOS, IceCube, and Double Chooz, Super-Kamiokande set stringent limits on Lorentz violation in neutrino sector in terrestrial level.

Extra-terrestrial neutrinos from IceCube are one of the most sensitive tool to test fundamental physics, such as Lorentz violation.

Thank you for your attention!

backup

2. Comment: Is there preferred frame?

As we see, all observers are related with observer's Lorentz transformation, so there is no special "preferred" frame (all observer's are consistent)

But there is a frame where universe looks isotropic even with a Lorentz violating vector field. You may call that is the "preferred frame", and people often speculate the frame where CMB looks isotropic is such a frame (called "CMB frame").

However, we are not on CMB frame (e.g., dipole term of WMAP is nonzero), so we expect anisotropy by lab experiments even CMB frame is the preferred frame.

5. MiniBooNE experiment

MiniBooNE is a short-baseline neutrino oscillation experiment at Fermilab.


$$\nu_{\mu} \xrightarrow{\text{oscillation}} \nu_e + n \rightarrow e^{-} + p$$

$$\bar{\nu}_{\mu} \xrightarrow{\text{oscillation}} \bar{\nu}_e + p \rightarrow e^{+} + n$$


Neutrino mode analysis: MiniBooNE saw the 3.0σ excess at **low energy region**

Antineutrino mode analysis: MiniBooNE saw the 1.4σ excess at **low and high energy region**

MiniBooNE low E ν_e excess



MiniBooNE anti- ν_e excess



Intrinsic background errors are constraint from MiniBooNE data

Data driven corrections are applied to MisID backgrounds

5. MiniBooNE experiment

MiniBooNE is a short-baseline neutrino oscillation experiment at Fermilab.


$$\nu_{\mu} \xrightarrow{\text{oscillation}} \nu_e + n \rightarrow e^{-} + p$$

$$\bar{\nu}_{\mu} \xrightarrow{\text{oscillation}} \bar{\nu}_e + p \rightarrow e^{+} + n$$


Neutrino mode analysis: MiniBooNE saw the 3.0σ excess at **low energy region**

Antineutrino mode analysis: MiniBooNE saw the 1.4σ excess at **low and high energy region**

MiniBooNE low E ν_e excess



MiniBooNE anti- ν_e excess



These excesses are not predicted by neutrino Standard Model (ν SM), therefore it might sterile neutrino or other new physics, such as Lorentz violation

→ Oscillation candidate events may have sidereal time dependence!



6. Lorentz violation with MiniBooNE

Sidereal variation of neutrino oscillation probability for MiniBooNE (5 parameters)

$$P_{\nu_e \rightarrow \nu_\mu} = \left(\frac{L}{\hbar c} \right)^2 \left| (C)_{e\mu} + (A_s)_{e\mu} \sin w_{\oplus \oplus} T + (A_c)_{e\mu} \cos w_{\oplus \oplus} T + (B_s)_{e\mu} \sin 2w_{\oplus \oplus} T + (B_c)_{e\mu} \cos 2w_{\oplus \oplus} T \right|^2$$

Expression of 5 observables (14 SME parameters)

$$(C)_{e\mu} = (a_L)_{e\mu}^T - N^Z (a_L)_{e\mu}^Z + E \left[-\frac{1}{2} (3 - N^Z N^Z) (c_L)_{e\mu}^{TT} + 2N^Z (c_L)_{e\mu}^{TZ} + \frac{1}{2} (1 - 3N^Z N^Z) (c_L)_{e\mu}^{ZZ} \right]$$

$$(A_s)_{e\mu} = N^Y (a_L)_{e\mu}^X - N^X (a_L)_{e\mu}^Y + E \left[-2N^Y (c_L)_{e\mu}^{TX} + 2N^X (c_L)_{e\mu}^{TY} + 2N^Y N^Z (c_L)_{e\mu}^{XZ} - 2N^X N^Z (c_L)_{e\mu}^{YZ} \right]$$

$$(A_c)_{e\mu} = -N^X (a_L)_{e\mu}^X - N^Y (a_L)_{e\mu}^Y + E \left[2N^X (c_L)_{e\mu}^{TX} + 2N^Y (c_L)_{e\mu}^{TY} - 2N^X N^Z (c_L)_{e\mu}^{XZ} - 2N^Y N^Z (c_L)_{e\mu}^{YZ} \right]$$

$$(B_s)_{e\mu} = E \left[N^X N^Y \left((c_L)_{e\mu}^{XX} - (c_L)_{e\mu}^{YY} \right) - (N^X N^X - N^Y N^Y) (c_L)_{e\mu}^{XY} \right]$$

$$(B_c)_{e\mu} = E \left[-\frac{1}{2} (N^X N^X - N^Y N^Y) \left((c_L)_{e\mu}^{XX} - (c_L)_{e\mu}^{YY} \right) - 2N^X N^Y (c_L)_{e\mu}^{XY} \right]$$

$$\begin{pmatrix} N^X \\ N^Y \\ N^Z \end{pmatrix} = \begin{pmatrix} \cos \chi \sin \theta \cos \phi - \sin \chi \cos \theta \\ \sin \theta \sin \phi \\ -\sin \chi \sin \theta \cos \phi - \cos \chi \cos \theta \end{pmatrix}$$

coordinate dependent direction vector
(depends on the latitude of FNAL, location
of BNB and MiniBooNE detector)

8. Superluminal neutrinos

OPERA

$$\begin{aligned}v(\text{neutrino}) &= c + (2.37 \pm 0.32) \times 10^{-5} c \\ &= c + (16 \pm 2) \times 10^3 \text{ mph}\end{aligned}$$

It is fascinating result, but...

- time of flight is kinematic test (less sensitive than neutrino oscillations)
- no indication of Lorentz violation from any neutrino oscillation experiments
- superluminal neutrino is unstable (vacuum Cherenkov radiation) [ArXiv:1109.6562](#)
- pion phase space is limited to create such neutrinos [ArXiv:1109.6630](#)
- SN1987A neutrinos provide severe limit to superluminal neutrinos [PRL58\(1987\)1490](#)
- etc...

It is very difficult to interpret superluminal neutrinos at OPERA by Lorentz violation within field theory approach.